

Acta Crystallographica Section E

Structure Reports

Online

ISSN 1600-5368

3-Hexadecyl-1,5-dimethyl-1*H*-1,5-benzodiazepine-2,4(3*H*,5*H*)-dioneRachida Dardouri,^a Youssef Kandri Rodi,^a Sonia Ladeira,^b El Mokhtar Essassi^a and Seik Weng Ng^{c*}

^aLaboratoire de Chimie Organique Hétérocyclique, Pôle de Compétences Pharmacochimie, Université Mohammed V-Agdal, BP 1014 Avenue Ibn Batout, Rabat, Morocco, ^bService Commun Rayons-X, Laboratoire de Chimie de Coordination, Toulouse, France, and ^cDepartment of Chemistry, University of Malaya, 50603 Kuala Lumpur, Malaysia
Correspondence e-mail: seikweng@um.edu.my


Received 6 April 2011; accepted 11 April 2011

Key indicators: single-crystal X-ray study; $T = 293$ K; mean $\sigma(\text{C}-\text{C}) = 0.002$ Å; R factor = 0.050; wR factor = 0.159; data-to-parameter ratio = 26.1.

In the title molecule, $\text{C}_{27}\text{H}_{44}\text{N}_2\text{O}_2$, the seven-membered ring adopts a boat-shaped conformation, with two C atoms of the fused benzene ring forming the stern and the methine C atom forming the prow. The hexadecyl substituent occupies an equatorial position, with the aliphatic chain exhibiting an extended zigzag conformation.

Related literature

For the 3-tetradecyl-substituted analog, see: Dardouri *et al.* (2011).



Experimental

Crystal data

$\text{C}_{27}\text{H}_{44}\text{N}_2\text{O}_2$
 $M_r = 428.64$
Monoclinic, $P2_1/c$
 $a = 8.1426$ (1) Å
 $b = 36.2705$ (5) Å
 $c = 9.4090$ (1) Å
 $\beta = 114.611$ (1)°

$V = 2526.38$ (5) Å³
 $Z = 4$
Mo $K\alpha$ radiation
 $\mu = 0.07$ mm⁻¹
 $T = 293$ K
 $0.50 \times 0.30 \times 0.10$ mm

Data collection

Bruker X8 APEXII diffractometer
45223 measured reflections
7353 independent reflections

5105 reflections with $I > 2\sigma(I)$
 $R_{\text{int}} = 0.036$

Refinement

$R[F^2 > 2\sigma(F^2)] = 0.050$
 $wR(F^2) = 0.159$
 $S = 1.03$
7353 reflections

282 parameters
H-atom parameters constrained
 $\Delta\rho_{\text{max}} = 0.34$ e Å⁻³
 $\Delta\rho_{\text{min}} = -0.26$ e Å⁻³

Data collection: APEX2 (Bruker, 2008); cell refinement: SAINT (Bruker, 2008); data reduction: SAINT; program(s) used to solve structure: SHELXS97 (Sheldrick, 2008); program(s) used to refine structure: SHELXL97 (Sheldrick, 2008); molecular graphics: XSEED (Barbour, 2001); software used to prepare material for publication: publCIF (Westrip, 2010).

We thank Université Mohammed V-Agdal and the University of Malaya for supporting this study.

Supplementary data and figures for this paper are available from the IUCr electronic archives (Reference: LH5231).

References

- Barbour, L. J. (2001). *J. Supramol. Chem.* **1**, 189–191.
Bruker (2008). APEX2 and SAINT. Bruker AXS Inc., Madison, Wisconsin, USA.
Dardouri, R., Ouazzani Chahdi, F., Saffon, N., Essassi, E. M. & Ng, S. W. (2011). *Acta Cryst.* **E67**, o674.
Sheldrick, G. M. (2008). *Acta Cryst.* **A64**, 112–122.
Westrip, S. P. (2010). *J. Appl. Cryst.* **43**, 920–925.

supplementary materials

Acta Cryst. (2011). E67, o1155 [doi:10.1107/S1600536811013663]

3-Hexadecyl-1,5-dimethyl-1*H*-1,5-benzodiazepine-2,4(3*H*,5*H*)-dione

R. Dardouri, Y. K. Rodi, S. Ladeira, E. M. Essassi and S. W. Ng

Comment

The methylene part of 1,5-dimethyl-1,5-benzodiazepine-2,4-dione is relatively acidic, and one proton can be abstracted by using potassium *t*-butoxide. The resulting carbanion can undergo a nucleophilic substitution with a dibromoalkane to form a 3-substituted derivative. In a previous study, the compound was reacted with 1-bromotetradecane to give the tetradecyl substituted derivative (Dardouri *et al.*, 2011). The corresponding hexadecyl title compound (Fig. 1) was obtained by using 1-bromohexadecane.

Experimental

To a solution of the potassium *t*-butoxide (0.42 g, 3.6 mmol) in DMF (15 ml) was added 1,5-dimethyl-1,5-benzodiazepine-2,4-dione (0.50 g, 2.4 mmol) and 1-bromotetradecane (0.88 ml, 2.88 mmol). Stirring was continued for 24 h. The reaction was monitored by thin layer chromatography. The mixture was filtered and the solution evaporated to give colorless crystals.

Refinement

Carbon-bound H-atoms were placed in calculated positions (C—H 0.93–0.98 Å) and were included in the refinement in the riding model approximation, with $U(\text{H})$ set to 1.2–1.5 $U_{\text{eq}}(\text{C})$. Two reflections with bad agreements (0 2 0, 0 4 0) were omitted from the refinement.

Figures




Fig. 1. Thermal ellipsoid plot (Barbour, 2001) of $\text{C}_{27}\text{H}_{44}\text{N}_2\text{O}_2$ at the 50% probability level; hydrogen atoms are drawn as arbitrary radius.

3-Hexadecyl-1,5-dimethyl-1*H*-1,5-benzodiazepine- 2,4(3*H*,5*H*)-dione

Crystal data

$\text{C}_{27}\text{H}_{44}\text{N}_2\text{O}_2$

$M_r = 428.64$

Monoclinic, $P2_1/c$

Hall symbol: -P 2ybc

$a = 8.1426 (1) \text{ \AA}$

$b = 36.2705 (5) \text{ \AA}$

$F(000) = 944$

$D_x = 1.127 \text{ Mg m}^{-3}$

Mo $K\alpha$ radiation, $\lambda = 0.71073 \text{ \AA}$

Cell parameters from 9990 reflections

$\theta = 2.6\text{--}30.0^\circ$

$\mu = 0.07 \text{ mm}^{-1}$

supplementary materials

$c = 9.4090$ (1) Å
 $\beta = 114.611$ (1)°
 $V = 2526.38$ (5) Å³
 $Z = 4$

$T = 293$ K
Plate, colorless
 $0.50 \times 0.30 \times 0.10$ mm

Data collection

Bruker X8 APEXII diffractometer
Radiation source: fine-focus sealed tube graphite
 φ and ω scans
45223 measured reflections
7353 independent reflections

5105 reflections with $I > 2\sigma(I)$
 $R_{\text{int}} = 0.036$
 $\theta_{\text{max}} = 30.0^\circ$, $\theta_{\text{min}} = 2.5^\circ$
 $h = -10 \rightarrow 11$
 $k = -50 \rightarrow 51$
 $l = -12 \rightarrow 13$

Refinement

Refinement on F^2
Least-squares matrix: full
 $R[F^2 > 2\sigma(F^2)] = 0.050$
 $wR(F^2) = 0.159$
 $S = 1.03$
7353 reflections
282 parameters
0 restraints

Primary atom site location: structure-invariant direct methods
Secondary atom site location: difference Fourier map
Hydrogen site location: inferred from neighbouring sites
H-atom parameters constrained
 $w = 1/[\sigma^2(F_o^2) + (0.0871P)^2 + 0.3018P]$
where $P = (F_o^2 + 2F_c^2)/3$
 $(\Delta/\sigma)_{\text{max}} = 0.001$
 $\Delta\rho_{\text{max}} = 0.34 \text{ e \AA}^{-3}$
 $\Delta\rho_{\text{min}} = -0.26 \text{ e \AA}^{-3}$

Fractional atomic coordinates and isotropic or equivalent isotropic displacement parameters (Å²)

	<i>x</i>	<i>y</i>	<i>z</i>	$U_{\text{iso}}^*/U_{\text{eq}}$
O1	0.02825 (12)	0.20901 (2)	0.37201 (10)	0.0323 (2)
O2	-0.39162 (13)	0.17141 (3)	0.40983 (10)	0.0376 (2)
N1	-0.19184 (13)	0.22225 (3)	0.13447 (11)	0.0271 (2)
N2	-0.51023 (13)	0.19449 (3)	0.16414 (12)	0.0268 (2)
C1	-0.35060 (17)	0.21187 (3)	0.00186 (14)	0.0273 (3)
C2	-0.3522 (2)	0.21567 (4)	-0.14654 (15)	0.0400 (3)
H2	-0.2498	0.2243	-0.1560	0.048*
C3	-0.5039 (2)	0.20671 (4)	-0.27923 (16)	0.0491 (4)
H3	-0.5028	0.2088	-0.3773	0.059*
C4	-0.6573 (2)	0.19462 (4)	-0.26542 (17)	0.0485 (4)
H4	-0.7606	0.1891	-0.3546	0.058*
C5	-0.65797 (19)	0.19072 (3)	-0.11957 (16)	0.0384 (3)
H5	-0.7621	0.1825	-0.1117	0.046*
C6	-0.50464 (16)	0.19891 (3)	0.01618 (14)	0.0267 (3)
C7	-0.67424 (18)	0.20524 (4)	0.18255 (19)	0.0388 (3)
H7A	-0.6415	0.2151	0.2855	0.058*

H7B	-0.7501	0.1840	0.1681	0.058*
H7C	-0.7383	0.2236	0.1061	0.058*
C8	-0.37885 (16)	0.17653 (3)	0.28661 (13)	0.0249 (2)
C9	-0.21270 (15)	0.16463 (3)	0.26138 (13)	0.0233 (2)
H9A	-0.2522	0.1529	0.1587	0.028*
C10	-0.11159 (16)	0.20013 (3)	0.26219 (13)	0.0237 (2)
C11	-0.09947 (19)	0.25653 (4)	0.12803 (18)	0.0384 (3)
H11A	-0.0195	0.2638	0.2321	0.058*
H11B	-0.1873	0.2756	0.0806	0.058*
H11C	-0.0312	0.2526	0.0672	0.058*
C12	-0.09467 (16)	0.13780 (3)	0.38687 (13)	0.0268 (3)
H12A	-0.1690	0.1176	0.3940	0.032*
H12B	-0.0452	0.1504	0.4870	0.032*
C13	0.06032 (17)	0.12220 (3)	0.35425 (14)	0.0304 (3)
H13A	0.0108	0.1073	0.2601	0.036*
H13B	0.1266	0.1424	0.3354	0.036*
C14	0.19019 (17)	0.09876 (3)	0.48780 (14)	0.0279 (3)
H14A	0.1243	0.0782	0.5046	0.034*
H14B	0.2367	0.1134	0.5826	0.034*
C15	0.34868 (17)	0.08387 (3)	0.45879 (14)	0.0295 (3)
H15A	0.3025	0.0687	0.3655	0.035*
H15B	0.4133	0.1044	0.4398	0.035*
C16	0.47967 (16)	0.06121 (3)	0.59449 (14)	0.0283 (3)
H16A	0.4146	0.0409	0.6139	0.034*
H16B	0.5262	0.0765	0.6875	0.034*
C17	0.63784 (16)	0.04591 (3)	0.56685 (14)	0.0278 (3)
H17A	0.7022	0.0662	0.5462	0.033*
H17B	0.5914	0.0304	0.4746	0.033*
C18	0.77024 (16)	0.02355 (3)	0.70403 (14)	0.0278 (3)
H18A	0.8161	0.0390	0.7964	0.033*
H18B	0.7060	0.0031	0.7243	0.033*
C19	0.92947 (16)	0.00835 (3)	0.67688 (14)	0.0277 (3)
H19A	0.8839	-0.0079	0.5869	0.033*
H19B	0.9910	0.0287	0.6527	0.033*
C20	1.06521 (16)	-0.01274 (3)	0.81635 (14)	0.0292 (3)
H20A	1.1111	0.0036	0.9062	0.035*
H20B	1.0035	-0.0330	0.8408	0.035*
C21	1.22377 (17)	-0.02808 (3)	0.78946 (14)	0.0297 (3)
H21A	1.2842	-0.0079	0.7631	0.036*
H21B	1.1782	-0.0448	0.7009	0.036*
C22	1.36085 (17)	-0.04850 (4)	0.93023 (15)	0.0319 (3)
H22A	1.3000	-0.0685	0.9571	0.038*
H22B	1.4070	-0.0317	1.0185	0.038*
C23	1.51950 (17)	-0.06420 (4)	0.90372 (14)	0.0317 (3)
H23A	1.5760	-0.0445	0.8706	0.038*
H23B	1.4744	-0.0822	0.8199	0.038*
C24	1.66148 (17)	-0.08248 (3)	1.04886 (14)	0.0309 (3)
H24A	1.7070	-0.0644	1.1323	0.037*
H24B	1.6045	-0.1020	1.0824	0.037*

supplementary materials

C25	1.81995 (17)	-0.09856 (4)	1.02357 (15)	0.0318 (3)
H25A	1.7764	-0.1184	0.9479	0.038*
H25B	1.8696	-0.0796	0.9800	0.038*
C26	1.96936 (17)	-0.11326 (3)	1.17273 (15)	0.0331 (3)
H26A	2.0196	-0.0929	1.2450	0.040*
H26B	1.9174	-0.1306	1.2210	0.040*
C27	2.1211 (2)	-0.13226 (5)	1.1471 (2)	0.0527 (4)
H27A	2.2105	-0.1408	1.2454	0.079*
H27B	2.1755	-0.1151	1.1017	0.079*
H27C	2.0733	-0.1528	1.0778	0.079*

Atomic displacement parameters (\AA^2)

	U^{11}	U^{22}	U^{33}	U^{12}	U^{13}	U^{23}
O1	0.0250 (5)	0.0383 (5)	0.0285 (5)	-0.0006 (4)	0.0060 (4)	0.0013 (4)
O2	0.0388 (6)	0.0456 (5)	0.0359 (5)	0.0037 (4)	0.0231 (4)	0.0053 (4)
N1	0.0210 (5)	0.0307 (5)	0.0285 (5)	0.0020 (4)	0.0092 (4)	0.0081 (4)
N2	0.0197 (5)	0.0258 (5)	0.0348 (5)	0.0031 (4)	0.0112 (4)	0.0009 (4)
C1	0.0275 (6)	0.0265 (6)	0.0246 (5)	0.0091 (5)	0.0074 (5)	0.0048 (4)
C2	0.0447 (8)	0.0452 (8)	0.0309 (7)	0.0174 (6)	0.0166 (6)	0.0137 (6)
C3	0.0656 (11)	0.0486 (9)	0.0249 (7)	0.0224 (8)	0.0106 (7)	0.0087 (6)
C4	0.0545 (10)	0.0342 (7)	0.0314 (7)	0.0090 (7)	-0.0074 (7)	-0.0003 (6)
C5	0.0319 (7)	0.0271 (6)	0.0408 (7)	0.0028 (5)	0.0000 (6)	0.0011 (5)
C6	0.0249 (6)	0.0203 (5)	0.0288 (6)	0.0065 (4)	0.0052 (5)	0.0019 (4)
C7	0.0238 (7)	0.0365 (7)	0.0601 (9)	0.0034 (5)	0.0215 (7)	0.0009 (6)
C8	0.0238 (6)	0.0231 (5)	0.0279 (6)	-0.0004 (4)	0.0110 (5)	-0.0012 (4)
C9	0.0223 (6)	0.0246 (5)	0.0216 (5)	0.0048 (4)	0.0080 (4)	0.0023 (4)
C10	0.0220 (6)	0.0276 (5)	0.0236 (5)	0.0057 (4)	0.0116 (5)	0.0027 (4)
C11	0.0294 (7)	0.0373 (7)	0.0492 (8)	-0.0004 (5)	0.0171 (6)	0.0144 (6)
C12	0.0269 (6)	0.0282 (6)	0.0248 (5)	0.0070 (5)	0.0102 (5)	0.0076 (4)
C13	0.0315 (7)	0.0328 (6)	0.0271 (6)	0.0122 (5)	0.0125 (5)	0.0085 (5)
C14	0.0280 (6)	0.0282 (6)	0.0273 (6)	0.0082 (5)	0.0111 (5)	0.0059 (5)
C15	0.0283 (7)	0.0311 (6)	0.0292 (6)	0.0088 (5)	0.0120 (5)	0.0067 (5)
C16	0.0247 (6)	0.0318 (6)	0.0274 (6)	0.0073 (5)	0.0098 (5)	0.0049 (5)
C17	0.0244 (6)	0.0303 (6)	0.0275 (6)	0.0067 (5)	0.0097 (5)	0.0042 (5)
C18	0.0248 (6)	0.0297 (6)	0.0289 (6)	0.0060 (5)	0.0111 (5)	0.0054 (5)
C19	0.0247 (6)	0.0295 (6)	0.0283 (6)	0.0070 (5)	0.0103 (5)	0.0037 (5)
C20	0.0254 (6)	0.0327 (6)	0.0294 (6)	0.0069 (5)	0.0113 (5)	0.0067 (5)
C21	0.0269 (6)	0.0341 (6)	0.0280 (6)	0.0081 (5)	0.0113 (5)	0.0055 (5)
C22	0.0274 (7)	0.0370 (7)	0.0319 (6)	0.0096 (5)	0.0128 (5)	0.0077 (5)
C23	0.0284 (7)	0.0367 (6)	0.0303 (6)	0.0103 (5)	0.0124 (5)	0.0062 (5)
C24	0.0281 (7)	0.0342 (6)	0.0306 (6)	0.0084 (5)	0.0124 (5)	0.0048 (5)
C25	0.0291 (7)	0.0332 (6)	0.0341 (6)	0.0090 (5)	0.0140 (6)	0.0044 (5)
C26	0.0275 (7)	0.0293 (6)	0.0374 (7)	0.0039 (5)	0.0083 (6)	0.0017 (5)
C27	0.0346 (8)	0.0525 (9)	0.0647 (10)	0.0169 (7)	0.0145 (8)	0.0033 (8)

Geometric parameters (\AA , $^\circ$)

O1—C10	1.2197 (14)	C15—H15A	0.9700
--------	-------------	----------	--------

O2—C8	1.2206 (14)	C15—H15B	0.9700
N1—C10	1.3640 (14)	C16—C17	1.5209 (16)
N1—C1	1.4235 (16)	C16—H16A	0.9700
N1—C11	1.4676 (16)	C16—H16B	0.9700
N2—C8	1.3676 (15)	C17—C18	1.5254 (16)
N2—C6	1.4208 (15)	C17—H17A	0.9700
N2—C7	1.4690 (16)	C17—H17B	0.9700
C1—C6	1.3975 (18)	C18—C19	1.5252 (16)
C1—C2	1.3976 (17)	C18—H18A	0.9700
C2—C3	1.380 (2)	C18—H18B	0.9700
C2—H2	0.9300	C19—C20	1.5233 (16)
C3—C4	1.380 (2)	C19—H19A	0.9700
C3—H3	0.9300	C19—H19B	0.9700
C4—C5	1.382 (2)	C20—C21	1.5207 (16)
C4—H4	0.9300	C20—H20A	0.9700
C5—C6	1.3964 (18)	C20—H20B	0.9700
C5—H5	0.9300	C21—C22	1.5223 (17)
C7—H7A	0.9600	C21—H21A	0.9700
C7—H7B	0.9600	C21—H21B	0.9700
C7—H7C	0.9600	C22—C23	1.5244 (17)
C8—C9	1.5298 (16)	C22—H22A	0.9700
C9—C12	1.5230 (15)	C22—H22B	0.9700
C9—C10	1.5266 (16)	C23—C24	1.5251 (17)
C9—H9A	0.9800	C23—H23A	0.9700
C11—H11A	0.9600	C23—H23B	0.9700
C11—H11B	0.9600	C24—C25	1.5222 (17)
C11—H11C	0.9600	C24—H24A	0.9700
C12—C13	1.5264 (16)	C24—H24B	0.9700
C12—H12A	0.9700	C25—C26	1.5205 (17)
C12—H12B	0.9700	C25—H25A	0.9700
C13—C14	1.5210 (16)	C25—H25B	0.9700
C13—H13A	0.9700	C26—C27	1.5196 (19)
C13—H13B	0.9700	C26—H26A	0.9700
C14—C15	1.5248 (16)	C26—H26B	0.9700
C14—H14A	0.9700	C27—H27A	0.9600
C14—H14B	0.9700	C27—H27B	0.9600
C15—C16	1.5195 (16)	C27—H27C	0.9600
C10—N1—C1	122.93 (10)	C15—C16—C17	113.69 (10)
C10—N1—C11	117.97 (10)	C15—C16—H16A	108.8
C1—N1—C11	118.81 (10)	C17—C16—H16A	108.8
C8—N2—C6	123.12 (10)	C15—C16—H16B	108.8
C8—N2—C7	117.24 (10)	C17—C16—H16B	108.8
C6—N2—C7	119.16 (10)	H16A—C16—H16B	107.7
C6—C1—C2	119.60 (12)	C16—C17—C18	113.56 (10)
C6—C1—N1	121.90 (10)	C16—C17—H17A	108.9
C2—C1—N1	118.48 (12)	C18—C17—H17A	108.9
C3—C2—C1	120.87 (15)	C16—C17—H17B	108.9
C3—C2—H2	119.6	C18—C17—H17B	108.9
C1—C2—H2	119.6	H17A—C17—H17B	107.7

supplementary materials

C4—C3—C2	119.58 (14)	C19—C18—C17	113.64 (10)
C4—C3—H3	120.2	C19—C18—H18A	108.8
C2—C3—H3	120.2	C17—C18—H18A	108.8
C3—C4—C5	120.28 (14)	C19—C18—H18B	108.8
C3—C4—H4	119.9	C17—C18—H18B	108.8
C5—C4—H4	119.9	H18A—C18—H18B	107.7
C4—C5—C6	120.98 (14)	C20—C19—C18	113.56 (10)
C4—C5—H5	119.5	C20—C19—H19A	108.9
C6—C5—H5	119.5	C18—C19—H19A	108.9
C5—C6—C1	118.67 (12)	C20—C19—H19B	108.9
C5—C6—N2	119.41 (12)	C18—C19—H19B	108.9
C1—C6—N2	121.91 (11)	H19A—C19—H19B	107.7
N2—C7—H7A	109.5	C21—C20—C19	113.71 (10)
N2—C7—H7B	109.5	C21—C20—H20A	108.8
H7A—C7—H7B	109.5	C19—C20—H20A	108.8
N2—C7—H7C	109.5	C21—C20—H20B	108.8
H7A—C7—H7C	109.5	C19—C20—H20B	108.8
H7B—C7—H7C	109.5	H20A—C20—H20B	107.7
O2—C8—N2	121.78 (11)	C20—C21—C22	113.51 (10)
O2—C8—C9	122.36 (11)	C20—C21—H21A	108.9
N2—C8—C9	115.83 (10)	C22—C21—H21A	108.9
C12—C9—C10	111.73 (10)	C20—C21—H21B	108.9
C12—C9—C8	111.91 (9)	C22—C21—H21B	108.9
C10—C9—C8	105.74 (9)	H21A—C21—H21B	107.7
C12—C9—H9A	109.1	C21—C22—C23	113.81 (10)
C10—C9—H9A	109.1	C21—C22—H22A	108.8
C8—C9—H9A	109.1	C23—C22—H22A	108.8
O1—C10—N1	121.80 (11)	C21—C22—H22B	108.8
O1—C10—C9	122.55 (10)	C23—C22—H22B	108.8
N1—C10—C9	115.60 (10)	H22A—C22—H22B	107.7
N1—C11—H11A	109.5	C22—C23—C24	113.20 (10)
N1—C11—H11B	109.5	C22—C23—H23A	108.9
H11A—C11—H11B	109.5	C24—C23—H23A	108.9
N1—C11—H11C	109.5	C22—C23—H23B	108.9
H11A—C11—H11C	109.5	C24—C23—H23B	108.9
H11B—C11—H11C	109.5	H23A—C23—H23B	107.8
C9—C12—C13	112.87 (9)	C25—C24—C23	113.73 (10)
C9—C12—H12A	109.0	C25—C24—H24A	108.8
C13—C12—H12A	109.0	C23—C24—H24A	108.8
C9—C12—H12B	109.0	C25—C24—H24B	108.8
C13—C12—H12B	109.0	C23—C24—H24B	108.8
H12A—C12—H12B	107.8	H24A—C24—H24B	107.7
C14—C13—C12	112.94 (9)	C26—C25—C24	113.24 (10)
C14—C13—H13A	109.0	C26—C25—H25A	108.9
C12—C13—H13A	109.0	C24—C25—H25A	108.9
C14—C13—H13B	109.0	C26—C25—H25B	108.9
C12—C13—H13B	109.0	C24—C25—H25B	108.9
H13A—C13—H13B	107.8	H25A—C25—H25B	107.7
C13—C14—C15	113.52 (10)	C27—C26—C25	113.80 (12)

C13—C14—H14A	108.9	C27—C26—H26A	108.8
C15—C14—H14A	108.9	C25—C26—H26A	108.8
C13—C14—H14B	108.9	C27—C26—H26B	108.8
C15—C14—H14B	108.9	C25—C26—H26B	108.8
H14A—C14—H14B	107.7	H26A—C26—H26B	107.7
C16—C15—C14	113.04 (10)	C26—C27—H27A	109.5
C16—C15—H15A	109.0	C26—C27—H27B	109.5
C14—C15—H15A	109.0	H27A—C27—H27B	109.5
C16—C15—H15B	109.0	C26—C27—H27C	109.5
C14—C15—H15B	109.0	H27A—C27—H27C	109.5
H15A—C15—H15B	107.8	H27B—C27—H27C	109.5
C10—N1—C1—C6	50.83 (16)	N2—C8—C9—C10	70.80 (12)
C11—N1—C1—C6	-135.46 (12)	C1—N1—C10—O1	177.18 (11)
C10—N1—C1—C2	-130.67 (12)	C11—N1—C10—O1	3.42 (17)
C11—N1—C1—C2	43.04 (16)	C1—N1—C10—C9	-5.49 (16)
C6—C1—C2—C3	-0.15 (19)	C11—N1—C10—C9	-179.24 (10)
N1—C1—C2—C3	-178.69 (12)	C12—C9—C10—O1	-16.10 (15)
C1—C2—C3—C4	1.4 (2)	C8—C9—C10—O1	105.89 (12)
C2—C3—C4—C5	-1.4 (2)	C12—C9—C10—N1	166.59 (9)
C3—C4—C5—C6	0.0 (2)	C8—C9—C10—N1	-71.43 (12)
C4—C5—C6—C1	1.23 (18)	C10—C9—C12—C13	-68.09 (13)
C4—C5—C6—N2	-179.85 (11)	C8—C9—C12—C13	173.55 (10)
C2—C1—C6—C5	-1.16 (17)	C9—C12—C13—C14	173.66 (10)
N1—C1—C6—C5	177.32 (11)	C12—C13—C14—C15	-178.37 (10)
C2—C1—C6—N2	179.94 (11)	C13—C14—C15—C16	178.71 (11)
N1—C1—C6—N2	-1.57 (17)	C14—C15—C16—C17	179.55 (10)
C8—N2—C6—C5	132.06 (12)	C15—C16—C17—C18	179.34 (10)
C7—N2—C6—C5	-39.70 (15)	C16—C17—C18—C19	-179.70 (10)
C8—N2—C6—C1	-49.05 (16)	C17—C18—C19—C20	177.81 (10)
C7—N2—C6—C1	139.19 (12)	C18—C19—C20—C21	179.76 (10)
C6—N2—C8—O2	-176.04 (11)	C19—C20—C21—C22	178.91 (11)
C7—N2—C8—O2	-4.13 (17)	C20—C21—C22—C23	179.57 (11)
C6—N2—C8—C9	6.17 (15)	C21—C22—C23—C24	176.41 (11)
C7—N2—C8—C9	178.07 (10)	C22—C23—C24—C25	179.53 (11)
O2—C8—C9—C12	14.88 (16)	C23—C24—C25—C26	174.22 (11)
N2—C8—C9—C12	-167.34 (10)	C24—C25—C26—C27	175.17 (12)
O2—C8—C9—C10	-106.98 (12)		

Fig. 1

