

1-(2,3-Dimethoxybenzylidene)-2-(2,4-dinitrophenyl)hydrazine

Xianrong Xin,^a Min Li,^a Zhimin Chen^a and Ruitao Zhu^{b*}

^aDepartment of Chemistry and Chemical Engineering, Taiyuan Institute of Technology, Taiyuan 030008, People's Republic of China, and ^bDepartment of Chemistry, Taiyuan Normal University, Taiyuan 030031, People's Republic of China
Correspondence e-mail: ruitaozhu@126.com


Received 29 March 2011; accepted 13 April 2011

Key indicators: single-crystal X-ray study; $T = 298\text{ K}$; mean $\sigma(\text{C}-\text{C}) = 0.005\text{ \AA}$; R factor = 0.063; wR factor = 0.178; data-to-parameter ratio = 11.9.

In the title compound, $\text{C}_{15}\text{H}_{14}\text{N}_4\text{O}_6$, the dihedral angle between the aromatic rings is $3.7(4)^\circ$. The nitro groups make dihedral angles of $6.0(4)$ and $5.2(4)^\circ$ with the parent ring and are oriented at $6.0(6)^\circ$ with respect to each other. The methoxy groups are inclined at $54.0(2)$ and $2.5(3)^\circ$ with respect to the benzene ring to which they are attached. In the crystal, molecules are linked by weak $\text{C}-\text{H}\cdots\text{O}$ interactions. The molecular conformation is consolidated by an intramolecular $\text{N}-\text{H}\cdots\text{O}$ hydrogen bond.

Related literature

For general background to the properties of Schiff base compounds, see: Mufakkar *et al.* (2010); Tahir *et al.* (2010). For related structures, see: Salhin *et al.* (2007); Tameem *et al.* (2008); Shao *et al.* (2008).



Experimental

Crystal data

$\text{C}_{15}\text{H}_{14}\text{N}_4\text{O}_6$
 $M_r = 346.30$
Triclinic, $P\bar{1}$
 $a = 7.8409(8)\text{ \AA}$
 $b = 7.9200(9)\text{ \AA}$
 $c = 13.8961(14)\text{ \AA}$
 $\alpha = 85.038(2)^\circ$
 $\beta = 82.773(1)^\circ$

$\gamma = 65.894(1)^\circ$
 $V = 780.85(14)\text{ \AA}^3$
 $Z = 2$
Mo $K\alpha$ radiation

$\mu = 0.12\text{ mm}^{-1}$
 $T = 298\text{ K}$
 $0.43 \times 0.38 \times 0.37\text{ mm}$

Data collection

Bruker SMART CCD
diffractometer
Absorption correction: multi-scan
(*SADABS*; Bruker, 2007)
 $T_{\min} = 0.952$, $T_{\max} = 0.958$

4130 measured reflections
2725 independent reflections
1414 reflections with $I > 2\sigma(I)$
 $R_{\text{int}} = 0.032$

Refinement

$R[F^2 > 2\sigma(F^2)] = 0.063$
 $wR(F^2) = 0.178$
 $S = 1.06$
2725 reflections

229 parameters
H-atom parameters constrained
 $\Delta\rho_{\max} = 0.28\text{ e \AA}^{-3}$
 $\Delta\rho_{\min} = -0.25\text{ e \AA}^{-3}$

Table 1
Hydrogen-bond geometry (\AA , $^\circ$).

$D-\text{H}\cdots A$	$D-\text{H}$	$\text{H}\cdots A$	$D\cdots A$	$D-\text{H}\cdots A$
N1—H1 \cdots O1	0.86	1.99	2.625 (4)	130
C14—H14A \cdots O4 ⁱ	0.96	2.48	3.431 (4)	170

Symmetry code: (i) $-x + 1, -y, -z + 1$.

Data collection: *SMART* (Bruker, 2007); cell refinement: *SAINT* (Bruker, 2007); data reduction: *SAINT*; program(s) used to solve structure: *SHELXS97* (Sheldrick, 2008); program(s) used to refine structure: *SHELXL97* (Sheldrick, 2008); molecular graphics: *SHELXTL* (Sheldrick, 2008) and *PLATON* (Spek, 2009); software used to prepare material for publication: *SHELXTL*.

The authors thank Professor Daqi Wang, Liaocheng University, for use of the X-ray diffraction facility.

Supplementary data and figures for this paper are available from the IUCr electronic archives (Reference: PV2405).

References

- Bruker (2007). *SMART*, *SAINT* and *SADABS*. Bruker AXS Inc., Madison, Wisconsin, USA.
- Mufakkar, M., Tahir, M. N., Tariq, M. I., Ahmad, S. & Sarfraz, M. (2010). *Acta Cryst. E66*, o1887.
- Salhin, A., Tameem, A. A., Saad, B., Ng, S.-L. & Fun, H.-K. (2007). *Acta Cryst. E63*, o2880.
- Shao, J., Lin, I., Yu, M., Cai, Z. & Lin, I. (2008). *Talanta*, **75**, 551–555.
- Sheldrick, G. M. (2008). *Acta Cryst. A64*, 112–122.
- Spek, A. L. (2009). *Acta Cryst. D65*, 148–155.
- Tahir, M. N., Tariq, M. I., Ahmad, S., Sarfraz, M. & Ather, A. Q. (2010). *Acta Cryst. E66*, o1817.
- Tameem, A., Saad, B., Salhin, A. M., Jebas, S. R. & Fun, H.-K. (2008). *Acta Cryst. E64*, o679–o680.

supplementary materials

Acta Cryst. (2011). E67, o1169 [doi:10.1107/S1600536811013894]

1-(2,3-Dimethoxybenzylidene)-2-(2,4-dinitrophenyl)hydrazine

X. Xin, M. Li, Z. Chen and R. Zhu

Comment

In view of the importance of hydrazone derivatives in chemical and biological applications (Shao *et al.*, 2008), a series of hydrazone derivatives has been prepared, and several X-ray structures have been reported (Salhin *et al.*, 2007; Aameem *et al.*, 2008). Here, we report the crystal structure of the title compound.

The bond distances and bond angles in the title compound (Fig. 1) are in agreement with the corresponding bond distances and angles reported in the crystal structures of closely related compounds, (Salhin *et al.*, 2007; Tameem *et al.*, 2008). In the crystal structure (Fig. 2), the molecules are linked by C—H···O weak interactions (Table 1). The molecular conformation is consolidated by an intramolecular N—H···O hydrogen bonding interaction (Table 1).

Experimental

Equimolar quantities of 2,4-dinitrophenylhydrazine (0.99 g, 5.0 mmol) and 2,3-dimethoxybenzaldehyde (0.83 g, 5.0 mmol) were refluxed in ethanol (20 ml) for 30 min and rotary evaporated. The crystals of the title compound were grown by recrystallization from an ethanol solution.

Refinement

H atoms were placed in idealized positions and allowed to ride on their respective parent atoms, with C—H = 0.93 and 0.96 Å, for methylene and ary H-atoms, respectively, and N—H = 0.86 Å with $U_{\text{iso}}(\text{H}) = 1.2U_{\text{eq}}(\text{methylene C/N})$ and $1.5U_{\text{eq}}(\text{aryl C})$.

Figures




Fig. 1. A view of the molecular structure of the title compound; displacement ellipsoids are drawn at the 30% probability level.




Fig. 2. Part of the crystal structure of (I) with hydrogen bonds drawn as dashed lines.

supplementary materials

1-(2,3-Dimethoxybenzylidene)-2-(2,4-dinitrophenyl)hydrazine

Crystal data

C ₁₅ H ₁₄ N ₄ O ₆	Z = 2
M _r = 346.30	F(000) = 360
Triclinic, PT	D _x = 1.473 Mg m ⁻³
Hall symbol: -P 1	Mo K α radiation, λ = 0.71073 Å
a = 7.8409 (8) Å	Cell parameters from 855 reflections
b = 7.9200 (9) Å	θ = 2.8–22.8°
c = 13.8961 (14) Å	μ = 0.12 mm ⁻¹
α = 85.038 (2)°	T = 298 K
β = 82.773 (1)°	Block, orange
γ = 65.894 (1)°	0.43 × 0.38 × 0.37 mm
V = 780.85 (14) Å ³	

Data collection

Bruker SMART CCD diffractometer	2725 independent reflections
Radiation source: fine-focus sealed tube graphite	1414 reflections with $I > 2\sigma(I)$
φ and ω scans	$R_{\text{int}} = 0.032$
Absorption correction: multi-scan (SADABS; Bruker, 2007)	$\theta_{\text{max}} = 25.0^\circ$, $\theta_{\text{min}} = 2.8^\circ$
$T_{\text{min}} = 0.952$, $T_{\text{max}} = 0.958$	$h = -9 \rightarrow 9$
4130 measured reflections	$k = -9 \rightarrow 9$
	$l = -16 \rightarrow 8$

Refinement

Refinement on F^2	Secondary atom site location: difference Fourier map
Least-squares matrix: full	Hydrogen site location: inferred from neighbouring sites
$R[F^2 > 2\sigma(F^2)] = 0.063$	H-atom parameters constrained
$wR(F^2) = 0.178$	$w = 1/[\sigma^2(F_o^2) + (0.0737P)^2]$
$S = 1.06$	where $P = (F_o^2 + 2F_c^2)/3$
2725 reflections	$(\Delta/\sigma)_{\text{max}} < 0.001$
229 parameters	$\Delta\rho_{\text{max}} = 0.28 \text{ e \AA}^{-3}$
0 restraints	$\Delta\rho_{\text{min}} = -0.25 \text{ e \AA}^{-3}$
Primary atom site location: structure-invariant direct methods	Extinction correction: SHELXL97 (Sheldrick, 2008), $F_c^* = kF_c[1 + 0.001xF_c^2\lambda^3/\sin(2\theta)]^{1/4}$
	Extinction coefficient: 0.028 (5)

Special details

Geometry. All e.s.d.'s (except the e.s.d. in the dihedral angle between two l.s. planes) are estimated using the full covariance matrix. The cell e.s.d.'s are taken into account individually in the estimation of e.s.d.'s in distances, angles and torsion angles; correlations between e.s.d.'s in cell parameters are only used when they are defined by crystal symmetry. An approximate (isotropic) treatment of cell e.s.d.'s is used for estimating e.s.d.'s involving l.s. planes.

Refinement. Refinement of F^2 against ALL reflections. The weighted R -factor wR and goodness of fit S are based on F^2 , conventional R -factors R are based on F , with F set to zero for negative F^2 . The threshold expression of $F^2 > \sigma(F^2)$ is used only for calculating R -factors(gt) etc. and is not relevant to the choice of reflections for refinement. R -factors based on F^2 are statistically about twice as large as those based on F , and R -factors based on ALL data will be even larger.

Fractional atomic coordinates and isotropic or equivalent isotropic displacement parameters (\AA^2)

	<i>x</i>	<i>y</i>	<i>z</i>	$U_{\text{iso}}^*/U_{\text{eq}}$
N1	0.6288 (3)	0.1887 (3)	0.55122 (17)	0.0672 (7)
H1	0.7290	0.1995	0.5243	0.081*
N2	0.4685 (3)	0.2598 (3)	0.50406 (17)	0.0627 (6)
N3	0.9691 (3)	0.0341 (4)	0.6504 (3)	0.0789 (8)
N4	0.6248 (5)	-0.1660 (4)	0.9164 (2)	0.0889 (9)
O1	0.9777 (3)	0.1238 (4)	0.57412 (19)	0.0936 (8)
O2	1.1061 (3)	-0.0474 (4)	0.6963 (2)	0.1191 (10)
O3	0.7741 (4)	-0.2406 (4)	0.9539 (2)	0.1259 (11)
O4	0.4763 (4)	-0.1674 (4)	0.95499 (18)	0.1107 (9)
O5	0.5156 (3)	0.5336 (3)	0.25258 (14)	0.0734 (6)
O6	0.2347 (3)	0.6845 (3)	0.13346 (16)	0.0810 (7)
C1	0.6302 (3)	0.1025 (3)	0.6391 (2)	0.0549 (7)
C2	0.7926 (3)	0.0256 (4)	0.6903 (2)	0.0626 (8)
C3	0.7883 (4)	-0.0608 (4)	0.7802 (2)	0.0681 (8)
H3	0.8956	-0.1094	0.8130	0.082*
C4	0.6287 (4)	-0.0754 (4)	0.8214 (2)	0.0652 (8)
C5	0.4661 (4)	-0.0033 (4)	0.7723 (2)	0.0651 (8)
H5	0.3567	-0.0136	0.8008	0.078*
C6	0.4688 (4)	0.0809 (4)	0.6841 (2)	0.0609 (7)
H6	0.3609	0.1260	0.6518	0.073*
C7	0.4850 (4)	0.3361 (4)	0.4206 (2)	0.0627 (7)
H7	0.6000	0.3394	0.3968	0.075*
C8	0.3275 (4)	0.4181 (4)	0.3620 (2)	0.0588 (7)
C9	0.3505 (4)	0.5101 (4)	0.2741 (2)	0.0602 (7)
C10	0.2012 (4)	0.5909 (4)	0.2156 (2)	0.0641 (8)
C11	0.0346 (4)	0.5735 (4)	0.2461 (2)	0.0744 (9)
H11	-0.0643	0.6240	0.2074	0.089*
C12	0.0117 (4)	0.4825 (4)	0.3330 (3)	0.0756 (9)
H12	-0.1026	0.4740	0.3525	0.091*
C13	0.1564 (4)	0.4046 (4)	0.3907 (2)	0.0684 (8)
H13	0.1404	0.3428	0.4488	0.082*
C14	0.6191 (4)	0.4797 (5)	0.1608 (2)	0.0847 (10)
H14A	0.5771	0.3991	0.1323	0.127*

supplementary materials

H14B	0.7503	0.4159	0.1691	0.127*
H14C	0.5998	0.5877	0.1190	0.127*
C15	0.0951 (5)	0.7525 (5)	0.0675 (3)	0.1016 (12)
H15A	0.0718	0.6516	0.0472	0.152*
H15B	0.1378	0.8107	0.0118	0.152*
H15C	-0.0187	0.8412	0.0987	0.152*

Atomic displacement parameters (\AA^2)

	U^{11}	U^{22}	U^{33}	U^{12}	U^{13}	U^{23}
N1	0.0509 (13)	0.0903 (17)	0.0649 (17)	-0.0324 (12)	-0.0047 (11)	-0.0079 (13)
N2	0.0547 (14)	0.0696 (15)	0.0639 (16)	-0.0245 (12)	-0.0081 (11)	-0.0030 (12)
N3	0.0472 (15)	0.0818 (19)	0.110 (2)	-0.0237 (14)	-0.0100 (15)	-0.0246 (16)
N4	0.117 (3)	0.088 (2)	0.081 (2)	-0.0550 (19)	-0.0386 (19)	0.0088 (16)
O1	0.0661 (14)	0.133 (2)	0.0922 (18)	-0.0513 (14)	0.0068 (12)	-0.0250 (15)
O2	0.0526 (13)	0.124 (2)	0.180 (3)	-0.0300 (13)	-0.0379 (16)	0.0068 (18)
O3	0.136 (2)	0.138 (2)	0.114 (2)	-0.0557 (18)	-0.0725 (19)	0.0353 (17)
O4	0.142 (2)	0.140 (2)	0.0873 (19)	-0.095 (2)	-0.0317 (16)	0.0251 (15)
O5	0.0724 (13)	0.0967 (15)	0.0642 (13)	-0.0473 (12)	-0.0049 (10)	-0.0064 (10)
O6	0.0827 (14)	0.0920 (15)	0.0730 (15)	-0.0389 (12)	-0.0219 (11)	0.0126 (12)
C1	0.0492 (15)	0.0540 (15)	0.0625 (19)	-0.0194 (13)	-0.0076 (12)	-0.0111 (13)
C2	0.0461 (16)	0.0640 (18)	0.080 (2)	-0.0201 (13)	-0.0103 (14)	-0.0183 (16)
C3	0.0621 (19)	0.0584 (17)	0.087 (2)	-0.0196 (15)	-0.0281 (16)	-0.0119 (16)
C4	0.077 (2)	0.0615 (17)	0.066 (2)	-0.0324 (15)	-0.0250 (15)	-0.0003 (14)
C5	0.0628 (18)	0.0648 (18)	0.076 (2)	-0.0329 (15)	-0.0142 (15)	0.0020 (15)
C6	0.0515 (16)	0.0651 (17)	0.070 (2)	-0.0256 (13)	-0.0144 (13)	-0.0007 (15)
C7	0.0573 (17)	0.0741 (19)	0.0600 (19)	-0.0302 (15)	-0.0019 (13)	-0.0060 (15)
C8	0.0543 (16)	0.0630 (17)	0.0602 (18)	-0.0240 (14)	-0.0005 (13)	-0.0129 (13)
C9	0.0572 (17)	0.0653 (17)	0.0628 (19)	-0.0287 (14)	-0.0009 (13)	-0.0125 (14)
C10	0.0627 (18)	0.0639 (18)	0.067 (2)	-0.0250 (15)	-0.0085 (14)	-0.0074 (15)
C11	0.0620 (19)	0.075 (2)	0.089 (2)	-0.0260 (16)	-0.0185 (16)	-0.0074 (17)
C12	0.0578 (18)	0.078 (2)	0.095 (3)	-0.0327 (16)	0.0006 (17)	-0.0115 (18)
C13	0.0627 (18)	0.0689 (19)	0.076 (2)	-0.0293 (15)	-0.0014 (15)	-0.0098 (15)
C14	0.080 (2)	0.113 (3)	0.072 (2)	-0.051 (2)	-0.0011 (16)	-0.0033 (18)
C15	0.095 (3)	0.120 (3)	0.084 (3)	-0.035 (2)	-0.032 (2)	0.019 (2)

Geometric parameters (\AA , $^\circ$)

N1—C1	1.345 (3)	C5—C6	1.347 (4)
N1—N2	1.375 (3)	C5—H5	0.9300
N1—H1	0.8600	C6—H6	0.9300
N2—C7	1.278 (3)	C7—C8	1.455 (4)
N3—O2	1.230 (3)	C7—H7	0.9300
N3—O1	1.234 (3)	C8—C13	1.395 (4)
N3—C2	1.451 (4)	C8—C9	1.397 (4)
N4—O4	1.223 (3)	C9—C10	1.408 (4)
N4—O3	1.236 (3)	C10—C11	1.379 (4)
N4—C4	1.450 (4)	C11—C12	1.382 (4)
O5—C9	1.375 (3)	C11—H11	0.9300

O5—C14	1.420 (3)	C12—C13	1.372 (4)
O6—C10	1.363 (3)	C12—H12	0.9300
O6—C15	1.420 (3)	C13—H13	0.9300
C1—C6	1.407 (4)	C14—H14A	0.9600
C1—C2	1.421 (4)	C14—H14B	0.9600
C2—C3	1.375 (4)	C14—H14C	0.9600
C3—C4	1.355 (4)	C15—H15A	0.9600
C3—H3	0.9300	C15—H15B	0.9600
C4—C5	1.404 (4)	C15—H15C	0.9600
C1—N1—N2	120.6 (2)	C8—C7—H7	119.4
C1—N1—H1	119.7	C13—C8—C9	119.6 (3)
N2—N1—H1	119.7	C13—C8—C7	121.7 (3)
C7—N2—N1	114.9 (2)	C9—C8—C7	118.7 (2)
O2—N3—O1	122.2 (3)	O5—C9—C8	117.6 (2)
O2—N3—C2	117.5 (3)	O5—C9—C10	121.8 (3)
O1—N3—C2	120.3 (3)	C8—C9—C10	120.2 (3)
O4—N4—O3	123.2 (3)	O6—C10—C11	125.7 (3)
O4—N4—C4	119.1 (3)	O6—C10—C9	115.8 (3)
O3—N4—C4	117.7 (3)	C11—C10—C9	118.4 (3)
C9—O5—C14	118.8 (2)	C10—C11—C12	121.4 (3)
C10—O6—C15	117.7 (2)	C10—C11—H11	119.3
N1—C1—C6	120.8 (2)	C12—C11—H11	119.3
N1—C1—C2	122.6 (3)	C13—C12—C11	120.4 (3)
C6—C1—C2	116.5 (3)	C13—C12—H12	119.8
C3—C2—C1	121.0 (3)	C11—C12—H12	119.8
C3—C2—N3	117.3 (3)	C12—C13—C8	120.0 (3)
C1—C2—N3	121.8 (3)	C12—C13—H13	120.0
C4—C3—C2	120.4 (3)	C8—C13—H13	120.0
C4—C3—H3	119.8	O5—C14—H14A	109.5
C2—C3—H3	119.8	O5—C14—H14B	109.5
C3—C4—C5	120.2 (3)	H14A—C14—H14B	109.5
C3—C4—N4	120.1 (3)	O5—C14—H14C	109.5
C5—C4—N4	119.6 (3)	H14A—C14—H14C	109.5
C6—C5—C4	119.9 (3)	H14B—C14—H14C	109.5
C6—C5—H5	120.0	O6—C15—H15A	109.5
C4—C5—H5	120.0	O6—C15—H15B	109.5
C5—C6—C1	121.9 (2)	H15A—C15—H15B	109.5
C5—C6—H6	119.0	O6—C15—H15C	109.5
C1—C6—H6	119.0	H15A—C15—H15C	109.5
N2—C7—C8	121.3 (3)	H15B—C15—H15C	109.5
N2—C7—H7	119.4		
C1—N1—N2—C7	-179.1 (2)	C2—C1—C6—C5	2.0 (4)
N2—N1—C1—C6	0.9 (4)	N1—N2—C7—C8	-179.8 (2)
N2—N1—C1—C2	179.5 (2)	N2—C7—C8—C13	-5.0 (4)
N1—C1—C2—C3	179.7 (2)	N2—C7—C8—C9	175.9 (2)
C6—C1—C2—C3	-1.7 (4)	C14—O5—C9—C8	129.0 (3)
N1—C1—C2—N3	-0.4 (4)	C14—O5—C9—C10	-57.3 (4)
C6—C1—C2—N3	178.2 (2)	C13—C8—C9—O5	174.7 (2)

supplementary materials


O2—N3—C2—C3	4.9 (4)	C7—C8—C9—O5	-6.3 (4)
O1—N3—C2—C3	-173.6 (3)	C13—C8—C9—C10	0.9 (4)
O2—N3—C2—C1	-174.9 (3)	C7—C8—C9—C10	180.0 (2)
O1—N3—C2—C1	6.6 (4)	C15—O6—C10—C11	-7.7 (4)
C1—C2—C3—C4	0.6 (4)	C15—O6—C10—C9	173.9 (3)
N3—C2—C3—C4	-179.3 (2)	O5—C9—C10—O6	3.7 (4)
C2—C3—C4—C5	0.3 (4)	C8—C9—C10—O6	177.2 (2)
C2—C3—C4—N4	-179.8 (2)	O5—C9—C10—C11	-174.8 (2)
O4—N4—C4—C3	175.4 (3)	C8—C9—C10—C11	-1.4 (4)
O3—N4—C4—C3	-5.5 (4)	O6—C10—C11—C12	-177.0 (3)
O4—N4—C4—C5	-4.7 (4)	C9—C10—C11—C12	1.3 (4)
O3—N4—C4—C5	174.4 (3)	C10—C11—C12—C13	-0.9 (5)
C3—C4—C5—C6	0.0 (4)	C11—C12—C13—C8	0.4 (5)
N4—C4—C5—C6	-179.9 (2)	C9—C8—C13—C12	-0.5 (4)
C4—C5—C6—C1	-1.2 (4)	C7—C8—C13—C12	-179.5 (3)
N1—C1—C6—C5	-179.3 (2)		

Hydrogen-bond geometry (\AA , °)

$D\cdots H$	$D—H$	$H\cdots A$	$D\cdots A$	$D—H\cdots A$
N1—H1···O1	0.86	1.99	2.625 (4)	130
N1—H1···N3	0.86	2.60	2.913 (4)	103
C3—H3···O2	0.93	2.33	2.654 (4)	100
C6—H6···N2	0.93	2.44	2.766 (4)	100
C7—H7···O5	0.93	2.41	2.737 (4)	101
C14—H14A···O4 ⁱ	0.96	2.48	3.431 (4)	170

Symmetry codes: (i) $-x+1, -y, -z+1$.

Fig. 1



supplementary materials

Fig. 2

