

Diethyl 4,6-diacetamidoisophthalate

Peishen Li, Xianghui Li, Chao Chen, Lihua Yuan and Wen Feng*

Key Laboratory for Radiation Physics and Technology of the Ministry of Education,
 College of Chemistry, Institute of Nuclear Science and Technology, Sichuan
 University, Chengdu 610064, Sichuan, People's Republic of China
 Correspondence e-mail: wfeng9510@scu.edu.cn


Received 26 March 2011; accepted 12 April 2011

Key indicators: single-crystal X-ray study; $T = 292\text{ K}$; mean $\sigma(\text{C}-\text{C}) = 0.005\text{ \AA}$;
 R factor = 0.079; wR factor = 0.268; data-to-parameter ratio = 13.9.

In the title compound, $\text{C}_{16}\text{H}_{20}\text{N}_2\text{O}_6$, two intramolecular N—H···O hydrogen bonds occur, in which the carbonyl O atoms of the ethyl acetate groups serve as the acceptor atoms; both motifs generate $S(6)$ rings. In the crystal, molecules are linked by weak C—H···O links (with the acceptor O atoms part of the amide groups), generating [001] chains.

Related literature

For background to intramolecular hydrogen bonds this class of compound, see: Zhu *et al.* (2000); Yuan *et al.* (2004); Feng *et al.* (2009); Yan *et al.* (2010); Zhang *et al.* (2008). For a related structure, see: Zhang *et al.* (2006).



Experimental

Crystal data

$\text{C}_{16}\text{H}_{20}\text{N}_2\text{O}_6$	$\gamma = 76.50(2)^\circ$
$M_r = 336.34$	$V = 843.7(5)\text{ \AA}^3$
Triclinic, $P\bar{1}$	$Z = 2$
$a = 7.951(3)\text{ \AA}$	Mo $K\alpha$ radiation
$b = 10.249(3)\text{ \AA}$	$\mu = 0.10\text{ mm}^{-1}$
$c = 11.109(4)\text{ \AA}$	$T = 292\text{ K}$
$\alpha = 76.70(3)^\circ$	$0.50 \times 0.46 \times 0.40\text{ mm}$
$\beta = 77.42(3)^\circ$	

Data collection

Enraf–Nonius CAD-4
 diffractometer
 3149 measured reflections
 3094 independent reflections
 1826 reflections with $I > 2\sigma(I)$

$R_{\text{int}} = 0.004$
 3 standard reflections every 150
 reflections
 intensity decay: 4.5%

Refinement

$R[F^2 > 2\sigma(F^2)] = 0.079$
 $wR(F^2) = 0.268$
 $S = 1.09$
 3094 reflections
 223 parameters
 4 restraints

H atoms treated by a mixture of
 independent and constrained
 refinement
 $\Delta\rho_{\text{max}} = 0.59\text{ e \AA}^{-3}$
 $\Delta\rho_{\text{min}} = -0.65\text{ e \AA}^{-3}$

Table 1
 Hydrogen-bond geometry (\AA , $^\circ$).

$D-\text{H}\cdots A$	$D-\text{H}$	$\text{H}\cdots A$	$D\cdots A$	$D-\text{H}\cdots A$
N1—H1N···O2	0.88 (4)	1.92 (3)	2.676 (4)	144 (3)
N2—H2N···O6 ⁱ	0.81 (4)	1.96 (4)	2.656 (4)	143 (3)
C9—H9B···O3 ^j	0.96	2.54	3.445 (6)	157
C16—H16A···O4 ⁱⁱ	0.96	2.57	3.485 (7)	160

Symmetry codes: (i) $x, y, z - 1$; (ii) $-x + 1, -y, -z + 2$.

Data collection: *DIFRAC* (Gabe *et al.*, 1993); cell refinement: *DIFRAC*; data reduction: *NRCVAX* (Gabe *et al.*, 1989); program(s) used to solve structure: *SHELXS97* (Sheldrick, 2008); program(s) used to refine structure: *SHELXL97* (Sheldrick, 2008); molecular graphics: *ORTEP-3* (Farrugia, 1997); software used to prepare material for publication: *SHELXL97*.

The authors acknowledge the National Natural Science Foundation of China (20774059) for funding this work, and the Analytical & Testing Center of Sichuan University for the X-ray analysis.

Supplementary data and figures for this paper are available from the IUCr electronic archives (Reference: HB5828).

References

- Farrugia, L. J. (1997). *J. Appl. Cryst.* **30**, 565.
- Feng, W., Yamato, K., Yang, L. Q., Ferguson, J. S., Zhong, L. J., Zou, S. L., Yuan, L. H., Zeng, X. C. & Gong, B. (2009). *J. Am. Chem. Soc.* **131**, 2629–2637.
- Gabe, E. J., Le Page, Y., Charland, J.-P., Lee, F. L. & White, P. S. (1989). *J. Appl. Cryst.* **22**, 384–387.
- Gabe, E. J., White, P. S. & Enright, G. D. (1993). *DIFRAC*. American Crystallographic Association, Pittsburgh Meeting Abstract, PA 104.
- Sheldrick, G. M. (2008). *Acta Cryst. A* **64**, 112–122.
- Yan, Y., Qin, B., Ren, C. L., Chen, X. Y., Yip, Y. K., Ye, R. J., Zhang, D. W., Su, H. B. & Zeng, H. Q. (2010). *J. Am. Chem. Soc.* **132**, 5869–5879.
- Yuan, L. H., Feng, W., Yamato, K., Sanford, A. R., Xu, D. G., Guo, H. & Gong, B. (2004). *J. Am. Chem. Soc.* **126**, 11120–11121.
- Zhang, A. M., Han, Y. H., Yamato, K., Zeng, X. C. & Gong, B. (2006). *Org. Lett.* **8**, 803–806.
- Zhang, Y. F., Yamato, K., Zhong, K., Zhu, J., Deng, J. G. & Gong, B. (2008). *Org. Lett.* **10**, 4338–4342.
- Zhu, J., Parra, R. D., Zeng, H. Q., Skrzypczak-Jankun, E., Zeng, X. C. & Gong, B. (2000). *J. Am. Chem. Soc.* **122**, 4219–4220.

supplementary materials

Acta Cryst. (2011). E67, o1185 [doi:10.1107/S1600536811013845]

Diethyl 4,6-diacetamidoisophthalate

P. Li, X. Li, C. Chen, L. Yuan and W. Feng

Comment

Hydrogen bonds play a vital role in assisting formation of aromatic oligoamide foldmers and macrocycles (Zhu *et al.*, 2000; Yuan *et al.*, 2004; Feng *et al.*, 2009). It is important to examine the structural feature of the backbones that is responsible for the formation of foldmers and macrocycles. Research has focused on the role of intramolecular hydrogen bonds in regulating conformations and interactions both in solution and the solid state (Yan *et al.*, 2010; Zhang *et al.* 2008). Herein, we report on the crystal structure of the aromatic monomer containing intramolecular hydrogen bonds, which could be applied as important building blocks to construct macrocycles.

The title compound comprises two similar six-member hydrogen bonds. There are two possible conformations: (a) one with the ethoxy O atoms involving in the intramolecular NH···OC₂H₅ six-member hydrogen bonds (conformation **a**), and (b) the other with carbonyl O atoms that form intramolecular H-bonds (conformation **b**) (Fig. 1). However, the crystal structure revealed the existence of two six-member hydrogen bonds that involve carbonyl O atoms, indicating that conformation **b** is more stable and is sustained by two intramolecular hydrogen bond N1H1···O2 (Table 1, Fig. 2). It is expected that this type of hydrogen bonds may be exploited to construct macrocycles and supamolecular architecture with folded conformations. In the crystal, the molecules are linked by C—H···O interactions.

Experimental

For synthesis of the title compound, Pd/C (50 mg) was added to the solution of diethyl 4,6-dinitroisophthalate (500 mg, 1.60 mmol) in CH₂Cl₂ and the mixture was stirred at room temperature under H₂ atmosphere for 6 h. After removal of Pd/C and solvent, the white solid was obtained and then dissolved in CH₂Cl₂ (50 ml). Et₃N (335 mg, 3.31 mmol) was added to the above solution followed by dropping acetyl chloride (350 mg, 4.46 mmol). After stirring 2 h at room temperature, the mixture was washed with distilled water, dried over anhydrous Na₂SO₄. Removal of solvent under reduced pressure gave the crude product, which was recrystallized from methanol to yield the white product (430 mg, yield 79.8%). ¹H NMR (400 MHz, CDCl₃): δ 10.95 (s, 2H), 9.98 (s, 1H), 8.74 (s, 1H), 4.35 (m, 4H), 2.25 (s, 6H), 1.20 (t, J = 6.4 Hz, 6H).

The crystal of the title compound was grown from the THF solution. This compound was dissolved in THF and filtered, then the filtrate was allowed to stand undisturbed to give the single-crystal.

Refinement

The H atoms bonded to atom N1 and N2 of the title compound were located from difference Fourier maps and refined isotropically [N1—H = 1.88 (4), N2—H = 0.81 (4) Å]. H atoms bonded to C atoms were placed in calculated positions and refined using a riding model, with C—H = 0.9300 Å and *U*_{iso}(H) = 1.2Ueq(C) for aryl H atoms, C—H = 0.9700 Å and *U*_{iso}(H) = 1.2Ueq(C) for methylene H atoms, C—H = 0.9600 Å and *U*_{iso}(H) = 1.5Ueq(C) for methyl H atoms. The deepest difference hole of -0.65 e.Å⁻³ is 0.33 Å from atom C16.

supplementary materials

Figures




Fig. 1. The two possible intramolecular hydrogen bond conformations of the title compound.




Fig. 2. A view of the title compound with displacement ellipsoids shown at the 50% probability level. Hydrogen bonds are indicated by broken lines.

Diethyl 4,6-diacetamidoisophthalate

Crystal data

C ₁₆ H ₂₀ N ₂ O ₆	Z = 2
M _r = 336.34	F(000) = 356
Triclinic, PT	D _x = 1.324 Mg m ⁻³
a = 7.951 (3) Å	Mo K α radiation, λ = 0.71073 Å
b = 10.249 (3) Å	Cell parameters from 20 reflections
c = 11.109 (4) Å	θ = 5.4–6.7°
α = 76.70 (3)°	μ = 0.10 mm ⁻¹
β = 77.42 (3)°	T = 292 K
γ = 76.50 (2)°	Block, colourless
V = 843.7 (5) Å ³	0.50 × 0.46 × 0.40 mm

Data collection

Enraf–Nonius CAD-4 diffractometer	R _{int} = 0.004
Radiation source: fine-focus sealed tube graphite	$\theta_{\text{max}} = 25.5^\circ$, $\theta_{\text{min}} = 1.9^\circ$
$\omega/2\theta$ scans	$h = -9 \rightarrow 9$
3149 measured reflections	$k = -1 \rightarrow 12$
3094 independent reflections	$l = -12 \rightarrow 13$
1826 reflections with $I > 2\sigma(I)$	3 standard reflections every 150 reflections intensity decay: 4.5%

Refinement

Refinement on F^2	Primary atom site location: structure-invariant direct methods
Least-squares matrix: full	Secondary atom site location: difference Fourier map
$R[F^2 > 2\sigma(F^2)] = 0.079$	Hydrogen site location: inferred from neighbouring sites

$wR(F^2) = 0.268$	H atoms treated by a mixture of independent and constrained refinement
$S = 1.09$	$w = 1/[\sigma^2(F_o^2) + (0.1639P)^2 + 0.1078P]$
3094 reflections	where $P = (F_o^2 + 2F_c^2)/3$
223 parameters	$(\Delta/\sigma)_{\text{max}} < 0.001$
4 restraints	$\Delta\rho_{\text{max}} = 0.59 \text{ e \AA}^{-3}$
	$\Delta\rho_{\text{min}} = -0.65 \text{ e \AA}^{-3}$

Special details

Geometry. All e.s.d.'s (except the e.s.d. in the dihedral angle between two l.s. planes) are estimated using the full covariance matrix. The cell e.s.d.'s are taken into account individually in the estimation of e.s.d.'s in distances, angles and torsion angles; correlations between e.s.d.'s in cell parameters are only used when they are defined by crystal symmetry. An approximate (isotropic) treatment of cell e.s.d.'s is used for estimating e.s.d.'s involving l.s. planes.

Refinement. Refinement of F^2 against ALL reflections. The weighted R -factor wR and goodness of fit S are based on F^2 , conventional R -factors R are based on F , with F set to zero for negative F^2 . The threshold expression of $F^2 > \sigma(F^2)$ is used only for calculating R -factors(gt) etc. and is not relevant to the choice of reflections for refinement. R -factors based on F^2 are statistically about twice as large as those based on F , and R -factors based on ALL data will be even larger.

Fractional atomic coordinates and isotropic or equivalent isotropic displacement parameters (\AA^2)

	x	y	z	$U_{\text{iso}}^*/U_{\text{eq}}$
O1	0.9467 (4)	-0.1465 (3)	0.6464 (2)	0.0692 (9)
O2	1.1334 (4)	-0.2976 (3)	0.7586 (2)	0.0706 (9)
O3	1.0473 (5)	-0.3487 (3)	1.2179 (3)	0.0988 (12)
O4	0.7415 (4)	-0.0447 (3)	1.3243 (3)	0.0796 (9)
O5	0.5572 (4)	0.2164 (3)	0.7774 (3)	0.0692 (8)
O6	0.5009 (4)	0.2628 (3)	0.9676 (3)	0.0718 (8)
N1	1.0637 (4)	-0.3063 (3)	1.0068 (3)	0.0503 (8)
H1N	1.113 (4)	-0.338 (4)	0.938 (3)	0.046 (9)*
N2	0.6413 (4)	0.0776 (3)	1.1468 (3)	0.0514 (8)
H2N	0.581 (5)	0.150 (4)	1.120 (3)	0.044 (10)*
C1	0.9150 (4)	-0.1332 (4)	0.8583 (3)	0.0477 (8)
C2	0.9442 (4)	-0.1847 (3)	0.9824 (3)	0.0427 (8)
C3	0.8541 (4)	-0.1138 (3)	1.0777 (3)	0.0440 (8)
H3	0.8745	-0.1477	1.1594	0.053*
C4	0.7336 (4)	0.0075 (3)	1.0518 (3)	0.0413 (8)
C5	0.7021 (4)	0.0591 (3)	0.9291 (3)	0.0449 (8)
C6	0.7941 (4)	-0.0127 (3)	0.8356 (3)	0.0460 (8)
H6	0.7736	0.0216	0.7539	0.055*
C7	1.0103 (5)	-0.2022 (4)	0.7527 (3)	0.0539 (9)
C8	1.0398 (7)	-0.1987 (5)	0.5319 (4)	0.0871 (15)
H8A	1.1655	-0.2192	0.5310	0.104*
H8B	1.0160	-0.1303	0.4581	0.104*
C9	0.9772 (8)	-0.3269 (6)	0.5304 (5)	0.1008 (17)
H9A	1.0175	-0.3987	0.5961	0.151*
H9B	1.0233	-0.3543	0.4507	0.151*

supplementary materials

H9C	0.8512	-0.3090	0.5434	0.151*
C10	1.1082 (5)	-0.3819 (4)	1.1169 (4)	0.0574 (10)
C11	1.2385 (6)	-0.5100 (4)	1.1039 (4)	0.0680 (11)
H11A	1.2382	-0.5696	1.1844	0.102*
H11B	1.3533	-0.4888	1.0722	0.102*
H11C	1.2081	-0.5544	1.0466	0.102*
C12	0.6424 (5)	0.0486 (4)	1.2735 (3)	0.0544 (9)
C13	0.5091 (6)	0.1480 (5)	1.3433 (4)	0.0723 (12)
H13A	0.3951	0.1530	1.3249	0.108*
H13B	0.5399	0.2367	1.3177	0.108*
H13C	0.5073	0.1177	1.4320	0.108*
C14	0.5778 (5)	0.1888 (4)	0.8965 (4)	0.0531 (9)
C15	0.4366 (9)	0.3443 (6)	0.7425 (5)	0.1109 (14)
H15A	0.4848	0.4216	0.7463	0.133*
H15B	0.3245	0.3465	0.7987	0.133*
C16	0.4146 (9)	0.3488 (6)	0.6092 (5)	0.1109 (14)
H16A	0.3796	0.2666	0.6051	0.166*
H16B	0.5239	0.3565	0.5534	0.166*
H16C	0.3262	0.4263	0.5848	0.166*

Atomic displacement parameters (\AA^2)

	U^{11}	U^{22}	U^{33}	U^{12}	U^{13}	U^{23}
O1	0.0823 (19)	0.0772 (19)	0.0456 (15)	0.0082 (15)	-0.0164 (13)	-0.0255 (13)
O2	0.0655 (17)	0.0820 (19)	0.0598 (17)	0.0178 (15)	-0.0127 (13)	-0.0357 (14)
O3	0.130 (3)	0.084 (2)	0.0577 (19)	0.040 (2)	-0.0244 (18)	-0.0185 (16)
O4	0.094 (2)	0.082 (2)	0.0529 (16)	0.0193 (18)	-0.0208 (15)	-0.0222 (14)
O5	0.0774 (19)	0.0613 (16)	0.0625 (17)	0.0094 (14)	-0.0272 (14)	-0.0076 (13)
O6	0.083 (2)	0.0520 (15)	0.0749 (19)	0.0166 (14)	-0.0225 (15)	-0.0227 (14)
N1	0.0532 (17)	0.0460 (16)	0.0500 (17)	0.0046 (13)	-0.0107 (14)	-0.0184 (13)
N2	0.0526 (18)	0.0501 (18)	0.0507 (18)	0.0036 (15)	-0.0115 (14)	-0.0189 (14)
C1	0.0476 (19)	0.0513 (19)	0.0465 (19)	-0.0098 (16)	-0.0054 (15)	-0.0161 (15)
C2	0.0392 (17)	0.0438 (17)	0.0469 (18)	-0.0060 (14)	-0.0068 (14)	-0.0147 (14)
C3	0.0455 (18)	0.0449 (18)	0.0428 (18)	-0.0052 (15)	-0.0074 (14)	-0.0147 (14)
C4	0.0353 (16)	0.0430 (17)	0.0480 (18)	-0.0066 (14)	-0.0029 (13)	-0.0182 (14)
C5	0.0436 (18)	0.0433 (18)	0.0492 (19)	-0.0035 (15)	-0.0102 (15)	-0.0143 (15)
C6	0.0485 (19)	0.0480 (18)	0.0430 (17)	-0.0067 (15)	-0.0100 (14)	-0.0120 (14)
C7	0.055 (2)	0.058 (2)	0.051 (2)	-0.0040 (18)	-0.0062 (16)	-0.0240 (17)
C8	0.105 (4)	0.096 (3)	0.054 (2)	0.007 (3)	-0.009 (2)	-0.032 (2)
C9	0.121 (4)	0.107 (4)	0.081 (3)	0.002 (3)	-0.026 (3)	-0.048 (3)
C10	0.058 (2)	0.051 (2)	0.062 (2)	0.0021 (17)	-0.0145 (18)	-0.0186 (18)
C11	0.076 (3)	0.048 (2)	0.077 (3)	0.0043 (19)	-0.021 (2)	-0.0148 (19)
C12	0.056 (2)	0.060 (2)	0.050 (2)	-0.0067 (18)	-0.0058 (17)	-0.0237 (17)
C13	0.071 (3)	0.081 (3)	0.063 (2)	0.005 (2)	-0.005 (2)	-0.035 (2)
C14	0.058 (2)	0.0453 (19)	0.058 (2)	-0.0045 (17)	-0.0150 (17)	-0.0146 (16)
C15	0.130 (3)	0.087 (3)	0.103 (3)	0.021 (2)	-0.050 (3)	-0.008 (2)
C16	0.130 (3)	0.087 (3)	0.103 (3)	0.021 (2)	-0.050 (3)	-0.008 (2)

Geometric parameters (Å, °)

O1—C7	1.338 (4)	C5—C14	1.482 (5)
O1—C8	1.474 (5)	C6—H6	0.9300
O2—C7	1.212 (4)	C8—C9	1.514 (6)
O3—C10	1.212 (5)	C8—H8A	0.9700
O4—C12	1.204 (5)	C8—H8B	0.9700
O5—C14	1.326 (4)	C9—H9A	0.9600
O5—C15	1.459 (5)	C9—H9B	0.9600
O6—C14	1.192 (4)	C9—H9C	0.9600
N1—C10	1.359 (5)	C10—C11	1.485 (5)
N1—C2	1.390 (4)	C11—H11A	0.9600
N1—H1N	0.88 (4)	C11—H11B	0.9600
N2—C12	1.371 (5)	C11—H11C	0.9600
N2—C4	1.391 (4)	C12—C13	1.505 (5)
N2—H2N	0.81 (4)	C13—H13A	0.9600
C1—C6	1.386 (5)	C13—H13B	0.9600
C1—C2	1.408 (5)	C13—H13C	0.9600
C1—C7	1.482 (5)	C15—C16	1.517 (7)
C2—C3	1.393 (4)	C15—H15A	0.9700
C3—C4	1.395 (4)	C15—H15B	0.9700
C3—H3	0.9300	C16—H16A	0.9600
C4—C5	1.398 (5)	C16—H16B	0.9600
C5—C6	1.388 (4)	C16—H16C	0.9600
C7—O1—C8	117.5 (3)	H9A—C9—H9B	109.5
C14—O5—C15	114.2 (3)	C8—C9—H9C	109.5
C10—N1—C2	130.8 (3)	H9A—C9—H9C	109.5
C10—N1—H1N	117 (2)	H9B—C9—H9C	109.5
C2—N1—H1N	112 (2)	O3—C10—N1	123.3 (3)
C12—N2—C4	131.2 (3)	O3—C10—C11	122.3 (4)
C12—N2—H2N	116 (3)	N1—C10—C11	114.4 (3)
C4—N2—H2N	112 (3)	C10—C11—H11A	109.5
C6—C1—C2	118.0 (3)	C10—C11—H11B	109.5
C6—C1—C7	119.7 (3)	H11A—C11—H11B	109.5
C2—C1—C7	122.3 (3)	C10—C11—H11C	109.5
N1—C2—C3	121.5 (3)	H11A—C11—H11C	109.5
N1—C2—C1	118.7 (3)	H11B—C11—H11C	109.5
C3—C2—C1	119.9 (3)	O4—C12—N2	124.0 (3)
C2—C3—C4	120.7 (3)	O4—C12—C13	122.9 (4)
C2—C3—H3	119.6	N2—C12—C13	113.1 (3)
C4—C3—H3	119.6	C12—C13—H13A	109.5
N2—C4—C3	121.0 (3)	C12—C13—H13B	109.5
N2—C4—C5	119.0 (3)	H13A—C13—H13B	109.5
C3—C4—C5	120.0 (3)	C12—C13—H13C	109.5
C6—C5—C4	118.3 (3)	H13A—C13—H13C	109.5
C6—C5—C14	119.7 (3)	H13B—C13—H13C	109.5
C4—C5—C14	122.0 (3)	O6—C14—O5	121.7 (3)
C1—C6—C5	123.0 (3)	O6—C14—C5	125.2 (3)

supplementary materials


C1—C6—H6	118.5	O5—C14—C5	113.1 (3)
C5—C6—H6	118.5	O5—C15—C16	106.1 (4)
O2—C7—O1	122.4 (3)	O5—C15—H15A	110.5
O2—C7—C1	125.2 (3)	C16—C15—H15A	110.5
O1—C7—C1	112.4 (3)	O5—C15—H15B	110.5
O1—C8—C9	108.6 (4)	C16—C15—H15B	110.5
O1—C8—H8A	110.0	H15A—C15—H15B	108.7
C9—C8—H8A	110.0	C15—C16—H16A	109.5
O1—C8—H8B	110.0	C15—C16—H16B	109.5
C9—C8—H8B	110.0	H16A—C16—H16B	109.5
H8A—C8—H8B	108.3	C15—C16—H16C	109.5
C8—C9—H9A	109.5	H16A—C16—H16C	109.5
C8—C9—H9B	109.5	H16B—C16—H16C	109.5
C10—N1—C2—C3	4.6 (6)	C14—C5—C6—C1	178.7 (3)
C10—N1—C2—C1	−175.5 (3)	C8—O1—C7—O2	−4.3 (6)
C6—C1—C2—N1	179.3 (3)	C8—O1—C7—C1	175.0 (3)
C7—C1—C2—N1	−1.5 (5)	C6—C1—C7—O2	170.3 (4)
C6—C1—C2—C3	−0.8 (5)	C2—C1—C7—O2	−8.9 (6)
C7—C1—C2—C3	178.4 (3)	C6—C1—C7—O1	−9.1 (5)
N1—C2—C3—C4	−179.5 (3)	C2—C1—C7—O1	171.8 (3)
C1—C2—C3—C4	0.6 (5)	C7—O1—C8—C9	83.1 (5)
C12—N2—C4—C3	−1.0 (6)	C2—N1—C10—O3	−2.1 (7)
C12—N2—C4—C5	178.0 (3)	C2—N1—C10—C11	177.7 (3)
C2—C3—C4—N2	179.0 (3)	C4—N2—C12—O4	6.1 (7)
C2—C3—C4—C5	0.0 (5)	C4—N2—C12—C13	−175.2 (3)
N2—C4—C5—C6	−179.4 (3)	C15—O5—C14—O6	1.0 (6)
C3—C4—C5—C6	−0.4 (5)	C15—O5—C14—C5	−179.4 (4)
N2—C4—C5—C14	2.1 (5)	C6—C5—C14—O6	−174.9 (4)
C3—C4—C5—C14	−178.9 (3)	C4—C5—C14—O6	3.6 (6)
C2—C1—C6—C5	0.4 (5)	C6—C5—C14—O5	5.6 (5)
C7—C1—C6—C5	−178.8 (3)	C4—C5—C14—O5	−176.0 (3)
C4—C5—C6—C1	0.2 (5)	C14—O5—C15—C16	−173.5 (4)

Hydrogen-bond geometry (\AA , °)

$D\text{—H}\cdots A$	$D\text{—H}$	$H\cdots A$	$D\cdots A$	$D\text{—H}\cdots A$
N1—H1N···O2	0.88 (4)	1.92 (3)	2.676 (4)	144 (3)
N2—H2N···O6	0.81 (4)	1.96 (4)	2.656 (4)	143 (3)
C9—H9B···O3 ⁱ	0.96	2.54	3.445 (6)	157
C16—H16A···O4 ⁱⁱ	0.96	2.57	3.485 (7)	160

Symmetry codes: (i) $x, y, z-1$; (ii) $-x+1, -y, -z+2$.

Fig. 1



supplementary materials

Fig. 2

