

Acta Crystallographica Section E

Structure Reports

Online

ISSN 1600-5368

(2E)-2-[2-(Cyclohexylcarbamothioyl)-hydrazinylidene]propanoic acidMd. Abu Affan,^{a,‡} Md. Abdus Salam,^a Fasihuddin B. Ahmad,^a Seik Weng Ng^b and Edward R. T. Tiekink^{b*}^aFaculty of Resource Science and Technology, Universiti Malaysia Sarawak, 94300 Kota Samarahan, Sarawak, Malaysia, and ^bDepartment of Chemistry, University of Malaya, 50603 Kuala Lumpur, Malaysia

Correspondence e-mail: edward.tiekink@gmail.com


Received 16 April 2011; accepted 18 April 2011

Key indicators: single-crystal X-ray study; $T = 100$ K; mean $\sigma(\text{C}-\text{C}) = 0.003$ Å; R factor = 0.041; wR factor = 0.143; data-to-parameter ratio = 17.2.

In the title thiourea derivative, $\text{C}_{10}\text{H}_{17}\text{N}_3\text{O}_2\text{S}$, the carboxyl group and the least-squares plane through the cyclohexyl ring are twisted out of the plane through the central CN_3S residue; the respective dihedral angles are 7.18 (8) and 62.29 (4)°. The conformation about the azomethine bond [1.275 (2) Å] is *E*. The NH groups are *anti*, with one forming an intramolecular $\text{N}-\text{H}\cdots\text{N}$ hydrogen bond. The main feature of the crystal structure is the formation of linear supramolecular chains along [110] mediated by alternating pairs of $\text{O}-\text{H}\cdots\text{O}$ and pairs of $\text{N}-\text{H}\cdots\text{S}$ hydrogen bonds.

Related literature

For related thiourea structures, see: Normaya *et al.* (2011); Salam *et al.* (2011).



Experimental

Crystal data

 $\text{C}_{10}\text{H}_{17}\text{N}_3\text{O}_2\text{S}$ $M_r = 243.33$ Monoclinic, $P2_1/n$ $a = 8.9204$ (2) Å $b = 6.0350$ (2) Å $c = 22.4750$ (6) Å $\beta = 90.051$ (3)° $V = 1209.93$ (6) Å³ $Z = 4$ Mo $K\alpha$ radiation $\mu = 0.26$ mm⁻¹ $T = 100$ K $0.30 \times 0.20 \times 0.10$ mm

Data collection

Agilent Supernova Dual diffractometer with an Atlas detector

Absorption correction: multi-scan (*CrysAlis PRO*; Agilent, 2010) $T_{\min} = 0.687$, $T_{\max} = 1.000$

7394 measured reflections

2723 independent reflections

2251 reflections with $I > 2\sigma(I)$ $R_{\text{int}} = 0.036$

Refinement

 $R[F^2 > 2\sigma(F^2)] = 0.041$ $wR(F^2) = 0.143$ $S = 0.98$

2723 reflections

158 parameters

3 restraints

H atoms treated by a mixture of independent and constrained refinement

 $\Delta\rho_{\text{max}} = 0.36$ e Å⁻³ $\Delta\rho_{\text{min}} = -0.32$ e Å⁻³

Table 1

Hydrogen-bond geometry (Å, °).

$D-\text{H}\cdots A$	$D-\text{H}$	$\text{H}\cdots A$	$D\cdots A$	$D-\text{H}\cdots A$
$\text{N3}-\text{H3}\cdots\text{N1}$	0.87 (1)	2.20 (2)	2.574 (2)	106 (2)
$\text{O1}-\text{H1}\cdots\text{O2}^{\text{i}}$	0.85 (1)	1.80 (1)	2.6416 (17)	172 (3)
$\text{N2}-\text{H2}\cdots\text{S1}^{\text{ii}}$	0.89 (1)	2.65 (1)	3.5384 (15)	176 (2)

Symmetry codes: (i) $-x + 1, -y + 1, -z + 1$; (ii) $-x + 2, -y + 3, -z + 1$.

Data collection: *CrysAlis PRO* (Agilent, 2010); cell refinement: *CrysAlis PRO*; data reduction: *CrysAlis PRO*; program(s) used to solve structure: *SHELXS97* (Sheldrick, 2008); program(s) used to refine structure: *SHELXL97* (Sheldrick, 2008); molecular graphics: *ORTEP-3* (Farrugia, 1997) and *DIAMOND* (Brandenburg, 2006); software used to prepare material for publication: *PUBLICIF* (Westrip, 2010).

This work was supported financially by the Ministry of Science, Technology and Innovation (MOSTI) under a research grant (No. 06-01-09-SF0046). The authors thank Universiti Malaysia Sarawak (UNIMAS) for the facilities to carry out the research work. The authors also thank the University of Malaya for support of the crystallographic facility.

Supplementary data and figures for this paper are available from the IUCr electronic archives (Reference: HB5851).

References

- Agilent (2010). *CrysAlis PRO*. Agilent Technologies, Yarnton, Oxfordshire, England.
- Brandenburg, K. (2006). *DIAMOND*. Crystal Impact GbR, Bonn, Germany.
- Farrugia, L. J. (1997). *J. Appl. Cryst.* **30**, 565.
- Normaya, E., Farina, Y., Halim, S. N. A. & Tiekink, E. R. T. (2011). *Acta Cryst. E* **67**, o943–o944.
- Salam, M. A., Affan, M. A., Ahmad, F. B., Ng, S. W. & Tiekink, E. R. T. (2011). *Acta Cryst. E* **67**, o955.
- Sheldrick, G. M. (2008). *Acta Cryst. A* **64**, 112–122.
- Westrip, S. P. (2010). *J. Appl. Cryst.* **43**, 920–925.

‡ Additional correspondence author, e-mail: maaffan@yahoo.com.

supplementary materials

Acta Cryst. (2011). E67, o1193 [doi:10.1107/S1600536811014449]

(2E)-2-[2-(Cyclohexylcarbamothioyl)hydrazinylidene]propanoic acid

M. A. Affan, M. A. Salam, F. B. Ahmad, S. W. Ng and E. R. T. Tiekink

Comment

In continuation of structural investigations into conformation and hydrogen bonding in thiourea derivatives (Normaya *et al.* 2011; Salam *et al.*, 2011), the title compound, (I), was investigated.

The central CN₃S chromophore in (I), Fig. 1, is planar (r.m.s. = 0.0039 Å). The carboxylate residue is slightly twisted out this plane (dihedral angle = 7.18 (8) °), and, by contrast, the least-squares plane through the cyclohexyl group (which has the conformation of a chair) is twisted significantly out of the central plane (dihedral angle = 62.29 (4) °). The H atoms of the NH groups are *anti*, and the conformation about the azomethine bond [1.275 (2) Å] is *E*. The N3—H forms an intramolecular hydrogen bond with the imino-N3 atom, Table 1. Finally, the thione and carboxylic acid groups lie to opposite sides of the molecule. This arrangements enables the formation of linear supramolecular chains *via* hydrogen bonds along [110] in the crystal packing, Fig. 2 and Table 1. The carboxylic acid residues self-associate *via* a centrosymmetric eight-membered {···HOC=O}₂ synthon. Similarly, the thiourea entity with the NH atom not involved in the intramolecular N—H···N interaction, self-associates *via* a centrosymmetric eight-membered {···HNC=S}₂ synthon. Chains lie in the *ab* plane with the cyclohexyl rings inter-digitating along the *c* axis, Fig. 3.

Experimental

Cyclohexylisothiocyanate (0.706 g, 5 mmol) and hydrazine hydrate (0.250 g, 5 mmol), each dissolved in 10 ml ethanol were mixed with constant stirring. The stirring was continued for 30 min and the white product that formed, *N*(4)-cyclohexylthiosemicarbazide, was washed with ethanol and dried *in vacuo*. A solution of this (0.51 g, 3 mmol) in 10 ml methanol was refluxed with a methanolic solution of pyruvic acid (0.261 g, 3 mmol) for 5 h after the addition of 1–2 drops of glacial acetic acid. On cooling the solution to room temperature, white precipitate separated, which were filtered and washed with methanol. The white precipitate was recrystallized from methanol to yield colourless prisms and dried *in vacuo* over silica gel. (*M.pt.* 465–467 K. Yield 0.621 g (80%). Anal. Found: C, 49.31; H, 7.01; N, 17.18%. C₁₀H₁₇N₃O₂S requires: C, 49.36; H, 7.04; N, 17.26%. FT—IR (KBr, cm⁻¹) ν_{\max} : 3322 (m, OH), 3197 (s, NH), 2922, 2851 (s, cyclohexyl), 1692 (m, C=O), 1619 (w, C=N), 980 (m, N—N), 1249, 873 (w, C=S).

Refinement

Carbon-bound H-atoms were placed in calculated positions (C—H = 0.98 to 1.00 Å) and were included in the refinement in the riding model approximation, with $U_{\text{iso}}(\text{H})$ set to 1.2–1.5 $U_{\text{eq}}(\text{C})$. The O- and N-bound H-atoms were located in a difference Fourier map and were refined with distance restraints of O—H = 0.84±0.01 Å and N—H 0.88±0.01 Å, and with $U_{\text{iso}}(\text{H}) = yU_{\text{eq}}(\text{N})$ for $y = 1.5$ (O) and 1.2 (N).

Figures




Fig. 1. The molecular structure of (I) showing displacement ellipsoids at the 50% probability level.




Fig. 2. A view of the supramolecular chain aligned along [110] in (I). The O—H...O and N—H...S hydrogen bonds are shown as orange and blue dashed lines, respectively.




Fig. 3. A view in projection down the *b* axis of the crystal packing in (I) showing the inter-digitation of the cyclohexyl rings along the *c* direction. The O—H...O and N—H...S hydrogen bonds are shown as orange and blue dashed lines, respectively.

(2*E*)-2-[2-(Cyclohexylcarbamothioyl)hydrazinylidene]propanoic acid

Crystal data

$C_{10}H_{17}N_3O_2S$

$M_r = 243.33$

Monoclinic, $P2_1/n$

Hall symbol: -P 2yn

$a = 8.9204 (2) \text{ \AA}$

$b = 6.0350 (2) \text{ \AA}$

$c = 22.4750 (6) \text{ \AA}$

$\beta = 90.051 (3)^\circ$

$V = 1209.93 (6) \text{ \AA}^3$

$Z = 4$

$F(000) = 520$

$D_x = 1.336 \text{ Mg m}^{-3}$

Mo $K\alpha$ radiation, $\lambda = 0.71073 \text{ \AA}$

Cell parameters from 3312 reflections

$\theta = 2.3\text{--}29.1^\circ$

$\mu = 0.26 \text{ mm}^{-1}$

$T = 100 \text{ K}$

Prism, colourless

$0.30 \times 0.20 \times 0.10 \text{ mm}$

Data collection

Agilent Supernova Dual

2723 independent reflections

diffractometer with an Atlas detector

Radiation source: SuperNova (Mo) X-ray Source 2251 reflections with $I > 2\sigma(I)$

Mirror $R_{\text{int}} = 0.036$

Detector resolution: 10.4041 pixels mm^{-1} $\theta_{\text{max}} = 27.5^\circ$, $\theta_{\text{min}} = 2.5^\circ$

ω scans $h = -11 \rightarrow 11$

Absorption correction: multi-scan
(*CrysAlis PRO*; Agilent, 2010) $k = -6 \rightarrow 7$

$T_{\text{min}} = 0.687$, $T_{\text{max}} = 1.000$ $l = -29 \rightarrow 29$

7394 measured reflections

Refinement

Refinement on F^2 Primary atom site location: structure-invariant direct methods

Least-squares matrix: full Secondary atom site location: difference Fourier map

$R[F^2 > 2\sigma(F^2)] = 0.041$ Hydrogen site location: inferred from neighbouring sites

$wR(F^2) = 0.143$ H atoms treated by a mixture of independent and constrained refinement

$S = 0.98$ $w = 1/[\sigma^2(F_o^2) + (0.1P)^2]$

where $P = (F_o^2 + 2F_c^2)/3$

2723 reflections $(\Delta/\sigma)_{\text{max}} = 0.001$

158 parameters $\Delta\rho_{\text{max}} = 0.36 \text{ e } \text{\AA}^{-3}$

3 restraints $\Delta\rho_{\text{min}} = -0.32 \text{ e } \text{\AA}^{-3}$

Special details

Geometry. All e.s.d.'s (except the e.s.d. in the dihedral angle between two l.s. planes) are estimated using the full covariance matrix. The cell e.s.d.'s are taken into account individually in the estimation of e.s.d.'s in distances, angles and torsion angles; correlations between e.s.d.'s in cell parameters are only used when they are defined by crystal symmetry. An approximate (isotropic) treatment of cell e.s.d.'s is used for estimating e.s.d.'s involving l.s. planes.

Refinement. Refinement of F^2 against ALL reflections. The weighted R -factor wR and goodness of fit S are based on F^2 , conventional R -factors R are based on F , with F set to zero for negative F^2 . The threshold expression of $F^2 > \sigma(F^2)$ is used only for calculating R -factors(gt) *etc.* and is not relevant to the choice of reflections for refinement. R -factors based on F^2 are statistically about twice as large as those based on F , and R -factors based on ALL data will be even larger.

Fractional atomic coordinates and isotropic or equivalent isotropic displacement parameters (\AA^2)

	<i>x</i>	<i>y</i>	<i>z</i>	$U_{\text{iso}}^*/U_{\text{eq}}$
S1	1.07001 (5)	1.46426 (7)	0.59008 (2)	0.01940 (18)
O1	0.65476 (14)	0.6300 (2)	0.54143 (6)	0.0209 (3)
O2	0.53180 (13)	0.7099 (2)	0.45694 (6)	0.0196 (3)
N1	0.80028 (16)	1.0010 (2)	0.53480 (7)	0.0156 (3)
N2	0.88614 (16)	1.1878 (2)	0.53543 (7)	0.0170 (3)
N3	0.94326 (16)	1.0938 (3)	0.63114 (7)	0.0169 (3)
C1	0.62814 (18)	0.7522 (3)	0.49413 (8)	0.0164 (4)
C2	0.72145 (19)	0.9564 (3)	0.48899 (8)	0.0162 (4)

supplementary materials

C3	0.7098 (2)	1.0904 (3)	0.43363 (8)	0.0211 (4)
H3A	0.6904	1.2455	0.4440	0.032*
H3B	0.6273	1.0339	0.4091	0.032*
H3C	0.8039	1.0800	0.4113	0.032*
C4	0.96146 (18)	1.2376 (3)	0.58689 (8)	0.0156 (4)
C5	1.00682 (18)	1.1176 (3)	0.69060 (7)	0.0157 (4)
H5	1.1055	1.1951	0.6871	0.019*
C6	1.0336 (2)	0.8900 (3)	0.71759 (9)	0.0210 (4)
H6A	1.1037	0.8054	0.6921	0.025*
H6B	0.9377	0.8076	0.7194	0.025*
C7	1.0991 (2)	0.9113 (3)	0.78024 (8)	0.0221 (4)
H7A	1.1116	0.7618	0.7977	0.026*
H7B	1.1994	0.9811	0.7779	0.026*
C8	0.9984 (2)	1.0501 (3)	0.82063 (9)	0.0235 (4)
H8A	0.9019	0.9725	0.8267	0.028*
H8B	1.0471	1.0689	0.8599	0.028*
C9	0.9696 (2)	1.2771 (3)	0.79289 (9)	0.0250 (4)
H9A	0.8988	1.3610	0.8182	0.030*
H9B	1.0648	1.3611	0.7912	0.030*
C10	0.9048 (2)	1.2561 (3)	0.73033 (8)	0.0212 (4)
H10A	0.8046	1.1858	0.7325	0.025*
H10B	0.8924	1.4055	0.7128	0.025*
H1	0.602 (2)	0.514 (3)	0.5431 (12)	0.040 (7)*
H2	0.892 (2)	1.275 (3)	0.5035 (6)	0.026 (6)*
H3	0.888 (2)	0.976 (2)	0.6260 (11)	0.029 (6)*

Atomic displacement parameters (\AA^2)

	U^{11}	U^{22}	U^{33}	U^{12}	U^{13}	U^{23}
S1	0.0251 (3)	0.0177 (3)	0.0154 (3)	-0.00825 (16)	-0.0030 (2)	0.00088 (18)
O1	0.0233 (6)	0.0208 (7)	0.0185 (7)	-0.0089 (5)	-0.0048 (5)	0.0055 (6)
O2	0.0200 (6)	0.0205 (7)	0.0182 (7)	-0.0064 (5)	-0.0030 (5)	0.0018 (5)
N1	0.0148 (7)	0.0144 (7)	0.0178 (8)	-0.0030 (5)	0.0009 (6)	-0.0010 (6)
N2	0.0214 (7)	0.0140 (7)	0.0157 (8)	-0.0039 (6)	-0.0007 (6)	0.0036 (6)
N3	0.0196 (7)	0.0172 (7)	0.0140 (8)	-0.0068 (6)	-0.0047 (6)	0.0006 (6)
C1	0.0159 (8)	0.0178 (9)	0.0155 (9)	0.0010 (7)	0.0000 (7)	-0.0017 (7)
C2	0.0153 (8)	0.0168 (9)	0.0166 (9)	-0.0015 (6)	0.0005 (7)	0.0006 (7)
C3	0.0247 (9)	0.0215 (9)	0.0169 (9)	-0.0069 (7)	-0.0040 (7)	0.0022 (8)
C4	0.0154 (8)	0.0160 (8)	0.0155 (8)	-0.0002 (6)	0.0004 (7)	-0.0008 (7)
C5	0.0173 (8)	0.0184 (9)	0.0112 (8)	-0.0030 (7)	-0.0038 (6)	-0.0006 (7)
C6	0.0255 (9)	0.0168 (9)	0.0208 (9)	0.0006 (7)	0.0006 (8)	-0.0003 (8)
C7	0.0266 (9)	0.0186 (9)	0.0210 (10)	0.0007 (7)	-0.0032 (8)	0.0036 (8)
C8	0.0301 (10)	0.0263 (10)	0.0141 (9)	-0.0050 (8)	-0.0001 (8)	-0.0016 (8)
C9	0.0318 (10)	0.0241 (10)	0.0191 (9)	0.0032 (8)	-0.0004 (8)	-0.0057 (8)
C10	0.0255 (9)	0.0197 (9)	0.0185 (9)	0.0037 (7)	-0.0007 (8)	-0.0011 (8)

Geometric parameters (\AA , $^\circ$)

S1—C4	1.6774 (18)	C5—C10	1.525 (2)
-------	-------------	--------	-----------

O1—C1	1.315 (2)	C5—H5	1.0000
O1—H1	0.845 (10)	C6—C7	1.530 (3)
O2—C1	1.225 (2)	C6—H6A	0.9900
N1—C2	1.275 (2)	C6—H6B	0.9900
N1—N2	1.363 (2)	C7—C8	1.528 (3)
N2—C4	1.370 (2)	C7—H7A	0.9900
N2—H2	0.892 (9)	C7—H7B	0.9900
N3—C4	1.330 (2)	C8—C9	1.527 (3)
N3—C5	1.458 (2)	C8—H8A	0.9900
N3—H3	0.870 (9)	C8—H8B	0.9900
C1—C2	1.491 (2)	C9—C10	1.525 (3)
C2—C3	1.488 (3)	C9—H9A	0.9900
C3—H3A	0.9800	C9—H9B	0.9900
C3—H3B	0.9800	C10—H10A	0.9900
C3—H3C	0.9800	C10—H10B	0.9900
C5—C6	1.520 (3)		
C1—O1—H1	113.7 (18)	C5—C6—H6A	109.5
C2—N1—N2	119.52 (15)	C7—C6—H6A	109.5
N1—N2—C4	117.71 (14)	C5—C6—H6B	109.5
N1—N2—H2	121.1 (14)	C7—C6—H6B	109.5
C4—N2—H2	121.2 (14)	H6A—C6—H6B	108.1
C4—N3—C5	124.97 (14)	C8—C7—C6	111.63 (15)
C4—N3—H3	120.2 (16)	C8—C7—H7A	109.3
C5—N3—H3	114.8 (16)	C6—C7—H7A	109.3
O2—C1—O1	124.08 (15)	C8—C7—H7B	109.3
O2—C1—C2	120.71 (15)	C6—C7—H7B	109.3
O1—C1—C2	115.18 (15)	H7A—C7—H7B	108.0
N1—C2—C3	126.77 (15)	C9—C8—C7	110.38 (16)
N1—C2—C1	114.80 (15)	C9—C8—H8A	109.6
C3—C2—C1	118.38 (15)	C7—C8—H8A	109.6
C2—C3—H3A	109.5	C9—C8—H8B	109.6
C2—C3—H3B	109.5	C7—C8—H8B	109.6
H3A—C3—H3B	109.5	H8A—C8—H8B	108.1
C2—C3—H3C	109.5	C10—C9—C8	111.43 (15)
H3A—C3—H3C	109.5	C10—C9—H9A	109.3
H3B—C3—H3C	109.5	C8—C9—H9A	109.3
N3—C4—N2	115.34 (15)	C10—C9—H9B	109.3
N3—C4—S1	124.83 (14)	C8—C9—H9B	109.3
N2—C4—S1	119.82 (13)	H9A—C9—H9B	108.0
N3—C5—C6	109.71 (14)	C5—C10—C9	111.06 (14)
N3—C5—C10	111.03 (14)	C5—C10—H10A	109.4
C6—C5—C10	110.79 (15)	C9—C10—H10A	109.4
N3—C5—H5	108.4	C5—C10—H10B	109.4
C6—C5—H5	108.4	C9—C10—H10B	109.4
C10—C5—H5	108.4	H10A—C10—H10B	108.0
C5—C6—C7	110.56 (15)		
C2—N1—N2—C4	175.80 (15)	C4—N3—C5—C6	-151.10 (16)
N2—N1—C2—C3	-0.7 (3)	C4—N3—C5—C10	86.1 (2)

supplementary materials

N2—N1—C2—C1	-178.00 (14)	N3—C5—C6—C7	-179.54 (14)
O2—C1—C2—N1	169.21 (15)	C10—C5—C6—C7	-56.59 (19)
O1—C1—C2—N1	-8.9 (2)	C5—C6—C7—C8	56.5 (2)
O2—C1—C2—C3	-8.4 (2)	C6—C7—C8—C9	-55.6 (2)
O1—C1—C2—C3	173.52 (15)	C7—C8—C9—C10	55.2 (2)
C5—N3—C4—N2	-177.12 (15)	N3—C5—C10—C9	178.84 (14)
C5—N3—C4—S1	4.2 (2)	C6—C5—C10—C9	56.7 (2)
N1—N2—C4—N3	0.7 (2)	C8—C9—C10—C5	-56.1 (2)
N1—N2—C4—S1	179.49 (12)		

Hydrogen-bond geometry (\AA , $^\circ$)

$D-H\cdots A$	$D-H$	$H\cdots A$	$D\cdots A$	$D-H\cdots A$
N3—H3 \cdots N1	0.87 (1)	2.20 (2)	2.574 (2)	106.(2)
O1—H1 \cdots O2 ⁱ	0.85 (1)	1.80 (1)	2.6416 (17)	172 (3)
N2—H2 \cdots S1 ⁱⁱ	0.89 (1)	2.65 (1)	3.5384 (15)	176.(2)

Symmetry codes: (i) $-x+1, -y+1, -z+1$; (ii) $-x+2, -y+3, -z+1$.

Fig. 1




Fig. 2




Fig. 3

