

1,4-Bis(5-methyl-1*H*-1,2,4-triazol-3-yl)-benzene tetrahydrate

Ai-Xin Zhu,^{a*} Xiu-Li Chen,^b Zhen Li,^a Yuan-Chao Du^a and Hong-Can Wang^a

^aFaculty of Chemistry and Chemical Engineering, Yunnan Normal University, Kunming 650092, People's Republic of China, and ^bDepartment of Chemistry, Zhengzhou Normal University, Zhengzhou 450044, People's Republic of China
Correspondence e-mail: zaxchem@126.com


Received 9 April 2011; accepted 21 April 2011

Key indicators: single-crystal X-ray study; $T = 293$ K; mean $\sigma(\text{C}-\text{C}) = 0.003$ Å; R factor = 0.052; wR factor = 0.148; data-to-parameter ratio = 15.1.

In the title compound, $\text{C}_{12}\text{H}_{12}\text{N}_6 \cdot 4\text{H}_2\text{O}$, the two triazole rings adopt a *cis* configuration with a crystallographic twofold axis passing through the central benzene group. The benzene and triazole rings are almost coplanar with a dihedral angle of 5.5 (1)°. In the crystal, water molecules are joined together by $\text{OW}-\text{H}\cdots\text{OW}$ hydrogen bonds to form a one-dimensional zigzag chain. These water chains are further connected to the organic molecule, forming a three-dimensional network by intermolecular $\text{OW}-\text{H}\cdots\text{N}$ and $\text{N}-\text{H}\cdots\text{OW}$ hydrogen bonds. Moreover, $\pi-\pi$ stacking interactions between triazole rings [centroid-centroid distances = 3.667 (1)– 3.731 (1) Å] are observed. One of the water molecules shows one of the H atoms to be disordered over two positions.

Related literature

For applications of 1,2,4-triazole and its derivatives in coordination chemistry, see: Zhang *et al.* (2005); Ouellette *et al.* (2006); Zhu *et al.* (2009). For the structures of ruthenium complexes with pyridine-2-yl-1,2,4-triazole-based ligands, see: Passaniti *et al.* (2002). For the previous synthesis of the title compound, see: Bahçeci *et al.* (2005).



Experimental

Crystal data

$\text{C}_{12}\text{H}_{12}\text{N}_6 \cdot 4\text{H}_2\text{O}$
 $M_r = 312.34$
Monoclinic, $C2/c$
 $a = 12.7343$ (19) Å

$b = 13.937$ (2) Å
 $c = 9.0648$ (14) Å
 $\beta = 100.893$ (3)°
 $V = 1579.8$ (4) Å³

$Z = 4$
Mo $K\alpha$ radiation
 $\mu = 0.10$ mm⁻¹

$T = 293$ K
 $0.35 \times 0.28 \times 0.08$ mm

Data collection

Bruker APEX CCD diffractometer
Absorption correction: multi-scan
(*SADABS*; Sheldrick, 1996)
 $T_{\min} = 0.966$, $T_{\max} = 0.992$

4670 measured reflections
1542 independent reflections
1286 reflections with $I > 2\sigma(I)$
 $R_{\text{int}} = 0.021$

Refinement

$R[F^2 > 2\sigma(F^2)] = 0.052$
 $wR(F^2) = 0.148$
 $S = 1.04$
1542 reflections

102 parameters
H-atom parameters constrained
 $\Delta\rho_{\text{max}} = 0.27$ e Å⁻³
 $\Delta\rho_{\text{min}} = -0.25$ e Å⁻³

Table 1

Hydrogen-bond geometry (Å, °).

$D-\text{H}\cdots A$	$D-\text{H}$	$\text{H}\cdots A$	$D\cdots A$	$D-\text{H}\cdots A$
$\text{N1}-\text{H1D}\cdots\text{O1W}$	0.86	1.88	2.736 (2)	173
$\text{O1W}-\text{H1WA}\cdots\text{N2}^i$	0.85	2.08	2.926 (2)	172
$\text{O1W}-\text{H1WB}\cdots\text{O2W}^{ii}$	0.85	1.96	2.801 (2)	170
$\text{O2W}-\text{H2WA}\cdots\text{N3}^{iii}$	0.85	1.95	2.800 (2)	173
$\text{O2W}-\text{H2WB}\cdots\text{O2W}^{iii}$	0.85	1.93	2.754 (3)	164
$\text{O2W}-\text{H2WC}\cdots\text{O2W}^{iv}$	0.85	1.92	2.774 (3)	178

Symmetry codes: (i) $x, -y + 1, z - \frac{1}{2}$; (ii) $x + \frac{1}{2}, -y + \frac{1}{2}, z - \frac{1}{2}$; (iii) $-x + 1, -y, -z$; (iv) $-x + 1, y, -z + \frac{1}{2}$.

Data collection: *SMART* (Bruker, 2004); cell refinement: *SAINTE* (Bruker, 2004); data reduction: *SAINTE*; program(s) used to solve structure: *SHELXS97* (Sheldrick, 2008); program(s) used to refine structure: *SHELXL97* (Sheldrick, 2008); molecular graphics: *DIAMOND* (Brandenburg, 1999); software used to prepare material for publication: *SHELXTL* (Sheldrick, 2008).

The authors thank the Youth Foundation (grant No. 10QZ02) of Yunnan Normal University, the Science Foundation of the Education Department (grant No. 2010Y004) as well as the Science and Technology Department (grant No. 2010ZC070) of Yunnan Province for supporting this work.

Supplementary data and figures for this paper are available from the IUCr electronic archives (Reference: IM2280).

References

- Bahçeci, S., Yükek, H. & Serdar, M. (2005). *Indian J. Chem. Sect. B*, **44**, 568–572.
- Brandenburg, K. (1999). *DIAMOND*. Crystal Impact GbR, Bonn, Germany.
- Bruker (2004). *SMART* and *SAINTE*. Bruker AXS Inc., Madison, Wisconsin, USA.
- Ouellette, W., Prosvirin, A. V., Chieffo, V., Dunbar, K. R., Hudson, B. & Zubieta, J. (2006). *Inorg. Chem.* **45**, 9346–9366.
- Passaniti, P., Browne, W. R., Lynch, F. C., Hughes, D., Nieuwenhuysen, M., James, P., Maestri, M. & Vos, J. G. (2002). *J. Chem. Soc. Dalton Trans.* pp. 1740–1746.
- Sheldrick, G. M. (1996). *SADABS*. University of Göttingen, Germany.
- Sheldrick, G. M. (2008). *Acta Cryst.* **A64**, 112–122.
- Zhang, J.-P., Lin, Y.-Y., Huang, X.-C. & Chen, X.-M. (2005). *J. Am. Chem. Soc.* **127**, 5495–5506.
- Zhu, A.-X., Lin, J.-B., Zhang, J.-P. & Chen, X.-M. (2009). *Inorg. Chem.* **48**, 3882–3889.

supplementary materials

Acta Cryst. (2011). E67, o1255 [doi:10.1107/S1600536811015133]

1,4-Bis(5-methyl-1*H*-1,2,4-triazol-3-yl)benzene tetrahydrate

A.-X. Zhu, X.-L. Chen, Z. Li, Y.-C. Du and H.-C. Wang

Comment

In the past few years, 1,2,4-triazole and its derivatives have attracted increasing attention as *N*-heterocyclic aromatic ligands, since they are effective bridging ligands combining the coordination modes of both imidazoles and pyrazoles. In addition, metal-triazolate frameworks have demonstrated high thermal and chemical stabilities, and interesting luminescent, magnetic and gas-adsorption properties (Zhang *et al.* 2005; Ouellette *et al.* 2006; Zhu *et al.* 2009). However, there are rare crystal structure reports of 1,2,4-triazole derivatives attached to aromatic groups (Passaniti *et al.* 2002). Although the synthesis of the title compound 1,4-bis(5-methyl-1*H*-1,2,4-triazol-3-yl)benzene has been reported by Bahçeci *et al.* (2005), no crystallographic study has been reported on this ligand and related metal coordination compounds. We reported herein another synthetic method and the crystal structure of the title compound.

The asymmetric unit of the title compound contains one-half organic molecule, which adopts a *cis*-configuration with a crystallographic mirror plane passing through the central benzene group, and two water molecules (Fig. 1). The bond lengths and angles are within normal ranges in accordance with the corresponding values reported (Passaniti *et al.* 2002). The benzene and the triazole rings are almost coplanar, with a dihedral angle of 5.4 (1)°. In the crystal structure, water molecules are joined together by OW—H···OW hydrogen bonds to form a one-dimensional zig-zag water chain (Fig. 2, Table 1). These water chains are further connected to the organic molecule producing a three-dimensional network (Fig. 2) by intermolecular OW—H···N and N—H···OW hydrogen bonds (Table 1). Moreover, π - π stacking interactions between triazole rings (centroid-centroid distance = 3.665 (1)–3.732 (1) Å) are observed (Fig. 3).

Experimental

The ligand 1,4-bis(5-methyl-1*H*-1,2,4-triazol-3-yl)benzene was synthesized according to a literature method (Bahçeci *et al.* 2005). Yellow, plate-like single crystals of the title compound are obtained from a solution of 1,4-bis(5-methyl-1*H*-1,2,4-triazol-3-yl)benzene (24 mg, 0.1 mmol) in methanol (1 ml) and water (5 ml) if the solution is placed in a Teflon-lined stainless steel vessel (15 ml), heated at 453 K for 24 h and then cooled to room temperature at a rate of 5 K h⁻¹.

Refinement

All H atoms were placed in idealized positions (O—H = 0.85 Å, N—H = 0.86 Å and C—H = 0.95 Å) and refined as riding atoms with $U_{\text{iso}}(\text{H}) = 1.2U_{\text{eq}}(\text{C}, \text{N})$ and $U_{\text{iso}}(\text{H}) = 1.5U_{\text{eq}}(\text{O})$. One hydrogen atom from O2W is disordered over two positions in a 0.52 (3):0.48 (3) ratio, which is freely refined with the command 'PART'.

Figures




Fig. 1. Molecular structure of the title compound, non-H atoms are depicted as 30% probability displacement ellipsoids.




Fig. 2. Packing diagram of the title compound showing the hydrogen bonding interactions as dashed lines. H atoms not involved in hydrogen bondings have been omitted.




Fig. 3. π - π Stacking interactions between triazole rings, H and O atoms are omitted for clarity.

5-methyl-3-[4-(5-methyl-1H-1,2,4-triazol-3-yl)phenyl]-1H-1,2,4-triazole tetrahydrate

Crystal data

$C_{12}H_{12}N_6 \cdot 4H_2O$	$Z = 4$
$M_r = 312.34$	$F(000) = 664$
Monoclinic, $C2/c$	$D_x = 1.313 \text{ Mg m}^{-3}$
Hall symbol: $-C 2yc$	Mo $K\alpha$ radiation, $\lambda = 0.71073 \text{ \AA}$
$a = 12.7343 (19) \text{ \AA}$	$\mu = 0.10 \text{ mm}^{-1}$
$b = 13.937 (2) \text{ \AA}$	$T = 293 \text{ K}$
$c = 9.0648 (14) \text{ \AA}$	Plate, yellow
$\beta = 100.893 (3)^\circ$	$0.35 \times 0.28 \times 0.08 \text{ mm}$
$V = 1579.8 (4) \text{ \AA}^3$	

Data collection

Bruker APEX CCD diffractometer	1542 independent reflections
Radiation source: fine-focus sealed tube graphite	1286 reflections with $I > 2\sigma(I)$
ω scans	$R_{\text{int}} = 0.021$
Absorption correction: multi-scan (SADABS; Sheldrick, 1996)	$\theta_{\text{max}} = 26.0^\circ$, $\theta_{\text{min}} = 2.2^\circ$
$T_{\text{min}} = 0.966$, $T_{\text{max}} = 0.992$	$h = -15 \rightarrow 15$
4670 measured reflections	$k = -15 \rightarrow 17$
	$l = -11 \rightarrow 11$

Refinement

Refinement on F^2	Primary atom site location: structure-invariant direct methods
---------------------	--

Least-squares matrix: full

$$R[F^2 > 2\sigma(F^2)] = 0.052$$

$$wR(F^2) = 0.148$$

$$S = 1.04$$

1542 reflections

102 parameters

0 restraints

Secondary atom site location: difference Fourier map

Hydrogen site location: inferred from neighbouring sites

H-atom parameters constrained

$$w = 1/[\sigma^2(F_o^2) + (0.0816P)^2 + 0.7602P]$$

$$\text{where } P = (F_o^2 + 2F_c^2)/3$$

$$(\Delta/\sigma)_{\max} < 0.001$$

$$\Delta\rho_{\max} = 0.27 \text{ e } \text{\AA}^{-3}$$

$$\Delta\rho_{\min} = -0.25 \text{ e } \text{\AA}^{-3}$$

Special details

Geometry. All e.s.d.'s (except the e.s.d. in the dihedral angle between two l.s. planes) are estimated using the full covariance matrix. The cell e.s.d.'s are taken into account individually in the estimation of e.s.d.'s in distances, angles and torsion angles; correlations between e.s.d.'s in cell parameters are only used when they are defined by crystal symmetry. An approximate (isotropic) treatment of cell e.s.d.'s is used for estimating e.s.d.'s involving l.s. planes.

Refinement. Refinement of F^2 against ALL reflections. The weighted R -factor wR and goodness of fit S are based on F^2 , conventional R -factors R are based on F , with F set to zero for negative F^2 . The threshold expression of $F^2 > \sigma(F^2)$ is used only for calculating R -factors(gt) *etc.* and is not relevant to the choice of reflections for refinement. R -factors based on F^2 are statistically about twice as large as those based on F , and R -factors based on ALL data will be even larger. One hydrogen atom from O2W is disordered over two positions in a 0.52 (3):0.48 (3) ratio, which is freely refined with command 'PART'.

Fractional atomic coordinates and isotropic or equivalent isotropic displacement parameters (\AA^2)

	<i>x</i>	<i>y</i>	<i>z</i>	$U_{\text{iso}}^*/U_{\text{eq}}$	Occ. (<1)
N1	0.66106 (13)	0.34969 (11)	-0.19565 (17)	0.0531 (5)	
H1D	0.6874	0.3847	-0.2578	0.064*	
N2	0.62354 (14)	0.38395 (11)	-0.07499 (17)	0.0522 (5)	
N3	0.60877 (12)	0.22386 (10)	-0.09140 (16)	0.0454 (4)	
C1	0.68428 (18)	0.19667 (15)	-0.3248 (2)	0.0590 (6)	
H1A	0.7292	0.1449	-0.2805	0.088*	
H1B	0.7230	0.2361	-0.3830	0.088*	
H1C	0.6218	0.1711	-0.3887	0.088*	
C2	0.65166 (14)	0.25529 (13)	-0.20444 (18)	0.0448 (4)	
C3	0.59264 (14)	0.30486 (12)	-0.01538 (19)	0.0425 (4)	
C4	0.54472 (14)	0.30544 (12)	0.12072 (19)	0.0421 (4)	
C5	0.52248 (17)	0.22014 (13)	0.1863 (2)	0.0510 (5)	
H5A	0.5379	0.1622	0.1443	0.061*	
C6	0.52170 (16)	0.39108 (13)	0.1860 (2)	0.0510 (5)	
H6A	0.5358	0.4491	0.1429	0.061*	
O1W	0.73472 (14)	0.45314 (11)	-0.41166 (19)	0.0757 (5)	
H1WA	0.6969	0.4969	-0.4609	0.091*	
H1WB	0.8001	0.4691	-0.4017	0.091*	
O2W	0.44076 (12)	-0.02791 (9)	0.10493 (16)	0.0602 (4)	
H2WA	0.4240	-0.0868	0.0928	0.072*	
H2WB	0.4790	-0.0006	0.0500	0.072*	0.52 (3)
H2WC	0.4784	-0.0277	0.1930	0.072*	0.48 (3)

supplementary materials

Atomic displacement parameters (\AA^2)

	U^{11}	U^{22}	U^{33}	U^{12}	U^{13}	U^{23}
N1	0.0657 (11)	0.0516 (10)	0.0484 (8)	-0.0032 (8)	0.0274 (8)	0.0074 (7)
N2	0.0662 (11)	0.0437 (9)	0.0523 (9)	-0.0025 (7)	0.0257 (8)	0.0035 (7)
N3	0.0540 (9)	0.0428 (8)	0.0425 (8)	-0.0035 (7)	0.0166 (7)	-0.0015 (6)
C1	0.0649 (13)	0.0672 (13)	0.0495 (10)	-0.0045 (10)	0.0232 (9)	-0.0060 (9)
C2	0.0467 (10)	0.0488 (10)	0.0401 (9)	-0.0022 (8)	0.0115 (7)	0.0016 (7)
C3	0.0452 (10)	0.0437 (9)	0.0397 (9)	-0.0003 (7)	0.0108 (7)	0.0012 (7)
C4	0.0448 (10)	0.0431 (10)	0.0396 (9)	-0.0001 (7)	0.0109 (7)	0.0010 (7)
C5	0.0774 (14)	0.0368 (9)	0.0429 (9)	0.0004 (9)	0.0218 (9)	-0.0027 (7)
C6	0.0603 (12)	0.0380 (10)	0.0604 (11)	-0.0004 (8)	0.0260 (9)	0.0049 (8)
O1W	0.0804 (11)	0.0679 (10)	0.0871 (11)	0.0058 (8)	0.0373 (9)	0.0277 (8)
O2W	0.0809 (11)	0.0447 (7)	0.0611 (9)	-0.0064 (7)	0.0289 (8)	-0.0014 (6)

Geometric parameters (\AA , $^\circ$)

N1—C2	1.322 (3)	C4—C5	1.382 (2)
N1—N2	1.360 (2)	C4—C6	1.388 (2)
N1—H1D	0.8600	C5—C5 ⁱ	1.383 (4)
N2—C3	1.320 (2)	C5—H5A	0.9300
N3—C2	1.324 (2)	C6—C6 ⁱ	1.376 (4)
N3—C3	1.358 (2)	C6—H6A	0.9300
C1—C2	1.484 (3)	O1W—H1WA	0.8500
C1—H1A	0.9600	O1W—H1WB	0.8500
C1—H1B	0.9600	O2W—H2WA	0.8500
C1—H1C	0.9600	O2W—H2WB	0.8501
C3—C4	1.476 (2)	O2W—H2WC	0.8499
C2—N1—N2	110.87 (14)	N2—C3—C4	122.67 (15)
C2—N1—H1D	124.6	N3—C3—C4	123.69 (15)
N2—N1—H1D	124.6	C5—C4—C6	118.63 (17)
C3—N2—N1	102.31 (15)	C5—C4—C3	120.35 (15)
C2—N3—C3	104.00 (15)	C6—C4—C3	121.01 (15)
C2—C1—H1A	109.5	C4—C5—C5 ⁱ	120.67 (10)
C2—C1—H1B	109.5	C4—C5—H5A	119.7
H1A—C1—H1B	109.5	C5 ⁱ —C5—H5A	119.7
C2—C1—H1C	109.5	C6 ⁱ —C6—C4	120.69 (10)
H1A—C1—H1C	109.5	C6 ⁱ —C6—H6A	119.7
H1B—C1—H1C	109.5	C4—C6—H6A	119.7
N1—C2—N3	109.18 (15)	H1WA—O1W—H1WB	108.3
N1—C2—C1	123.88 (16)	H2WA—O2W—H2WB	121.0
N3—C2—C1	126.93 (17)	H2WA—O2W—H2WC	102.0
N2—C3—N3	113.64 (16)	H2WB—O2W—H2WC	105.4

Symmetry codes: (i) $-x+1, y, -z+1/2$.

Hydrogen-bond geometry (Å, °)

<i>D</i> —H··· <i>A</i>	<i>D</i> —H	H··· <i>A</i>	<i>D</i> ··· <i>A</i>	<i>D</i> —H··· <i>A</i>
N1—H1D···O1W	0.86	1.88	2.736 (2)	173.
O1W—H1WA···N2 ⁱⁱ	0.85	2.08	2.926 (2)	172.
O1W—H1WB···O2W ⁱⁱⁱ	0.85	1.96	2.801 (2)	170.
O2W—H2WA···N3 ^{iv}	0.85	1.95	2.800 (2)	173.
O2W—H2WB···O2W ^{iv}	0.85	1.93	2.754 (3)	164.
O2W—H2WC···O2W ⁱ	0.85	1.92	2.774 (3)	178.

Symmetry codes: (ii) $x, -y+1, z-1/2$; (iii) $x+1/2, -y+1/2, z-1/2$; (iv) $-x+1, -y, -z$; (i) $-x+1, y, -z+1/2$.

Fig. 1




Fig. 2




Fig. 3

