

Poly[[triaqua(μ_3 -4-oxidopyridine-2,6-dicarboxylato)holmium(III)] mono-hydrate]

Zhu-Qing Gao,^{a*} Dong-Yu Lv,^b Jin-Zhong Gu^b and Hong-Jin Li^a

^aSchool of Chemistry and Biological Engineering, Taiyuan University of Science and Technology, Taiyuan 030021, People's Republic of China, and ^bKey Laboratory of Nonferrous Metal Chemistry and Resources Utilization of Gansu Province, College of Chemistry and Chemical Engineering, Lanzhou University, Lanzhou 730000, People's Republic of China

Correspondence e-mail: zqgao2008@163.com


Received 28 April 2011; accepted 5 May 2011

Key indicators: single-crystal X-ray study; $T = 296$ K; mean $\sigma(\text{C}-\text{C}) = 0.006$ Å; R factor = 0.021; wR factor = 0.047; data-to-parameter ratio = 10.2.

In the title coordination polymer, $\{[\text{Ho}(\text{C}_7\text{H}_2\text{NO}_5)(\text{H}_2\text{O})_3]\cdot\text{H}_2\text{O}\}_n$, the Ho^{III} atom is eight-coordinated by a tridentate 4-oxidopyridine-2,6-dicarboxylate trianion, two monodentate anions and three water molecules, forming a distorted bicapped trigonal-prismatic HoNO_7 coordination geometry. The anions bridge adjacent Ho^{III} ions into double chains. Adjacent chains are further connected into sheets. O—H···O hydrogen bonds involving both coordinated and uncoordinated water molecules generate a three-dimensional supramolecular framework.

Related literature

For the structures and properties of lanthanide coordination compounds, see: Wang *et al.* (2007); Lv *et al.* (2010); Gao *et al.* (2006). For bond lengths and angles in other complexes with eight-coordinate Ho^{III} , see: Wang *et al.* (2007); Munoz *et al.* (2005).



Experimental

Crystal data

$[\text{Ho}(\text{C}_7\text{H}_2\text{NO}_5)(\text{H}_2\text{O})_3]\cdot\text{H}_2\text{O}$	$V = 1107.5$ (9) Å ³
$M_r = 417.09$	$Z = 4$
Monoclinic, $P2_1/n$	Mo $K\alpha$ radiation
$a = 9.879$ (5) Å	$\mu = 7.19$ mm ⁻¹
$b = 7.557$ (4) Å	$T = 296$ K
$c = 15.386$ (8) Å	$0.32 \times 0.29 \times 0.26$ mm
$\beta = 105.386$ (5)°	

Data collection

Bruker APEXII CCD diffractometer	4776 measured reflections
Absorption correction: multi-scan (<i>SADABS</i> ; Bruker, 2004)	2003 independent reflections
$T_{\min} = 0.207$, $T_{\max} = 0.257$	1646 reflections with $I > 2\sigma(I)$
	$R_{\text{int}} = 0.025$

Refinement

$R[F^2 > 2\sigma(F^2)] = 0.021$	H atoms treated by a mixture of independent and constrained refinement
$wR(F^2) = 0.047$	$\Delta\rho_{\text{max}} = 0.71$ e Å ⁻³
$S = 1.02$	$\Delta\rho_{\text{min}} = -0.54$ e Å ⁻³
2003 reflections	
196 parameters	
12 restraints	

Table 1
Hydrogen-bond geometry (Å, °).

$D-\text{H}\cdots A$	$D-\text{H}$	$\text{H}\cdots A$	$D\cdots A$	$D-\text{H}\cdots A$
O9—H8W···O2 ⁱ	0.86 (1)	2.33 (4)	3.080 (5)	145 (6)
O9—H7W···O4	0.86 (1)	1.84 (1)	2.693 (5)	170 (5)
O8—H6W···O9 ⁱⁱ	0.86 (1)	2.28 (3)	3.026 (5)	145 (5)
O8—H5W···O9 ⁱⁱⁱ	0.86 (1)	1.84 (2)	2.689 (5)	168 (5)
O7—H4W···O3 ^{iv}	0.86 (1)	2.08 (3)	2.788 (4)	139 (4)
O7—H3W···O5 ^{iv}	0.87 (1)	1.92 (3)	2.711 (4)	152 (4)
O6—H2W···O1 ^v	0.86 (1)	1.85 (2)	2.671 (4)	159 (5)
O6—H1W···O4 ^{vi}	0.86 (1)	1.83 (1)	2.687 (4)	171 (5)

Symmetry codes: (i) $x - 1, y, z$; (ii) $x + \frac{1}{2}, -y + \frac{3}{2}, z - \frac{1}{2}$; (iii) $-x + \frac{1}{2}, y - \frac{1}{2}, -z + \frac{3}{2}$; (iv) $-x + 1, -y + 1, -z + 2$; (v) $-x + \frac{3}{2}, y + \frac{1}{2}, -z + \frac{3}{2}$; (vi) $-x + 1, -y + 2, -z + 2$.

Data collection: *APEX2* (Bruker, 2004); cell refinement: *SAINT* (Bruker, 2004); data reduction: *SAINT*; program(s) used to solve structure: *SHELXS97* (Sheldrick, 2008); program(s) used to refine structure: *SHELXL97* (Sheldrick, 2008); molecular graphics: *SHELXTL* (Sheldrick, 2008); software used to prepare material for publication: *SHELXTL*.

Supplementary data and figures for this paper are available from the IUCr electronic archives (Reference: SJ5136).

References

- Bruker (2004). *APEX2, SAINT* and *SADABS*. Bruker AXS Inc., Madison, Wisconsin, USA.
- Gao, H. L., Yi, L., Zhao, B., Zhao, X. Q., Cheng, P., Liao, D. Z. & Yan, S. P. (2006). *Inorg. Chem.* **45**, 5980–5988.
- Lv, D.-Y., Gao, Z.-Q. & Gu, J.-Z. (2010). *Acta Cryst. E66*, m1694–m1695.
- Munoz, J. C., Atria, A. M., Baggio, R., Garland, M. T., Pena, O. & Orrego, C. (2005). *Inorg. Chim. Acta*, **358**, 4027–4033.
- Sheldrick, G. M. (2008). *Acta Cryst. A64*, 112–122.
- Wang, H. S., Zhao, B., Zhai, B., Shi, W., Cheng, P., Liao, D. Z. & Yan, S. P. (2007). *Cryst. Growth Des.* **7**, 1851–1857.

supplementary materials

Acta Cryst. (2011). E67, m740 [doi:10.1107/S1600536811016953]

Poly[[triaqua(μ_3 -4-oxidopyridine-2,6-dicarboxylato)holmium(III)] monohydrate]

Z.-Q. Gao, D.-Y. Lv, J.-Z. Gu and H.-J. Li

Comment

Lanthanide coordination polymers have shown not only versatile architectures but also desirable properties, *e.g.*, luminescent, magnetic, catalytic, and gas absorption and separation properties (Wang *et al.*, 2007; Lv *et al.*, 2010). In order to extend our investigations in this field, we designed and synthesized one lanthanide coordination polymer $\{[\text{Ho}(\text{C}_7\text{H}_2\text{NO}_5)(\text{H}_2\text{O})_3]\cdot\text{H}_2\text{O}\}_n$ by choosing 4-oxidopyridine-2,6-dicarboxylic acid as a functional ligand, and report its structure here.

The title compound is isotopic with its Dy (Gao *et al.*, 2006) and Eu (Lv *et al.*, 2010) analogues. As shown in Fig. 1, the asymmetric unit of the cell contains one Ho(III) ion, one 4-oxidopyridine-2,6-dicarboxylate anion, three coordinated water molecules, and one water molecule of crystallization. The Ho atom is eight-coordinated by seven oxygen atoms from three anions and three coordinated water molecules and by one nitrogen atom from one tridentate anion (the other two anions are monodentate), forming a distorted bicapped trigonal-prismatic coordination environment.

Important bond distances are presented in Table 1. The Ho–O bond lengths [2.279 (3) to 2.400 (3) Å] are shorter than the Ho–N bond length [2.450 (3) Å], which is in agreement with the bond lengths observed in other Ho(III) complexes (Wang *et al.*, 2007; Munoz *et al.*, 2005). The anion adopts a μ_3 -pentadentate coordination mode, as shown in Fig. 1. The anions bridge the adjacent Ho^{III} ions to form infinite double chains (Fig. 2). Adjacent chains are further connected by the coordination of the anions and Ho(III) ions into a two-dimensional sheet (Fig. 3), which are further extended into a three-dimensional supramolecular framework through O–H···O hydrogen-bonding interactions including both coordinated and uncoordinated water molecules (Table 2).

Experimental

To a solution of holmium(III)nitrate hexahydrate (0.138 g, 0.3 mmol) in water (5 ml) was added an aqueous solution (5 ml) of the ligand (0.060 g, 0.3 mmol) and a drop of triethylamine. The reactants were sealed in a 25-ml Teflon-lined, stainless-steel Parr bomb. The bomb was heated at 433 K for 3 days. The cool solution yielded colorless single crystals in *ca* 75% yield. Anal. Calcd for C₇H₁₀TbNO₉: C, 20.16; H, 2.42; N, 3.36. Found: C, 20.45; H, 2.17; N, 3.64.

Refinement

The coordinated water H atoms were located in a different Fourier map and refined with distance constraints of O–H = 0.83 (3) Å. The free water H atoms were placed at calculated positions and refined with a riding model, considering the position of oxygen atoms and the quantity of H atoms. The carbon-bound H atoms were placed in geometrically idealized positions, with C–H = 0.93 Å and constrained to ride on their respective parent atoms, with $U_{\text{iso}}(\text{H}) = 1.2 U_{\text{eq}}(\text{C})$.

supplementary materials

Figures




Fig. 1. The asymmetric unit of the title complex, showing the atom-labeling scheme. Displacement ellipsoids are drawn at the 30% probability level.




Fig. 2. View along the b axis, showing the polymeric double chain.




Fig. 3. View along the a axis, showing the sheet structure of $\{[\text{Ho}(\text{C}_7\text{H}_2\text{NO}_5)(\text{H}_2\text{O})_3]\cdot\text{H}_2\text{O}\}_n$.

Poly[[triaqua(μ_3 -4-oxidopyridine-2,6-dicarboxylato)holmium(III)] monohydrate]

Crystal data

$[\text{Ho}(\text{C}_7\text{H}_2\text{NO}_5)(\text{H}_2\text{O})_3]\cdot\text{H}_2\text{O}$	$F(000) = 792$
$M_r = 417.09$	$D_x = 2.502 \text{ Mg m}^{-3}$
Monoclinic, $P2_1/n$	Mo $K\alpha$ radiation, $\lambda = 0.71073 \text{ \AA}$
Hall symbol: -P 2y/n	Cell parameters from 2292 reflections
$a = 9.879 (5) \text{ \AA}$	$\theta = 2.2\text{--}28.2^\circ$
$b = 7.557 (4) \text{ \AA}$	$\mu = 7.19 \text{ mm}^{-1}$
$c = 15.386 (8) \text{ \AA}$	$T = 296 \text{ K}$
$\beta = 105.386 (5)^\circ$	Block, colorless
$V = 1107.5 (9) \text{ \AA}^3$	$0.32 \times 0.29 \times 0.26 \text{ mm}$
$Z = 4$	

Data collection

Bruker APEXII CCD diffractometer	2003 independent reflections
Radiation source: fine-focus sealed tube graphite	1646 reflections with $I > 2\sigma(I)$
φ and ω scans	$R_{\text{int}} = 0.025$
Absorption correction: multi-scan (<i>SADABS</i> ; Bruker, 2004)	$\theta_{\text{max}} = 25.5^\circ, \theta_{\text{min}} = 2.2^\circ$
$T_{\text{min}} = 0.207, T_{\text{max}} = 0.257$	$h = -11 \rightarrow 7$
4776 measured reflections	$k = -8 \rightarrow 9$
	$l = -16 \rightarrow 17$

Refinement

Refinement on F^2	Secondary atom site location: difference Fourier map
Least-squares matrix: full	Hydrogen site location: inferred from neighbouring sites
$R[F^2 > 2\sigma(F^2)] = 0.021$	H atoms treated by a mixture of independent and constrained refinement
$wR(F^2) = 0.047$	$w = 1/[\sigma^2(F_o^2) + (0.0176P)^2]$ where $P = (F_o^2 + 2F_c^2)/3$
$S = 1.02$	$(\Delta/\sigma)_{\max} = 0.002$
2003 reflections	$\Delta\rho_{\max} = 0.71 \text{ e \AA}^{-3}$
196 parameters	$\Delta\rho_{\min} = -0.54 \text{ e \AA}^{-3}$
12 restraints	Extinction correction: <i>SHELXL97</i> (Sheldrick, 2008), $F_c^* = kF_c[1 + 0.001xF_c^2\lambda^3/\sin(2\theta)]^{1/4}$
Primary atom site location: structure-invariant direct methods	Extinction coefficient: 0.0119 (3)

Special details

Geometry. All e.s.d.'s (except the e.s.d. in the dihedral angle between two l.s. planes) are estimated using the full covariance matrix. The cell e.s.d.'s are taken into account individually in the estimation of e.s.d.'s in distances, angles and torsion angles; correlations between e.s.d.'s in cell parameters are only used when they are defined by crystal symmetry. An approximate (isotropic) treatment of cell e.s.d.'s is used for estimating e.s.d.'s involving l.s. planes.

Refinement. Refinement of F^2 against ALL reflections. The weighted R -factor wR and goodness of fit S are based on F^2 , conventional R -factors R are based on F , with F set to zero for negative F^2 . The threshold expression of $F^2 > \sigma(F^2)$ is used only for calculating R -factors(gt) etc. and is not relevant to the choice of reflections for refinement. R -factors based on F^2 are statistically about twice as large as those based on F , and R -factors based on ALL data will be even larger.

Fractional atomic coordinates and isotropic or equivalent isotropic displacement parameters (\AA^2)

	<i>x</i>	<i>y</i>	<i>z</i>	$U_{\text{iso}}^*/U_{\text{eq}}$
Ho1	0.501337 (17)	0.82185 (2)	0.747304 (11)	0.01332 (10)
C1	0.7829 (4)	0.5891 (5)	0.8379 (3)	0.0184 (9)
C2	0.7250 (4)	0.6133 (5)	0.9175 (3)	0.0158 (9)
C3	0.7940 (4)	0.5582 (5)	1.0018 (3)	0.0172 (9)
H3	0.8780	0.4963	1.0112	0.021*
C4	0.7379 (4)	0.5951 (5)	1.0743 (2)	0.0163 (9)
C5	0.6087 (4)	0.6828 (5)	1.0537 (3)	0.0193 (9)
H5	0.5652	0.7073	1.0991	0.023*
C6	0.5460 (4)	0.7328 (5)	0.9665 (3)	0.0157 (9)
C7	0.4083 (4)	0.8304 (5)	0.9374 (3)	0.0162 (9)
H1W	0.613 (6)	1.087 (8)	0.8854 (11)	0.08 (2)*
H2W	0.661 (5)	1.128 (7)	0.806 (3)	0.08 (2)*
H3W	0.332 (5)	0.547 (8)	0.7874 (18)	0.09 (2)*
H4W	0.313 (4)	0.523 (6)	0.6913 (15)	0.040 (15)*
H5W	0.484 (6)	0.5513 (17)	0.606 (4)	0.08 (2)*

supplementary materials

H6W	0.524 (8)	0.714 (7)	0.569 (3)	0.12 (3)*
H7W	0.1510 (15)	0.841 (6)	0.945 (3)	0.057 (19)*
H8W	0.032 (5)	0.745 (8)	0.888 (4)	0.16 (4)*
N1	0.6032 (3)	0.7004 (4)	0.8981 (2)	0.0151 (7)
O1	0.7200 (3)	0.6723 (4)	0.76661 (19)	0.0274 (8)
O2	0.8899 (3)	0.4962 (4)	0.84514 (18)	0.0234 (7)
O3	0.3689 (3)	0.8732 (4)	0.85503 (18)	0.0201 (6)
O4	0.3418 (3)	0.8625 (4)	0.99379 (18)	0.0249 (7)
O5	0.8051 (3)	0.5493 (4)	1.15748 (17)	0.0200 (6)
O6	0.6030 (3)	1.0722 (4)	0.8286 (2)	0.0315 (8)
O7	0.3636 (3)	0.5656 (4)	0.7411 (2)	0.0295 (8)
O8	0.4963 (4)	0.6642 (4)	0.6113 (2)	0.0255 (7)
O9	0.0620 (4)	0.8221 (4)	0.9302 (3)	0.0392 (9)

Atomic displacement parameters (\AA^2)

	U^{11}	U^{22}	U^{33}	U^{12}	U^{13}	U^{23}
Ho1	0.01235 (13)	0.01734 (13)	0.01046 (14)	0.00009 (7)	0.00339 (8)	0.00057 (8)
C1	0.016 (2)	0.024 (2)	0.014 (2)	0.0019 (17)	0.0032 (17)	-0.0012 (17)
C2	0.015 (2)	0.018 (2)	0.014 (2)	0.0028 (16)	0.0033 (17)	-0.0011 (17)
C3	0.016 (2)	0.018 (2)	0.017 (2)	0.0040 (16)	0.0039 (17)	0.0012 (16)
C4	0.017 (2)	0.019 (2)	0.013 (2)	-0.0047 (16)	0.0028 (17)	0.0022 (17)
C5	0.019 (2)	0.022 (2)	0.019 (2)	-0.0011 (16)	0.0079 (18)	0.0022 (17)
C6	0.016 (2)	0.017 (2)	0.014 (2)	0.0005 (16)	0.0049 (17)	0.0004 (16)
C7	0.016 (2)	0.015 (2)	0.018 (2)	-0.0025 (15)	0.0059 (18)	-0.0032 (16)
N1	0.0145 (18)	0.0198 (19)	0.0126 (18)	0.0011 (13)	0.0063 (14)	0.0005 (13)
O1	0.0270 (17)	0.044 (2)	0.0138 (16)	0.0158 (14)	0.0108 (14)	0.0074 (13)
O2	0.0244 (17)	0.0273 (18)	0.0196 (16)	0.0151 (13)	0.0076 (13)	0.0013 (13)
O3	0.0150 (15)	0.0300 (17)	0.0150 (16)	0.0057 (12)	0.0033 (12)	0.0029 (12)
O4	0.0211 (16)	0.0410 (19)	0.0151 (16)	0.0065 (13)	0.0092 (13)	0.0003 (13)
O5	0.0158 (15)	0.0295 (17)	0.0124 (15)	-0.0036 (12)	-0.0005 (12)	0.0065 (12)
O6	0.036 (2)	0.041 (2)	0.021 (2)	-0.0177 (15)	0.0143 (16)	-0.0087 (15)
O7	0.0341 (19)	0.036 (2)	0.0202 (18)	-0.0165 (14)	0.0105 (16)	-0.0040 (15)
O8	0.036 (2)	0.0232 (19)	0.0186 (18)	-0.0012 (14)	0.0098 (15)	-0.0031 (13)
O9	0.027 (2)	0.029 (2)	0.059 (3)	-0.0001 (15)	0.0072 (19)	0.0033 (18)

Geometric parameters (\AA , $^\circ$)

Ho1—O5 ⁱ	2.279 (3)	C5—C6	1.373 (5)
Ho1—O6	2.342 (3)	C5—H5	0.9300
Ho1—O7	2.354 (3)	C6—N1	1.341 (5)
Ho1—O1	2.386 (3)	C6—C7	1.508 (5)
Ho1—O2 ⁱⁱ	2.393 (3)	C7—O4	1.243 (5)
Ho1—O8	2.397 (3)	C7—O3	1.265 (5)
Ho1—O3	2.400 (3)	O2—Ho1 ⁱⁱⁱ	2.393 (3)
Ho1—N1	2.450 (3)	O5—Ho1 ^{iv}	2.279 (3)
C1—O2	1.249 (5)	O6—H1W	0.860 (10)
C1—O1	1.275 (5)	O6—H2W	0.858 (10)

C1—C2	1.493 (5)	O7—H3W	0.865 (10)
C2—N1	1.334 (5)	O7—H4W	0.861 (10)
C2—C3	1.361 (5)	O8—H5W	0.862 (10)
C3—C4	1.399 (5)	O8—H6W	0.861 (10)
C3—H3	0.9300	O9—H7W	0.860 (10)
C4—O5	1.322 (4)	O9—H8W	0.861 (10)
C4—C5	1.398 (5)		
O5 ⁱ —H01—O6	98.86 (11)	C3—C2—C1	122.7 (3)
O5 ⁱ —H01—O7	86.39 (12)	C2—C3—C4	119.7 (4)
O6—H01—O7	147.96 (12)	C2—C3—H3	120.2
O5 ⁱ —H01—O1	150.97 (10)	C4—C3—H3	120.2
O6—H01—O1	94.04 (12)	O5—C4—C5	122.5 (4)
O7—H01—O1	96.31 (12)	O5—C4—C3	121.1 (4)
O5 ⁱ —H01—O2 ⁱⁱ	81.27 (10)	C5—C4—C3	116.5 (4)
O6—H01—O2 ⁱⁱ	71.09 (11)	C6—C5—C4	119.9 (4)
O7—H01—O2 ⁱⁱ	140.78 (10)	C6—C5—H5	120.0
O1—H01—O2 ⁱⁱ	78.49 (10)	C4—C5—H5	120.0
O5 ⁱ —H01—O8	82.29 (11)	N1—C6—C5	122.8 (4)
O6—H01—O8	140.74 (12)	N1—C6—C7	113.0 (3)
O7—H01—O8	71.19 (12)	C5—C6—C7	124.2 (4)
O1—H01—O8	71.43 (11)	O4—C7—O3	124.8 (4)
O2 ⁱⁱ —H01—O8	70.33 (11)	O4—C7—C6	119.4 (4)
O5 ⁱ —H01—O3	79.50 (10)	O3—C7—C6	115.8 (3)
O6—H01—O3	74.68 (11)	C2—N1—C6	117.2 (3)
O7—H01—O3	75.28 (11)	C2—N1—H01	121.0 (3)
O1—H01—O3	129.20 (9)	C6—N1—H01	121.4 (2)
O2 ⁱⁱ —H01—O3	137.27 (9)	C1—O1—H01	124.0 (2)
O8—H01—O3	142.60 (11)	C1—O2—H01 ⁱⁱⁱ	139.4 (3)
O5 ⁱ —H01—N1	143.51 (10)	C7—O3—H01	124.8 (2)
O6—H01—N1	77.69 (12)	C4—O5—H01 ^{iv}	127.1 (2)
O7—H01—N1	79.53 (11)	H01—O6—H1W	124 (3)
O1—H01—N1	64.77 (10)	H01—O6—H2W	115 (3)
O2 ⁱⁱ —H01—N1	129.28 (10)	H1W—O6—H2W	116 (2)
O8—H01—N1	123.29 (11)	H01—O7—H3W	116 (3)
O3—H01—N1	64.44 (10)	H01—O7—H4W	123 (3)
O2—C1—O1	124.2 (4)	H3W—O7—H4W	114.8 (19)
O2—C1—C2	119.7 (3)	H01—O8—H5W	123 (3)
O1—C1—C2	116.1 (3)	H01—O8—H6W	120 (4)
N1—C2—C3	123.9 (4)	H5W—O8—H6W	115.3 (19)
N1—C2—C1	113.4 (3)	H7W—O9—H8W	116 (2)
O2—C1—C2—N1	-173.7 (4)	O3—H01—N1—C2	-179.5 (3)
O1—C1—C2—N1	8.7 (5)	O5 ⁱ —H01—N1—C6	-15.3 (4)
O2—C1—C2—C3	9.1 (6)	O6—H01—N1—C6	72.9 (3)
O1—C1—C2—C3	-168.6 (4)	O7—H01—N1—C6	-84.4 (3)
N1—C2—C3—C4	-1.0 (6)	O1—H01—N1—C6	173.5 (3)

supplementary materials

C1—C2—C3—C4	175.9 (4)	O2 ⁱⁱ —Ho1—N1—C6	125.6 (3)
C2—C3—C4—O5	−177.4 (4)	O8—Ho1—N1—C6	−143.5 (3)
C2—C3—C4—C5	2.3 (6)	O3—Ho1—N1—C6	−5.8 (3)
O5—C4—C5—C6	177.7 (4)	O2—C1—O1—Ho1	172.6 (3)
C3—C4—C5—C6	−2.0 (6)	C2—C1—O1—Ho1	−9.9 (5)
C4—C5—C6—N1	0.3 (6)	O5 ⁱ —Ho1—O1—C1	−163.5 (3)
C4—C5—C6—C7	−178.9 (4)	O6—Ho1—O1—C1	80.1 (3)
N1—C6—C7—O4	177.2 (3)	O7—Ho1—O1—C1	−69.5 (3)
C5—C6—C7—O4	−3.4 (6)	O2 ⁱⁱ —Ho1—O1—C1	149.8 (3)
N1—C6—C7—O3	−2.4 (5)	O8—Ho1—O1—C1	−137.3 (3)
C5—C6—C7—O3	176.9 (4)	O3—Ho1—O1—C1	6.6 (4)
C3—C2—N1—C6	−0.8 (6)	N1—Ho1—O1—C1	5.7 (3)
C1—C2—N1—C6	−178.0 (3)	O1—C1—O2—Ho1 ⁱⁱⁱ	−28.4 (7)
C3—C2—N1—Ho1	173.2 (3)	C2—C1—O2—Ho1 ⁱⁱⁱ	154.1 (3)
C1—C2—N1—Ho1	−4.0 (5)	O4—C7—O3—Ho1	177.4 (3)
C5—C6—N1—C2	1.1 (6)	C6—C7—O3—Ho1	−3.0 (5)
C7—C6—N1—C2	−179.6 (3)	O5 ⁱ —Ho1—O3—C7	178.8 (3)
C5—C6—N1—Ho1	−172.8 (3)	O6—Ho1—O3—C7	−78.9 (3)
C7—C6—N1—Ho1	6.5 (4)	O7—Ho1—O3—C7	89.8 (3)
O5 ⁱ —Ho1—N1—C2	171.0 (3)	O1—Ho1—O3—C7	3.6 (3)
O6—Ho1—N1—C2	−100.7 (3)	O2 ⁱⁱ —Ho1—O3—C7	−116.6 (3)
O7—Ho1—N1—C2	101.9 (3)	O8—Ho1—O3—C7	116.6 (3)
O1—Ho1—N1—C2	−0.2 (3)	N1—Ho1—O3—C7	4.5 (3)
O2 ⁱⁱ —Ho1—N1—C2	−48.1 (3)	C5—C4—O5—Ho1 ^{iv}	−106.4 (4)
O8—Ho1—N1—C2	42.8 (3)	C3—C4—O5—Ho1 ^{iv}	73.3 (4)


Symmetry codes: (i) $x-1/2, -y+3/2, z-1/2$; (ii) $-x+3/2, y+1/2, -z+3/2$; (iii) $-x+3/2, y-1/2, -z+3/2$; (iv) $x+1/2, -y+3/2, z+1/2$.

Hydrogen-bond geometry (\AA , °)

$D\cdots H$	$H\cdots A$	$D\cdots A$	$D\cdots H\cdots A$
O9—H8W ^v —O2 ^v	0.86 (1)	2.33 (4)	3.080 (5)
O9—H7W ^v —O4	0.86 (1)	1.84 (1)	2.693 (5)
O8—H6W ^{vi} —O9 ^{vi}	0.86 (1)	2.28 (3)	3.026 (5)
O8—H5W ^{vii} —O9 ^{vii}	0.86 (1)	1.84 (2)	2.689 (5)
O7—H4W ^{vii} —O3 ^{vii}	0.86 (1)	2.08 (3)	2.788 (4)
O7—H3W ^{viii} —O5 ^{viii}	0.87 (1)	1.92 (3)	2.711 (4)
O6—H2W ⁱⁱ —O1 ⁱⁱ	0.86 (1)	1.85 (2)	2.671 (4)
O6—H1W ^{ix} —O4 ^{ix}	0.86 (1)	1.83 (1)	2.687 (4)

Symmetry codes: (v) $x-1, y, z$; (vi) $x+1/2, -y+3/2, z-1/2$; (vii) $-x+1/2, y-1/2, -z+3/2$; (viii) $-x+1, -y+1, -z+2$; (ii) $-x+3/2, y+1/2, -z+3/2$; (ix) $-x+1, -y+2, -z+2$.

Fig. 1



supplementary materials

Fig. 2




Fig. 3

