

# 1,1-(Biphenyl-2,2'-diylldioxy)-3,3,5,5-tetrakis(4-bromomethylphenoxy)cyclotriphosphazene

Rui Han, Mei-Mei Chai, Jun-Liang Yang and Yong Ye\*

Phosphorus Chemical Engineering Research Center of Henan Province, Department of Chemistry, Zhengzhou University, Zhengzhou 450052, People's Republic of China

Correspondence e-mail: yeyong@zzu.edu.cn


Received 6 April 2011; accepted 27 April 2011

Key indicators: single-crystal X-ray study;  $T = 293$  K; mean  $\sigma(\text{C}-\text{C}) = 0.009$  Å; R factor = 0.077;  $wR$  factor = 0.192; data-to-parameter ratio = 19.8.

In the title compound,  $\text{C}_{40}\text{H}_{32}\text{Br}_4\text{N}_3\text{O}_6\text{P}_3$ , the cyclotriphosphazene ring adopts a planar conformation, with an r.m.s. deviation of 0.0247 Å. In the crystal, there is a weak intermolecular  $\text{C}-\text{H}\cdots\text{O}$  hydrogen bond as well as short intermolecular  $\text{Br}\cdots\text{Br}$  contacts [3.3352 (12) Å].

## Related literature

For general background to cyclotriphosphazenes, see: Manners (1996). For the applications of cyclotriphosphazene derivatives as flame retardants, see: Allcock (1977); as elastomers, see: Allcock (2000); as biomaterials, see: Trollsa & Hedrick (1998); as artificial nucleases, see: Wang, Ye, Zhong *et al.* (2009); Wang, Ye, Ju *et al.* (2009).



## Experimental

### Crystal data

$\text{C}_{40}\text{H}_{32}\text{Br}_4\text{N}_3\text{O}_6\text{P}_3$   
 $M_r = 1063.24$   
 Monoclinic,  $P2_1/n$   
 $a = 10.991$  (2) Å  
 $b = 28.417$  (6) Å  
 $c = 14.008$  (3) Å  
 $\beta = 105.47$  (3)°

$V = 4216.7$  (15) Å<sup>3</sup>  
 $Z = 4$   
 Mo  $K\alpha$  radiation  
 $\mu = 3.98$  mm<sup>-1</sup>  
 $T = 293$  K  
 $0.24 \times 0.20 \times 0.18$  mm

### Data collection

Bruker P4 CCD diffractometer  
 Absorption correction: multi-scan  
 (SADABS; Bruker, 2008)  
 $T_{\min} = 0.392$ ,  $T_{\max} = 0.520$

52073 measured reflections  
 9978 independent reflections  
 6049 reflections with  $I > 2\sigma(I)$   
 $R_{\text{int}} = 0.075$

### Refinement

$R[F^2 > 2\sigma(F^2)] = 0.077$   
 $wR(F^2) = 0.192$   
 $S = 1.11$   
 9978 reflections

505 parameters  
 H-atom parameters constrained  
 $\Delta\rho_{\text{max}} = 0.82$  e Å<sup>-3</sup>  
 $\Delta\rho_{\text{min}} = -0.71$  e Å<sup>-3</sup>

**Table 1**

Hydrogen-bond geometry (Å, °).

$D-\text{H}\cdots A$	$D-\text{H}$	$\text{H}\cdots A$	$D\cdots A$	$D-\text{H}\cdots A$
$\text{C33}-\text{H33A}\cdots\text{O1}^1$	0.97	2.60	3.525 (8)	160

Symmetry code: (i)  $x + \frac{1}{2}, -y + \frac{1}{2}, z - \frac{1}{2}$ .

Data collection: XSCANS (Bruker, 2008); cell refinement: XSCANS; data reduction: SAINT; program(s) used to solve structure: SHELXS97 (Sheldrick, 2008); program(s) used to refine structure: SHELXL97 (Sheldrick, 2008); molecular graphics: SHELXTL (Sheldrick, 2008); software used to prepare material for publication: SHELXTL.

This work was supported by Natural Science Foundation of China (Grant Nos. 20972143, 20602032).

Supplementary data and figures for this paper are available from the IUCr electronic archives (Reference: ZS2109).

## References

- Allcock, H. R. (1977). *Angew. Chem. Int. Ed. Engl.* **16**, 147–150.  
 Allcock, H. R. (2000). *Polym. Prep.* **14**, 553–558.  
 Bruker (2008). XSCANS, SAINT and SADABS. Bruker AXS Inc., Madison, Wisconsin, USA.  
 Manners, I. (1996). *Angew. Chem. Int. Ed. Engl.* **35**, 1602–1607.  
 Sheldrick, G. M. (2008). *Acta Cryst.* **A64**, 112–122.  
 Trollsa, M. & Hedrick, J. L. (1998). *J. Am. Chem. Soc.* **120**, 4644–4648.  
 Wang, L., Ye, Y., Ju, Z., Zhong, S. & Zhao, Y. F. (2009). *Phosphorus Sulfur*, **184**, 1958–1963.  
 Wang, L., Ye, Y., Zhong, S., Zhang, D. & Zhao, Y. F. (2009). *Chem. J. Chin. Univ.* **30**, 493–496.

**supplementary materials**

*Acta Cryst.* (2011). E67, o1298 [ doi:10.1107/S1600536811016047 ]

## 1,1-(Biphenyl-2,2'-diyldioxy)-3,3,5,5-tetrakis(4-bromomethylphenoxy)cyclotriphosphazene

R. Han, M.-M. Chai, J.-L. Yang and Y. Ye

### Comment

Cyclotriphosphazenes are of considerable interest not only because of their wide spectrum of chemical and physical properties, but also their importance in synthetic chemistry (Manners, 1996). Different side-group substituents affect the chemical and physical properties of the ring systems in high polymers based on a phosphazene skeleton. Various cyclotriphosphazenes have been successfully developed for a variety of applications, such as flame retardants (Allcock, 1977), elastomers (Allcock, 2000) and biomaterials (Trollsa & Hedrick, 1998), achieved by varying the nature of the substituent side group. Recently, some polydentate cyclotriphosphazene ligands were reported, which showed good nuclease activity with hydrolytic cleavage ability (Wang, Ye, Zhong *et al.*, 2009; Wang, Ye, Ju *et al.*, 2009). To obtain more insight into the selective recognition and efficient cleavage of DNA by different metal complexes of cyclotriphosphazene, the title compound, C<sub>40</sub>H<sub>32</sub>Br<sub>4</sub>N<sub>3</sub>O<sub>6</sub>P<sub>3</sub> (I), was synthesized and its crystal structure is reported here.

In the crystal structure of (I) (Fig. 1), the part of cyclotriphosphazene ring adopts a planar conformation with an r.m.s. deviation of 0.0247 Å. There are short intermolecular Br $\cdots$ Br contacts: Br1 $\cdots$ Br2<sup>ii</sup> [3.3352 (12) Å] and Br3 $\cdots$ Br4<sup>iii</sup> [3.5868 (15) Å] [for symmetry codes: (ii)  $-x - 1/2, y - 1/2, -z + 1/2$ ; (iii)  $-x + 1/2, y - 1/2, -z + 1/2$ ]. The crystal structure is stabilized by weak intermolecular C—H $\cdots$ O hydrogen bond (Table 1).

### Experimental

To 30 ml of THF 2,2'-biphenyldioxy-3,3,5,5-tetrakis(4-hydroxymethylphenoxy)- cyclotriphosphazene (1.622 g, 2 mmol) was added under argon. Then, 1.1 ml of phosphorus tribromide in 10 ml of THF was added dropwise and the mixture was stirred for 5 h. The organic solvent was removed under reduced pressure and the residue was dissolved in CHCl<sub>3</sub>. The residue washed with aq. K<sub>2</sub>CO<sub>3</sub> and extracted with CHCl<sub>3</sub>, the organic layer was dried over Na<sub>2</sub>SO<sub>4</sub> and then evaporated under reduced pressure. The remaining residue was purified by silica gel column chromatography (CH<sub>2</sub>Cl<sub>2</sub>/PE) to provide title compound which was recrystallized from CH<sub>2</sub>Cl<sub>2</sub> and single crystals of (I) were obtained by slow evaporation.

### Refinement

All H atoms were placed in calculated positions, with C—H = 0.93 or 0.97 Å, and included in the final cycles of refinement using a riding model, with  $U_{\text{iso}}(\text{H}) = 1.2U_{\text{eq}}(\text{C})$ . There is a single reflection which is considered to be affected by the beamstop.

## Figures




Fig. 1. Molecular configuration and atom numbering scheme for (I), with displacement ellipsoids drawn at the 20% probability level.

## 1,1-(Biphenyl-2,2'-diylldioxy)-3,3,5,5-tetrakis(4-bromomethylphenoxy)cyclotriphosphazene

### Crystal data

$C_{40}H_{32}Br_4N_3O_6P_3$

$M_r = 1063.24$

Monoclinic,  $P2_1/n$

Hall symbol: -P 2yn

$a = 10.991 (2) \text{ \AA}$

$b = 28.417 (6) \text{ \AA}$

$c = 14.008 (3) \text{ \AA}$

$\beta = 105.47 (3)^\circ$

$V = 4216.7 (15) \text{ \AA}^3$

$Z = 4$

$F(000) = 2104$

$D_x = 1.675 \text{ Mg m}^{-3}$

Mo  $K\alpha$  radiation,  $\lambda = 0.71073 \text{ \AA}$

Cell parameters from 7462 reflections

$\theta = 2.1\text{--}27.9^\circ$

$\mu = 3.98 \text{ mm}^{-1}$

$T = 293 \text{ K}$

Prism, colorless

$0.24 \times 0.20 \times 0.18 \text{ mm}$

### Data collection

Bruker P4 CCD  
diffractometer

Radiation source: fine-focus sealed tube  
graphite

$\omega$  scans

Absorption correction: multi-scan  
(*SADABS*; Bruker, 2008)

$T_{\min} = 0.392$ ,  $T_{\max} = 0.520$

52073 measured reflections

9978 independent reflections

6049 reflections with  $I > 2\sigma(I)$

$R_{\text{int}} = 0.075$

$\theta_{\max} = 27.9^\circ$ ,  $\theta_{\min} = 1.7^\circ$

$h = -14 \rightarrow 14$

$k = -37 \rightarrow 37$

$l = -18 \rightarrow 18$

### Refinement

Refinement on  $F^2$

Least-squares matrix: full

$R[F^2 > 2\sigma(F^2)] = 0.077$

$wR(F^2) = 0.192$

$S = 1.11$

9978 reflections

Primary atom site location: structure-invariant direct methods

Secondary atom site location: difference Fourier map

Hydrogen site location: inferred from neighbouring sites

H-atom parameters constrained

$w = 1/[\sigma^2(F_o^2) + (0.0761P)^2]$

where  $P = (F_o^2 + 2F_c^2)/3$

$(\Delta/\sigma)_{\max} = 0.001$

505 parameters

$$\Delta\rho_{\max} = 0.82 \text{ e } \text{\AA}^{-3}$$

0 restraints

$$\Delta\rho_{\min} = -0.71 \text{ e } \text{\AA}^{-3}$$

### Special details

**Geometry.** All e.s.d.'s (except the e.s.d. in the dihedral angle between two l.s. planes) are estimated using the full covariance matrix. The cell e.s.d.'s are taken into account individually in the estimation of e.s.d.'s in distances, angles and torsion angles; correlations between e.s.d.'s in cell parameters are only used when they are defined by crystal symmetry. An approximate (isotropic) treatment of cell e.s.d.'s is used for estimating e.s.d.'s involving l.s. planes.

**Refinement.** Refinement of  $F^2$  against ALL reflections. The weighted  $R$ -factor  $wR$  and goodness of fit  $S$  are based on  $F^2$ , conventional  $R$ -factors  $R$  are based on  $F$ , with  $F$  set to zero for negative  $F^2$ . The threshold expression of  $F^2 > \sigma(F^2)$  is used only for calculating  $R$ -factors(gt) *etc.* and is not relevant to the choice of reflections for refinement.  $R$ -factors based on  $F^2$  are statistically about twice as large as those based on  $F$ , and  $R$ -factors based on ALL data will be even larger.

### Fractional atomic coordinates and isotropic or equivalent isotropic displacement parameters ( $\text{\AA}^2$ )

	<i>x</i>	<i>y</i>	<i>z</i>	$U_{\text{iso}}^*/U_{\text{eq}}$
Br1	-0.08346 (6)	0.09289 (3)	0.36653 (6)	0.0732 (3)
Br2	-0.10180 (6)	0.59356 (3)	0.19304 (6)	0.0773 (3)
Br3	0.41308 (8)	0.13542 (3)	0.06958 (6)	0.0843 (3)
Br4	0.24692 (10)	0.53274 (3)	0.55072 (7)	0.0985 (3)
C1	-0.1461 (5)	0.3971 (2)	0.2173 (4)	0.0397 (12)
C2	-0.1438 (5)	0.4452 (2)	0.2072 (4)	0.0478 (14)
H2	-0.0683	0.4617	0.2283	0.057*
C3	-0.2555 (6)	0.4686 (2)	0.1653 (5)	0.0596 (17)
H3	-0.2547	0.5010	0.1558	0.072*
C4	-0.3689 (6)	0.4443 (2)	0.1372 (5)	0.0605 (17)
H4	-0.4442	0.4604	0.1114	0.073*
C5	-0.3695 (5)	0.3956 (2)	0.1477 (4)	0.0505 (15)
H5	-0.4457	0.3794	0.1284	0.061*
C6	-0.2570 (5)	0.37038 (19)	0.1869 (4)	0.0405 (13)
C7	-0.2574 (5)	0.3188 (2)	0.1952 (4)	0.0397 (12)
C8	-0.3569 (5)	0.2945 (2)	0.2200 (4)	0.0531 (15)
H8	-0.4232	0.3114	0.2329	0.064*
C9	-0.3574 (6)	0.2458 (2)	0.2256 (5)	0.0616 (17)
H9	-0.4243	0.2304	0.2410	0.074*
C10	-0.2592 (6)	0.2202 (2)	0.2083 (5)	0.0599 (17)
H10	-0.2593	0.1876	0.2134	0.072*
C11	-0.1597 (5)	0.2428 (2)	0.1833 (4)	0.0514 (15)
H11	-0.0936	0.2255	0.1713	0.062*
C12	-0.1602 (5)	0.2911 (2)	0.1766 (4)	0.0408 (13)
C13	0.2596 (5)	0.21640 (18)	0.3066 (4)	0.0394 (12)
C14	0.1915 (6)	0.1900 (2)	0.2275 (4)	0.0555 (16)
H14	0.1855	0.1997	0.1630	0.067*
C15	0.1329 (6)	0.1493 (2)	0.2442 (5)	0.0600 (17)
H15	0.0845	0.1322	0.1910	0.072*
C16	0.1454 (5)	0.13370 (19)	0.3407 (5)	0.0506 (15)

## supplementary materials

---

C17	0.2122 (6)	0.1612 (2)	0.4168 (5)	0.0582 (16)
H17	0.2182	0.1518	0.4815	0.070*
C18	0.2703 (6)	0.2019 (2)	0.4018 (4)	0.0551 (16)
H18	0.3163	0.2195	0.4552	0.066*
C19	0.0912 (6)	0.0870 (2)	0.3589 (6)	0.069 (2)
H19A	0.1424	0.0739	0.4204	0.082*
H19B	0.0942	0.0655	0.3058	0.082*
C20	0.2409 (5)	0.46520 (19)	0.1584 (4)	0.0443 (14)
C21	0.2325 (6)	0.4852 (2)	0.2471 (5)	0.0619 (17)
H21	0.2668	0.4699	0.3069	0.074*
C22	0.1733 (6)	0.5276 (2)	0.2451 (5)	0.0606 (17)
H22	0.1700	0.5415	0.3044	0.073*
C23	0.1184 (5)	0.5500 (2)	0.1571 (6)	0.0575 (17)
C24	0.1274 (5)	0.5302 (2)	0.0719 (5)	0.0545 (16)
H24	0.0915	0.5455	0.0123	0.065*
C25	0.1886 (5)	0.4877 (2)	0.0705 (4)	0.0505 (15)
H25	0.1943	0.4747	0.0109	0.061*
C26	0.0582 (6)	0.5975 (2)	0.1576 (7)	0.086 (3)
H26A	0.0436	0.6116	0.0924	0.104*
H26B	0.1156	0.6178	0.2045	0.104*
C27	0.2824 (5)	0.3118 (2)	0.0172 (4)	0.0453 (13)
C28	0.1792 (6)	0.2826 (2)	0.0044 (4)	0.0545 (15)
H28	0.1028	0.2937	0.0120	0.065*
C29	0.1932 (6)	0.2355 (2)	-0.0206 (4)	0.0580 (16)
H29	0.1246	0.2153	-0.0299	0.070*
C30	0.3068 (6)	0.2184 (2)	-0.0318 (4)	0.0518 (15)
C31	0.4050 (6)	0.2488 (2)	-0.0236 (5)	0.0591 (16)
H31	0.4797	0.2380	-0.0351	0.071*
C32	0.3950 (5)	0.2958 (2)	0.0020 (4)	0.0538 (15)
H32	0.4630	0.3162	0.0088	0.065*
C33	0.3209 (7)	0.1675 (2)	-0.0525 (5)	0.0693 (19)
H33A	0.3666	0.1641	-0.1025	0.083*
H33B	0.2383	0.1533	-0.0775	0.083*
C34	0.3556 (5)	0.3579 (2)	0.4642 (4)	0.0453 (14)
C35	0.4494 (6)	0.3902 (2)	0.4700 (5)	0.0605 (17)
H35	0.5128	0.3846	0.4388	0.073*
C36	0.4494 (7)	0.4315 (2)	0.5229 (5)	0.0670 (18)
H36	0.5131	0.4536	0.5274	0.080*
C37	0.3557 (8)	0.4398 (3)	0.5684 (5)	0.073 (2)
C38	0.2603 (8)	0.4073 (3)	0.5598 (5)	0.081 (2)
H38	0.1958	0.4134	0.5896	0.097*
C39	0.2586 (6)	0.3658 (2)	0.5074 (5)	0.0678 (19)
H39	0.1939	0.3440	0.5017	0.081*
C40	0.3541 (10)	0.4844 (3)	0.6263 (6)	0.104 (3)
H40A	0.4395	0.4964	0.6490	0.125*
H40B	0.3253	0.4771	0.6843	0.125*
N1	0.1362 (4)	0.31181 (15)	0.2933 (3)	0.0398 (10)
N2	0.3402 (4)	0.34123 (15)	0.2342 (3)	0.0408 (10)
N3	0.1042 (4)	0.37315 (15)	0.1405 (3)	0.0410 (11)

---

O1	-0.0320 (3)	0.37457 (12)	0.2693 (3)	0.0400 (9)
O2	-0.0652 (3)	0.31282 (12)	0.1399 (2)	0.0386 (8)
O3	0.3244 (3)	0.25632 (12)	0.2862 (3)	0.0473 (9)
O4	0.3618 (3)	0.31523 (13)	0.4151 (3)	0.0467 (9)
O5	0.3091 (4)	0.42359 (13)	0.1597 (4)	0.0642 (12)
O6	0.2786 (4)	0.35924 (13)	0.0467 (3)	0.0504 (10)
P1	0.04465 (12)	0.34309 (5)	0.21146 (10)	0.0365 (3)
P2	0.28306 (12)	0.30824 (5)	0.30222 (11)	0.0394 (3)
P3	0.25256 (13)	0.37218 (5)	0.14930 (11)	0.0419 (4)

*Atomic displacement parameters ( $\text{\AA}^2$ )*

	$U^{11}$	$U^{22}$	$U^{33}$	$U^{12}$	$U^{13}$	$U^{23}$
Br1	0.0496 (4)	0.0666 (5)	0.1086 (6)	-0.0111 (3)	0.0299 (4)	-0.0175 (4)
Br2	0.0483 (4)	0.0767 (5)	0.1062 (6)	0.0045 (3)	0.0193 (4)	-0.0218 (4)
Br3	0.1166 (7)	0.0600 (5)	0.0737 (5)	0.0086 (4)	0.0206 (5)	0.0056 (4)
Br4	0.1407 (8)	0.0686 (5)	0.0916 (6)	0.0241 (5)	0.0405 (6)	0.0054 (4)
C1	0.037 (3)	0.048 (3)	0.038 (3)	0.001 (2)	0.015 (2)	-0.009 (2)
C2	0.047 (3)	0.043 (3)	0.060 (4)	-0.004 (3)	0.026 (3)	-0.008 (3)
C3	0.081 (5)	0.044 (4)	0.063 (4)	0.013 (3)	0.034 (4)	0.001 (3)
C4	0.051 (4)	0.063 (4)	0.068 (4)	0.020 (3)	0.016 (3)	-0.001 (3)
C5	0.045 (3)	0.052 (4)	0.055 (4)	0.005 (3)	0.014 (3)	-0.008 (3)
C6	0.039 (3)	0.046 (3)	0.038 (3)	0.005 (2)	0.011 (2)	0.004 (2)
C7	0.032 (3)	0.047 (3)	0.040 (3)	-0.003 (2)	0.011 (2)	0.001 (2)
C8	0.044 (3)	0.069 (4)	0.048 (4)	-0.008 (3)	0.015 (3)	-0.001 (3)
C9	0.060 (4)	0.065 (4)	0.064 (4)	-0.031 (4)	0.024 (3)	-0.002 (3)
C10	0.080 (5)	0.039 (3)	0.060 (4)	-0.018 (3)	0.018 (3)	-0.001 (3)
C11	0.051 (3)	0.039 (3)	0.061 (4)	-0.004 (3)	0.011 (3)	-0.005 (3)
C12	0.031 (3)	0.049 (3)	0.041 (3)	-0.004 (2)	0.007 (2)	0.000 (2)
C13	0.035 (3)	0.036 (3)	0.049 (3)	0.004 (2)	0.014 (2)	-0.002 (2)
C14	0.073 (4)	0.048 (4)	0.041 (4)	0.001 (3)	0.008 (3)	0.004 (3)
C15	0.063 (4)	0.052 (4)	0.056 (4)	-0.007 (3)	0.001 (3)	-0.012 (3)
C16	0.039 (3)	0.036 (3)	0.079 (5)	0.003 (3)	0.020 (3)	0.000 (3)
C17	0.075 (4)	0.057 (4)	0.045 (4)	-0.013 (3)	0.020 (3)	0.005 (3)
C18	0.066 (4)	0.056 (4)	0.039 (4)	-0.013 (3)	0.007 (3)	-0.001 (3)
C19	0.063 (4)	0.048 (4)	0.107 (6)	-0.003 (3)	0.045 (4)	0.001 (4)
C20	0.038 (3)	0.032 (3)	0.066 (4)	-0.001 (2)	0.020 (3)	0.004 (3)
C21	0.075 (4)	0.063 (4)	0.050 (4)	-0.001 (4)	0.019 (3)	0.011 (3)
C22	0.064 (4)	0.054 (4)	0.067 (4)	-0.001 (3)	0.023 (3)	-0.019 (3)
C23	0.044 (3)	0.043 (3)	0.089 (5)	-0.004 (3)	0.024 (3)	-0.008 (4)
C24	0.050 (4)	0.044 (3)	0.062 (4)	-0.007 (3)	0.001 (3)	0.013 (3)
C25	0.056 (4)	0.052 (4)	0.045 (4)	-0.008 (3)	0.016 (3)	-0.009 (3)
C26	0.061 (4)	0.047 (4)	0.161 (8)	0.000 (3)	0.046 (5)	0.000 (5)
C27	0.052 (3)	0.047 (3)	0.038 (3)	0.002 (3)	0.015 (3)	0.006 (2)
C28	0.051 (4)	0.064 (4)	0.052 (4)	-0.003 (3)	0.020 (3)	0.001 (3)
C29	0.059 (4)	0.066 (4)	0.049 (4)	-0.014 (3)	0.014 (3)	0.000 (3)
C30	0.062 (4)	0.049 (4)	0.041 (3)	-0.006 (3)	0.008 (3)	-0.005 (3)
C31	0.065 (4)	0.058 (4)	0.057 (4)	0.012 (3)	0.022 (3)	-0.006 (3)

## supplementary materials

---

C32	0.041 (3)	0.062 (4)	0.057 (4)	-0.001 (3)	0.013 (3)	0.003 (3)
C33	0.101 (5)	0.049 (4)	0.054 (4)	0.008 (4)	0.014 (4)	-0.006 (3)
C34	0.047 (3)	0.047 (3)	0.040 (3)	0.005 (3)	0.008 (3)	-0.005 (3)
C35	0.059 (4)	0.059 (4)	0.063 (4)	-0.004 (3)	0.015 (3)	-0.002 (3)
C36	0.076 (5)	0.051 (4)	0.070 (5)	-0.004 (4)	0.012 (4)	0.007 (3)
C37	0.099 (6)	0.056 (4)	0.059 (4)	0.006 (4)	0.012 (4)	-0.005 (3)
C38	0.103 (6)	0.088 (6)	0.065 (5)	0.017 (5)	0.046 (4)	-0.016 (4)
C39	0.069 (4)	0.068 (5)	0.076 (5)	-0.013 (4)	0.035 (4)	-0.012 (4)
C40	0.181 (9)	0.056 (5)	0.068 (5)	0.022 (5)	0.022 (5)	-0.005 (4)
N1	0.033 (2)	0.042 (3)	0.047 (3)	0.0026 (19)	0.016 (2)	0.004 (2)
N2	0.031 (2)	0.035 (2)	0.059 (3)	-0.0023 (19)	0.018 (2)	0.002 (2)
N3	0.034 (2)	0.037 (2)	0.053 (3)	0.0041 (19)	0.014 (2)	0.010 (2)
O1	0.0313 (19)	0.043 (2)	0.047 (2)	0.0017 (16)	0.0130 (16)	-0.0061 (17)
O2	0.0345 (18)	0.043 (2)	0.042 (2)	-0.0042 (16)	0.0176 (16)	-0.0089 (16)
O3	0.048 (2)	0.032 (2)	0.067 (3)	0.0024 (17)	0.0243 (19)	-0.0022 (18)
O4	0.041 (2)	0.043 (2)	0.052 (2)	0.0017 (17)	0.0053 (18)	-0.0025 (18)
O5	0.045 (2)	0.032 (2)	0.125 (4)	0.0020 (18)	0.040 (2)	0.004 (2)
O6	0.062 (2)	0.044 (2)	0.057 (2)	0.0070 (19)	0.036 (2)	0.0122 (19)
P1	0.0320 (7)	0.0372 (7)	0.0431 (8)	-0.0004 (6)	0.0148 (6)	-0.0016 (6)
P2	0.0346 (7)	0.0346 (7)	0.0494 (9)	0.0015 (6)	0.0117 (6)	-0.0005 (6)
P3	0.0394 (8)	0.0330 (7)	0.0592 (10)	0.0005 (6)	0.0236 (7)	0.0036 (6)

### *Geometric parameters (Å, °)*

Br1—C19	1.958 (6)	C23—C24	1.347 (9)
Br2—C26	1.953 (7)	C23—C26	1.503 (8)
Br3—C33	1.963 (6)	C24—C25	1.385 (8)
Br4—C40	1.932 (8)	C24—H24	0.9300
C1—C2	1.374 (7)	C25—H25	0.9300
C1—C6	1.403 (7)	C26—H26A	0.9700
C1—O1	1.423 (6)	C26—H26B	0.9700
C2—C3	1.382 (8)	C27—C28	1.379 (8)
C2—H2	0.9300	C27—C32	1.387 (8)
C3—C4	1.388 (9)	C27—O6	1.413 (7)
C3—H3	0.9300	C28—C29	1.401 (9)
C4—C5	1.392 (8)	C28—H28	0.9300
C4—H4	0.9300	C29—C30	1.387 (8)
C5—C6	1.407 (7)	C29—H29	0.9300
C5—H5	0.9300	C30—C31	1.363 (8)
C6—C7	1.470 (7)	C30—C33	1.492 (8)
C7—C12	1.407 (7)	C31—C32	1.397 (8)
C7—C8	1.414 (7)	C31—H31	0.9300
C8—C9	1.385 (8)	C32—H32	0.9300
C8—H8	0.9300	C33—H33A	0.9700
C9—C10	1.375 (9)	C33—H33B	0.9700
C9—H9	0.9300	C34—C35	1.367 (8)
C10—C11	1.391 (8)	C34—C39	1.378 (8)
C10—H10	0.9300	C34—O4	1.405 (6)
C11—C12	1.374 (7)	C35—C36	1.389 (9)



C11—H11	0.9300	C35—H35	0.9300
C12—O2	1.422 (6)	C36—C37	1.368 (10)
C13—C18	1.370 (8)	C36—H36	0.9300
C13—C14	1.382 (7)	C37—C38	1.377 (10)
C13—O3	1.409 (6)	C37—C40	1.507 (10)
C14—C15	1.373 (8)	C38—C39	1.386 (9)
C14—H14	0.9300	C38—H38	0.9300
C15—C16	1.393 (9)	C39—H39	0.9300
C15—H15	0.9300	C40—H40A	0.9700
C16—C17	1.368 (8)	C40—H40B	0.9700
C16—C19	1.503 (8)	N1—P1	1.581 (4)
C17—C18	1.364 (8)	N1—P2	1.589 (4)
C17—H17	0.9300	N2—P2	1.581 (4)
C18—H18	0.9300	N2—P3	1.583 (4)
C19—H19A	0.9700	N3—P1	1.579 (4)
C19—H19B	0.9700	N3—P3	1.602 (4)
C20—C25	1.371 (8)	O1—P1	1.591 (4)
C20—C21	1.390 (8)	O2—P1	1.599 (4)
C20—O5	1.398 (6)	O3—P2	1.577 (4)
C21—C22	1.366 (9)	O4—P2	1.600 (4)
C21—H21	0.9300	O5—P3	1.579 (4)
C22—C23	1.377 (9)	O6—P3	1.582 (4)
C22—H22	0.9300		
C2—C1—C6	123.1 (5)	Br2—C26—H26A	109.5
C2—C1—O1	117.6 (5)	C23—C26—H26B	109.5
C6—C1—O1	119.1 (5)	Br2—C26—H26B	109.5
C1—C2—C3	118.9 (5)	H26A—C26—H26B	108.5
C1—C2—H2	120.0	C28—C27—C32	121.3 (6)
C3—C2—H2	120.0	C28—C27—O6	121.7 (5)
C2—C3—C4	120.6 (6)	C32—C27—O6	117.0 (5)
C4—C3—H3	120.0	C27—C28—C29	117.8 (6)
C2—C3—H3	120.0	C27—C28—H28	120.0
C3—C4—C5	119.7 (6)	C29—C28—H28	120.0
C3—C4—H4	120.0	C30—C29—C28	121.7 (6)
C5—C4—H4	120.0	C30—C29—H29	120.0
C4—C5—C6	121.2 (5)	C28—C29—H29	120.0
C4—C5—H5	120.0	C31—C30—C29	119.0 (6)
C6—C5—H5	120.0	C31—C30—C33	120.6 (6)
C1—C6—C5	116.4 (5)	C29—C30—C33	120.4 (6)
C1—C6—C7	122.5 (4)	C30—C31—C32	120.8 (6)
C5—C6—C7	121.1 (5)	C30—C31—H31	120.0
C12—C7—C8	116.5 (5)	C32—C31—H31	120.0
C12—C7—C6	121.7 (5)	C27—C32—C31	119.2 (6)
C8—C7—C6	121.8 (5)	C27—C32—H32	120.0
C9—C8—C7	121.1 (6)	C31—C32—H32	120.0
C9—C8—H8	120.0	C30—C33—Br3	109.7 (4)
C7—C8—H8	120.0	C30—C33—H33A	109.5
C10—C9—C8	120.2 (6)	Br3—C33—H33A	109.5
C10—C9—H9	120.0	C30—C33—H33B	109.5

## supplementary materials

---

C8—C9—H9	120.0	Br3—C33—H33B	109.5
C9—C10—C11	120.4 (6)	H33A—C33—H33B	108.5
C9—C10—H10	120.0	C35—C34—C39	121.6 (6)
C11—C10—H10	120.0	C35—C34—O4	118.3 (5)
C12—C11—C10	119.3 (6)	C39—C34—O4	120.1 (5)
C12—C11—H11	120.0	C34—C35—C36	119.5 (6)
C10—C11—H11	120.0	C34—C35—H35	120.0
C11—C12—C7	122.4 (5)	C36—C35—H35	120.0
C11—C12—O2	117.9 (5)	C37—C36—C35	120.1 (7)
C7—C12—O2	119.4 (5)	C37—C36—H36	120.0
C18—C13—C14	120.3 (5)	C35—C36—H36	120.0
C18—C13—O3	121.5 (5)	C36—C37—C38	119.6 (7)
C14—C13—O3	118.0 (5)	C36—C37—C40	120.9 (8)
C15—C14—C13	119.9 (6)	C38—C37—C40	119.5 (8)
C15—C14—H14	120.0	C37—C38—C39	121.2 (7)
C13—C14—H14	120.0	C37—C38—H38	120.0
C14—C15—C16	120.3 (6)	C39—C38—H38	120.0
C14—C15—H15	120.0	C34—C39—C38	118.0 (7)
C16—C15—H15	120.0	C34—C39—H39	120.0
C17—C16—C15	117.9 (5)	C38—C39—H39	120.0
C17—C16—C19	121.8 (6)	C37—C40—Br4	113.3 (5)
C15—C16—C19	120.3 (6)	C37—C40—H40A	109.5
C18—C17—C16	122.7 (6)	Br4—C40—H40A	109.5
C16—C17—H17	120.0	C37—C40—H40B	109.5
C18—C17—H17	120.0	Br4—C40—H40B	109.5
C17—C18—C13	118.9 (5)	H40A—C40—H40B	108.5
C13—C18—H18	120.0	P1—N1—P2	122.0 (3)
C17—C18—H18	120.0	P2—N2—P3	121.5 (2)
C16—C19—Br1	111.9 (4)	P1—N3—P3	121.8 (3)
C16—C19—H19A	109.5	C1—O1—P1	120.4 (3)
Br1—C19—H19A	109.5	C12—O2—P1	120.8 (3)
C16—C19—H19B	109.5	C13—O3—P2	123.0 (3)
Br1—C19—H19B	109.5	C34—O4—P2	120.5 (3)
H19A—C19—H19B	108.5	C20—O5—P3	125.8 (3)
C25—C20—C21	120.0 (5)	C27—O6—P3	120.9 (3)
C25—C20—O5	120.1 (5)	N3—P1—N1	118.0 (2)
C21—C20—O5	119.8 (5)	N3—P1—O1	112.4 (2)
C22—C21—C20	119.1 (6)	N1—P1—O1	105.5 (2)
C22—C21—H21	120.0	N3—P1—O2	105.5 (2)
C20—C21—H21	120.0	N1—P1—O2	112.3 (2)
C21—C22—C23	121.3 (6)	O1—P1—O2	101.99 (18)
C21—C22—H22	120.0	O3—P2—N2	107.0 (2)
C23—C22—H22	120.0	O3—P2—N1	111.9 (2)
C24—C23—C22	118.8 (6)	N2—P2—N1	118.2 (2)
C24—C23—C26	121.0 (7)	O3—P2—O4	99.0 (2)
C22—C23—C26	120.0 (7)	N2—P2—O4	108.9 (2)
C23—C24—C25	121.8 (6)	N1—P2—O4	109.9 (2)
C23—C24—H24	120.0	O5—P3—O6	97.9 (2)
C25—C24—H24	120.0	O5—P3—N2	107.4 (2)

C20—C25—C24	118.9 (6)	O6—P3—N2	109.4 (2)
C20—C25—H25	120.0	O5—P3—N3	111.0 (2)
C24—C25—H25	120.0	O6—P3—N3	111.2 (2)
C23—C26—Br2	112.2 (5)	N2—P3—N3	118.1 (2)
C23—C26—H26A	109.5		
O2—P1—O1—C1	-44.8 (4)	C5—C6—C7—C8	-36.7 (8)
N1—P1—O1—C1	-162.3 (4)	C1—C6—C7—C8	143.2 (6)
N3—P1—O1—C1	67.8 (4)	C6—C7—C12—O2	-6.0 (8)
O1—P1—O2—C12	-45.8 (4)	C12—C7—C8—C9	0.1 (8)
N1—P1—O2—C12	66.8 (4)	C8—C7—C12—C11	-0.9 (8)
N3—P1—O2—C12	-163.4 (4)	C6—C7—C12—C11	-179.5 (5)
N3—P1—N1—P2	-2.8 (4)	C8—C7—C12—O2	172.6 (5)
N1—P1—N3—P3	0.5 (4)	C6—C7—C8—C9	178.7 (5)
O1—P1—N3—P3	123.7 (3)	C7—C8—C9—C10	0.9 (9)
O2—P1—N3—P3	-126.0 (3)	C8—C9—C10—C11	-1.2 (10)
O1—P1—N1—P2	-129.4 (3)	C9—C10—C11—C12	0.4 (9)
O2—P1—N1—P2	120.2 (3)	C10—C11—C12—C7	0.7 (9)
N1—P2—O3—C13	-26.9 (5)	C10—C11—C12—O2	-172.9 (5)
O4—P2—O3—C13	88.9 (4)	O3—C13—C14—C15	176.3 (5)
O3—P2—N1—P1	-118.6 (3)	C18—C13—C14—C15	0.8 (9)
N2—P2—O3—C13	-158.0 (4)	O3—C13—C18—C17	-175.6 (5)
O3—P2—O4—C34	179.8 (4)	C14—C13—C18—C17	-0.3 (9)
N1—P2—O4—C34	-62.8 (4)	C13—C14—C15—C16	-2.2 (10)
N2—P2—O4—C34	68.2 (4)	C14—C15—C16—C19	-174.5 (6)
N1—P2—N2—P3	-8.1 (4)	C14—C15—C16—C17	3.0 (9)
N2—P2—N1—P1	6.5 (4)	C15—C16—C17—C18	-2.5 (10)
O3—P2—N2—P3	119.4 (3)	C19—C16—C17—C18	174.9 (6)
O4—P2—N1—P1	132.5 (3)	C15—C16—C19—Br1	-89.6 (7)
O4—P2—N2—P3	-134.4 (3)	C17—C16—C19—Br1	93.1 (7)
N2—P3—O5—C20	-132.4 (5)	C16—C17—C18—C13	1.2 (10)
N3—P3—O5—C20	-2.0 (6)	O5—C20—C21—C22	175.3 (6)
O6—P3—O5—C20	114.4 (5)	O5—C20—C25—C24	-176.5 (5)
O5—P3—N2—P2	132.2 (3)	C21—C20—C25—C24	-0.3 (9)
O6—P3—N2—P2	-122.6 (3)	C25—C20—C21—C22	-0.9 (9)
N3—P3—N2—P2	5.8 (4)	C20—C21—C22—C23	2.2 (10)
O5—P3—O6—C27	159.7 (4)	C21—C22—C23—C24	-2.2 (10)
N2—P3—O6—C27	48.1 (5)	C21—C22—C23—C26	-177.7 (6)
N3—P3—O6—C27	-84.1 (5)	C22—C23—C24—C25	1.0 (9)
O5—P3—N3—P1	-126.6 (3)	C22—C23—C26—Br2	-73.1 (7)
O6—P3—N3—P1	125.6 (3)	C24—C23—C26—Br2	111.5 (6)
N2—P3—N3—P1	-2.0 (4)	C26—C23—C24—C25	176.4 (6)
P1—O1—C1—C2	-108.8 (5)	C23—C24—C25—C20	0.3 (9)
P1—O1—C1—C6	76.9 (6)	C32—C27—C28—C29	2.6 (8)
P1—O2—C12—C7	75.7 (6)	O6—C27—C28—C29	-177.2 (5)
P1—O2—C12—C11	-110.5 (5)	O6—C27—C32—C31	177.7 (5)
P2—O3—C13—C18	-74.7 (6)	C28—C27—C32—C31	-2.1 (8)
P2—O3—C13—C14	109.9 (5)	C27—C28—C29—C30	0.3 (8)
P2—O4—C34—C39	84.1 (6)	C28—C29—C30—C33	176.4 (5)
P2—O4—C34—C35	-97.7 (6)	C28—C29—C30—C31	-3.7 (8)

## supplementary materials

---

P3—O5—C20—C21	94.2 (7)	C33—C30—C31—C32	-175.9 (6)
P3—O5—C20—C25	-89.6 (7)	C29—C30—C31—C32	4.1 (9)
P3—O6—C27—C32	-118.7 (5)	C29—C30—C33—Br3	-101.7 (6)
P3—O6—C27—C28	61.1 (7)	C31—C30—C33—Br3	78.3 (7)
O1—C1—C6—C5	171.9 (5)	C30—C31—C32—C27	-1.3 (9)
C6—C1—C2—C3	0.1 (9)	O4—C34—C35—C36	-176.4 (5)
O1—C1—C2—C3	-174.1 (5)	C39—C34—C35—C36	1.8 (10)
O1—C1—C6—C7	-8.0 (8)	O4—C34—C39—C38	176.4 (9)
C2—C1—C6—C5	-2.2 (8)	C35—C34—C39—C38	-1.7 (9)
C2—C1—C6—C7	178.0 (5)	C34—C35—C36—C37	-0.3 (10)
C1—C2—C3—C4	2.4 (9)	C35—C36—C37—C38	-1.2 (11)
C2—C3—C4—C5	-2.6 (10)	C35—C36—C37—C40	179.9 (7)
C3—C4—C5—C6	0.3 (9)	C36—C37—C38—C39	1.3 (11)
C4—C5—C6—C7	-178.2 (5)	C40—C37—C38—C39	-179.8 (7)
C4—C5—C6—C1	1.9 (8)	C36—C37—C40—Br4	96.5 (9)
C1—C6—C7—C12	-38.3 (8)	C38—C37—C40—Br4	-82.5 (8)
C5—C6—C7—C12	141.9 (6)	C37—C38—C39—C34	0.2 (10)

### Hydrogen-bond geometry ( $\text{\AA}$ , $^\circ$ )

$D-H\cdots A$	$D-H$	$H\cdots A$	$D\cdots A$	$D-H\cdots A$
C33—H33A $\cdots$ O1 <sup>i</sup>	0.97	2.60	3.525 (8)	160

Symmetry codes: (i)  $x+1/2, -y+1/2, z-1/2$ .

Fig. 1

