

N-(7-Ethoxy-1*H*-indazol-4-yl)-4-methylbenzenesulfonamide

Najat Abbassi,^a El Mostapha Rakib^{a*} and Hafid Zouihri^b

^aLaboratoire de Chimie Organique et Analytique, Université Sultan Moulay Slimane, Faculté des Sciences et Techniques, Béni-Mellal, BP 523, Morocco, and

^bLaboratoires de Diffraction des Rayons X, Centre Nationale pour la Recherche Scientifique et Technique, Rabat, Morocco

Correspondence e-mail: elmostapha1@ymail.com


Received 25 April 2011; accepted 2 May 2011

Key indicators: single-crystal X-ray study; $T = 296\text{ K}$, $P = 0.0\text{ kPa}$; mean $\sigma(\text{C}-\text{C}) = 0.003\text{ \AA}$; R factor = 0.053; wR factor = 0.158; data-to-parameter ratio = 32.1.

The molecule of the title heterocyclic compound, $\text{C}_{16}\text{H}_{17}\text{N}_3\text{O}_3\text{S}$, is bent at the S atom with an $\text{C}-\text{SO}_2-\text{NH}-\text{C}$ torsion angle of $80.17(8)^\circ$. The phenyl substituent at the S atom is rotated out of the plane of the 1H -indazole ring [interplanar angle = $46.24(8)^\circ$]. In the crystal, intermolecular $\text{N}-\text{H}\cdots\text{N}$ and $\text{N}-\text{H}\cdots\text{O}$ hydrogen bonds build up a ribbon developing parallel to the b -axis direction. $\text{C}-\text{H}\cdots\text{O}$ hydrogen bonds link these ribbons, forming a layer parallel to the bc plane.

Related literature

For related structures, see: Shakuntala *et al.* (2011a,b); Khan *et al.* (2010); Gowda *et al.* (2010). For the biological activity of similar sulfonamides, see: Soledade *et al.* (2006); Lee & Lee, (2002).



Experimental

Crystal data

$\text{C}_{16}\text{H}_{17}\text{N}_3\text{O}_3\text{S}$

$M_r = 331.39$

Monoclinic, $P2_1/c$
 $a = 16.2579(4)\text{ \AA}$
 $b = 5.0291(1)\text{ \AA}$
 $c = 20.4551(5)\text{ \AA}$
 $\beta = 97.269(1)^\circ$
 $V = 1659.02(7)\text{ \AA}^3$

$Z = 4$
 Mo $K\alpha$ radiation
 $\mu = 0.21\text{ mm}^{-1}$
 $T = 296\text{ K}$
 $0.23 \times 0.20 \times 0.14\text{ mm}$

Data collection

Bruker APEXII CCD detector
 diffractometer
 31243 measured reflections

6745 independent reflections
 5062 reflections with $I > 2\sigma(I)$
 $R_{\text{int}} = 0.025$

Refinement

$R[F^2 > 2\sigma(F^2)] = 0.053$
 $wR(F^2) = 0.158$
 $S = 1.07$
 6745 reflections

210 parameters
 H-atom parameters constrained
 $\Delta\rho_{\text{max}} = 0.42\text{ e \AA}^{-3}$
 $\Delta\rho_{\text{min}} = -0.25\text{ e \AA}^{-3}$

Table 1
 Hydrogen-bond geometry (\AA , $^\circ$).

$D-\text{H}\cdots A$	$D-\text{H}$	$\text{H}\cdots A$	$D\cdots A$	$D-\text{H}\cdots A$
C14—H14B \cdots O1 ⁱ	0.97	2.54	3.373 (3)	144
N1—H14 \cdots O2 ⁱⁱ	0.85	2.07	2.9159 (15)	172
N2—H2A \cdots N3 ⁱⁱⁱ	0.86	2.21	2.8974 (15)	136

Symmetry codes: (i) $x, -y + \frac{3}{2}, z + \frac{1}{2}$; (ii) $x, y - 1, z$; (iii) $-x + 1, -y + 3, -z + 1$.

Data collection: *APEX2* (Bruker, 2005); cell refinement: *SAINT* (Bruker, 2005); data reduction: *SAINT*; program(s) used to solve structure: *SHELXS97* (Sheldrick, 2008); program(s) used to refine structure: *SHELXL97* (Sheldrick, 2008); molecular graphics: *PLATON* (Spek, 2009); software used to prepare material for publication: *publCIF* (Westrip, 2010).

The authors thank the Unit of Support for Technical and Scientific Research (UATRS, CNRST) for the X-ray measurements.

Supplementary data and figures for this paper are available from the IUCr electronic archives (Reference: DN2681).

References

- Bruker (2005). *APEX2* and *SAINT*. Bruker AXS Inc., Madison, Wisconsin, USA.
- Gowda, B. T., Foro, S., Nirmala, P. G. & Fuess, H. (2010). *Acta Cryst. E66*, o1702.
- Khan, I. U., Ahmad, W., Arshad, M. N., Sharif, S. & Ahmed, J. (2010). *Acta Cryst. E66*, o2507.
- Lee, J. S. & Lee, C. H. (2002). *Bull. Korean Chem. Soc.* **23**, 167–169.
- Shakuntala, K., Foro, S. & Gowda, B. T. (2011a). *Acta Cryst. E67*, o104.
- Shakuntala, K., Foro, S. & Gowda, B. T. (2011b). *Acta Cryst. E67*, o142.
- Sheldrick, G. M. (2008). *Acta Cryst. A64*, 112–122.
- Soledade, M., Pedras, C. & Jha, M. (2006). *Bioorg. Med. Chem.* **14**, 4958–4979.
- Spek, A. L. (2009). *Acta Cryst. D65*, 148–155.
- Westrip, S. P. (2010). *J. Appl. Cryst.* **43**, 920–925.

supplementary materials

Acta Cryst. (2011). E67, o1354 [doi:10.1107/S1600536811016576]

N-(7-Ethoxy-1*H*-indazol-4-yl)-4-methylbenzenesulfonamide

N. Abbassi, E. M. Rakib and H. Zouihri

Comment

Similar sulfonamides have been studied in various previously works [Shakuntala *et al.* 2011a and 2011b; Khan *et al.* 2010; Gowda *et al.* 2010] and have proved important functionalities for biological and anti-hypertensive activities [Soledade *et al.*, 2006; Lee & Lee, 2002].

In the title compound, C₁₆H₁₇N₃O₃S, the molecule is bent at the S atom with an C—SO₂—NH—C torsion angle of 80.17 (8)° (Fig. 1). In the crystal structure, intermolecular N—H···N and N—H···O hydrogen bonds build up a ribbon developing parallel to the b direction and the C—H···O link these reibons to form a two D layer parallel to the bc plane (Fig. 2, Table 1).

The S atom has a distorted tetrahedral geometry [maximum deviation: O—S—O = 119.87 (9)°]. The phenyl substituent at S1 atom is rotated out of the plane of the 1*H*-indazol ring (the interplanar angles is 46.24 (8)°).

Experimental

A mixture of 4-nitroindazole (1.22 mmol) and anhydrous SnCl₂ (1.1 g, 6.1 mmol) in 25 mL of absolute ethanol was heated at 60 °C for 2 h. After reduction, the starting material disappeared, and the solution was allowed to cool down. The pH was made slightly basic (pH 7–8) by addition of 5% aqueous potassium bicarbonate before extraction with ethyl acetate. The organic phase was washed with brine and dried over magnesium sulfate. The solvent was removed to afford the amine, which was immediately dissolved in pyridine (5 ml) and then reacted with 4-methylbenzenesulfonyl chloride (0.26 g, 1.25 mmol) at room temperature for 24 h. After the reaction mixture was concentrated in vacuo, the resulting residue was purified by flash chromatography (eluted with Ethyl acetate: Hexane 1:9).

Yield: 45%; **mp:** 443–445 °K; **IR** (KBr, cm^{−1}): 3340, 3235 (NH), 1595 (CN), 1335, 1160 (SO₂); **¹H NMR** (**DMSO-d₆**): 2.27 (s, 3H, CH₃), 6.92 (dd, 1H, J=2.1 Hz and 6.1 Hz), 7.16 (d, 2H, J=6.2 Hz), 7.28 (d, 2H, J=8.3 Hz), 7.68 (d, 2H, J=8.3 Hz), 8.22 (s, 1H), 10.51 (s, 1H, NH), 13.05 (s, 1H, NH); **¹³C NMR** (**DMSO-d₆**): 21.3 (CH₃), 106.7, 111.0, 126.9, 127.2, 130.1, 132.4 (6 CH), 117.3, 130.5, 137.3, 141.4, 143.7 (5 C); **MS** m/z=288 [M+1].

Refinement

The H atoms bound to C were treated as riding with their parent atoms [C—H distances are 0.93 Å for CH groups and 0.96 Å for CH₂ with $U_{\text{iso}}(\text{H}) = 1.2 U_{\text{eq}}(\text{C})$, and 0.97 Å for CH₃ groups with $U_{\text{iso}}(\text{H}) = 1.5 U_{\text{eq}}(\text{C})$. The N₂—H₂₀ H atoms were treated as riding with $U_{\text{iso}}(\text{H}) = 1.2 U_{\text{eq}}(\text{N})$, and the N₁—H₁₀ H atoms were refined with restraints ($d_{\text{N}-\text{H}} = 0.88$ (2) Å) and then were treated as riding in the last cycles of refinement.

supplementary materials

Figures




Fig. 1. Molecular view of the title compound showing the atom-labeling scheme. Displacement ellipsoids are drawn at the 50% probability level. H atoms are represented as small spheres of arbitrary radii.




Fig. 2. Partial packing view showing the chain formed by C—H···O, N—H···O and N—H···N hydrogen bondings. H atoms not involved in hydrogen bonds have been omitted for clarity.

N-(7-Ethoxy-1*H*-indazol-4-yl)-4-methylbenzenesulfonamide

Crystal data

C ₁₆ H ₁₇ N ₃ O ₃ S	<i>F</i> (000) = 696
<i>M_r</i> = 331.39	<i>D_x</i> = 1.327 Mg m ⁻³
Monoclinic, <i>P</i> 2 ₁ /c	Melting point: 445 K
Hall symbol: -P 2ybc	Mo <i>K</i> α radiation, λ = 0.71073 Å
<i>a</i> = 16.2579 (4) Å	Cell parameters from 347 reflections
<i>b</i> = 5.0291 (1) Å	θ = 2.7–27.2°
<i>c</i> = 20.4551 (5) Å	μ = 0.21 mm ⁻¹
β = 97.269 (1)°	<i>T</i> = 296 K
<i>V</i> = 1659.02 (7) Å ³	Prism, yellow
<i>Z</i> = 4	0.23 × 0.20 × 0.14 mm

Data collection

Bruker APEXII CCD detector	5062 reflections with $I > 2\sigma(I)$
diffractometer	
Radiation source: fine-focus sealed tube	$R_{\text{int}} = 0.025$
graphite	$\theta_{\max} = 34.0^\circ$, $\theta_{\min} = 1.3^\circ$
ω and φ scans	$h = -25 \rightarrow 25$
31243 measured reflections	$k = -7 \rightarrow 7$
6745 independent reflections	$l = -32 \rightarrow 31$

Refinement

Refinement on F^2	Primary atom site location: structure-invariant direct methods
Least-squares matrix: full	Secondary atom site location: difference Fourier map
$R[F^2 > 2\sigma(F^2)] = 0.053$	Hydrogen site location: inferred from neighbouring sites
$wR(F^2) = 0.158$	H-atom parameters constrained

$S = 1.07$	$w = 1/[\sigma^2(F_o^2) + (0.0799P)^2 + 0.3561P]$
	where $P = (F_o^2 + 2F_c^2)/3$
6745 reflections	$(\Delta/\sigma)_{\max} = 0.011$
210 parameters	$\Delta\rho_{\max} = 0.42 \text{ e \AA}^{-3}$
0 restraints	$\Delta\rho_{\min} = -0.25 \text{ e \AA}^{-3}$

Special details

Geometry. All esds (except the esd in the dihedral angle between two l.s. planes) are estimated using the full covariance matrix. The cell esds are taken into account individually in the estimation of esds in distances, angles and torsion angles; correlations between esds in cell parameters are only used when they are defined by crystal symmetry. An approximate (isotropic) treatment of cell esds is used for estimating esds involving l.s. planes.

Refinement. Refinement of F^2 against ALL reflections. The weighted R -factor wR and goodness of fit S are based on F^2 , conventional R -factors R are based on F , with F set to zero for negative F^2 . The threshold expression of $F^2 > \sigma(F^2)$ is used only for calculating R -factors(gt) etc. and is not relevant to the choice of reflections for refinement. R -factors based on F^2 are statistically about twice as large as those based on F , and R -factors based on ALL data will be even larger.

Fractional atomic coordinates and isotropic or equivalent isotropic displacement parameters (\AA^2)

	x	y	z	$U_{\text{iso}}^*/U_{\text{eq}}$
S1	0.202461 (19)	0.91282 (6)	0.304901 (17)	0.03746 (10)
N1	0.27119 (7)	0.7334 (2)	0.35009 (6)	0.0391 (2)
H14	0.2635	0.5696	0.3412	0.047*
N2	0.43721 (7)	1.2628 (2)	0.50079 (6)	0.0384 (2)
H2A	0.4616	1.3343	0.5362	0.046*
N3	0.45193 (7)	1.3298 (3)	0.43916 (6)	0.0416 (3)
O1	0.19501 (7)	0.8007 (3)	0.24030 (5)	0.0519 (3)
O2	0.22674 (7)	1.1855 (2)	0.31483 (6)	0.0524 (3)
O3	0.37858 (8)	1.0205 (3)	0.61280 (6)	0.0578 (3)
C1	0.08572 (13)	1.0406 (4)	0.38431 (11)	0.0631 (5)
H1	0.1221	1.1725	0.4018	0.076*
C2	0.00971 (15)	1.0085 (5)	0.40702 (13)	0.0794 (7)
H2	-0.0048	1.1212	0.4398	0.095*
C3	-0.04508 (12)	0.8132 (5)	0.38215 (12)	0.0712 (6)
C4	-0.02207 (11)	0.6462 (5)	0.33422 (11)	0.0651 (5)
H4	-0.0581	0.5121	0.3174	0.078*
C5	0.05334 (10)	0.6742 (4)	0.31072 (9)	0.0515 (4)
H5	0.0681	0.5593	0.2785	0.062*
C6	0.10683 (8)	0.8748 (3)	0.33547 (7)	0.0385 (3)
C7	0.29527 (7)	0.8035 (2)	0.41776 (7)	0.0346 (2)
C8	0.26575 (9)	0.6722 (3)	0.46859 (8)	0.0451 (3)
H8	0.2269	0.5376	0.4590	0.054*
C9	0.29232 (10)	0.7346 (3)	0.53502 (8)	0.0501 (4)
H9	0.2716	0.6385	0.5682	0.060*
C10	0.34852 (9)	0.9356 (3)	0.55166 (7)	0.0415 (3)
C11	0.37907 (7)	1.0689 (2)	0.49948 (6)	0.0338 (2)
C12	0.35467 (7)	1.0062 (2)	0.43349 (6)	0.0323 (2)

supplementary materials

C13	0.40308 (8)	1.1763 (3)	0.39843 (7)	0.0381 (3)
H13	0.4008	1.1797	0.3528	0.046*
C14	0.36325 (16)	0.8578 (6)	0.66736 (10)	0.0814 (7)
H14A	0.3858	0.6811	0.6630	0.098*
H14B	0.3041	0.8422	0.6691	0.098*
C15	0.4038 (2)	0.9868 (8)	0.72770 (11)	0.1024 (9)
H15A	0.4610	1.0202	0.7232	0.154*
H15B	0.4005	0.8722	0.7648	0.154*
H15C	0.3765	1.1520	0.7344	0.154*
C16	-0.12866 (17)	0.7785 (10)	0.40632 (19)	0.1314 (14)
H16A	-0.1231	0.6624	0.4439	0.197*
H16B	-0.1671	0.7026	0.3718	0.197*
H16C	-0.1489	0.9484	0.4186	0.197*

Atomic displacement parameters (\AA^2)

	U^{11}	U^{22}	U^{33}	U^{12}	U^{13}	U^{23}
S1	0.03566 (15)	0.03045 (15)	0.04463 (19)	-0.00315 (11)	-0.00135 (12)	-0.00309 (12)
N1	0.0377 (5)	0.0267 (4)	0.0501 (6)	-0.0007 (4)	-0.0058 (4)	-0.0085 (4)
N2	0.0355 (5)	0.0403 (5)	0.0385 (5)	-0.0137 (4)	0.0015 (4)	-0.0025 (4)
N3	0.0373 (5)	0.0444 (6)	0.0427 (6)	-0.0155 (5)	0.0038 (4)	0.0018 (5)
O1	0.0506 (6)	0.0608 (7)	0.0429 (6)	-0.0006 (5)	0.0011 (4)	-0.0066 (5)
O2	0.0501 (6)	0.0289 (5)	0.0755 (8)	-0.0068 (4)	-0.0021 (5)	0.0035 (5)
O3	0.0651 (7)	0.0711 (8)	0.0380 (5)	-0.0210 (6)	0.0100 (5)	-0.0019 (5)
C1	0.0635 (10)	0.0507 (9)	0.0779 (13)	-0.0091 (8)	0.0196 (9)	-0.0217 (9)
C2	0.0760 (14)	0.0735 (13)	0.0963 (17)	-0.0007 (12)	0.0407 (13)	-0.0193 (13)
C3	0.0481 (9)	0.0836 (14)	0.0844 (14)	-0.0007 (9)	0.0185 (9)	0.0131 (12)
C4	0.0438 (8)	0.0742 (12)	0.0748 (12)	-0.0200 (8)	-0.0020 (8)	0.0052 (10)
C5	0.0452 (7)	0.0516 (8)	0.0561 (9)	-0.0130 (6)	-0.0006 (6)	-0.0089 (7)
C6	0.0352 (5)	0.0330 (6)	0.0457 (7)	-0.0009 (4)	-0.0012 (5)	-0.0021 (5)
C7	0.0296 (5)	0.0273 (5)	0.0457 (7)	-0.0034 (4)	-0.0001 (4)	-0.0027 (5)
C8	0.0384 (6)	0.0372 (6)	0.0587 (9)	-0.0146 (5)	0.0027 (6)	0.0028 (6)
C9	0.0454 (7)	0.0533 (8)	0.0528 (8)	-0.0178 (6)	0.0110 (6)	0.0081 (7)
C10	0.0382 (6)	0.0464 (7)	0.0404 (6)	-0.0082 (5)	0.0076 (5)	0.0017 (5)
C11	0.0288 (5)	0.0326 (5)	0.0400 (6)	-0.0053 (4)	0.0041 (4)	-0.0007 (5)
C12	0.0278 (4)	0.0299 (5)	0.0387 (6)	-0.0046 (4)	0.0019 (4)	-0.0013 (4)
C13	0.0360 (5)	0.0406 (6)	0.0375 (6)	-0.0094 (5)	0.0040 (4)	0.0004 (5)
C14	0.0892 (15)	0.1092 (19)	0.0474 (10)	-0.0299 (14)	0.0152 (10)	0.0098 (11)
C15	0.118 (2)	0.146 (3)	0.0443 (11)	-0.024 (2)	0.0118 (12)	-0.0018 (14)
C16	0.0683 (16)	0.171 (4)	0.165 (3)	-0.012 (2)	0.058 (2)	0.001 (3)

Geometric parameters (\AA , $^\circ$)

S1—O1	1.4277 (11)	C5—C6	1.3854 (19)
S1—O2	1.4345 (11)	C5—H5	0.9300
S1—N1	1.6296 (11)	C7—C8	1.369 (2)
S1—C6	1.7581 (14)	C7—C12	1.4133 (16)
N1—C7	1.4340 (17)	C8—C9	1.408 (2)
N1—H14	0.8492	C8—H8	0.9300

N2—N3	1.3550 (16)	C9—C10	1.376 (2)
N2—C11	1.3561 (15)	C9—H9	0.9300
N2—H2A	0.8600	C10—C11	1.4032 (18)
N3—C13	1.3241 (17)	C11—C12	1.3939 (17)
O3—C10	1.3529 (18)	C12—C13	1.4170 (17)
O3—C14	1.431 (2)	C13—H13	0.9300
C1—C6	1.378 (2)	C14—C15	1.474 (3)
C1—C2	1.384 (3)	C14—H14A	0.9700
C1—H1	0.9300	C14—H14B	0.9700
C2—C3	1.379 (3)	C15—H15A	0.9600
C2—H2	0.9300	C15—H15B	0.9600
C3—C4	1.378 (3)	C15—H15C	0.9600
C3—C16	1.514 (3)	C16—H16A	0.9600
C4—C5	1.379 (2)	C16—H16B	0.9600
C4—H4	0.9300	C16—H16C	0.9600
O1—S1—O2	119.88 (8)	C7—C8—H8	118.9
O1—S1—N1	106.15 (7)	C9—C8—H8	118.9
O2—S1—N1	107.00 (6)	C10—C9—C8	121.01 (13)
O1—S1—C6	108.12 (7)	C10—C9—H9	119.5
O2—S1—C6	107.07 (7)	C8—C9—H9	119.5
N1—S1—C6	108.17 (7)	O3—C10—C9	127.66 (14)
C7—N1—S1	119.69 (9)	O3—C10—C11	115.56 (12)
C7—N1—H14	117.4	C9—C10—C11	116.78 (13)
S1—N1—H14	110.1	N2—C11—C12	107.09 (11)
N3—N2—C11	111.41 (10)	N2—C11—C10	129.89 (12)
N3—N2—H2A	124.3	C12—C11—C10	122.95 (11)
C11—N2—H2A	124.3	C11—C12—C7	119.09 (11)
C13—N3—N2	106.15 (10)	C11—C12—C13	104.21 (10)
C10—O3—C14	117.55 (15)	C7—C12—C13	136.68 (12)
C6—C1—C2	119.19 (18)	N3—C13—C12	111.14 (12)
C6—C1—H1	120.4	N3—C13—H13	124.4
C2—C1—H1	120.4	C12—C13—H13	124.4
C3—C2—C1	121.5 (2)	O3—C14—C15	107.3 (2)
C3—C2—H2	119.3	O3—C14—H14A	110.3
C1—C2—H2	119.3	C15—C14—H14A	110.3
C4—C3—C2	118.40 (17)	O3—C14—H14B	110.3
C4—C3—C16	119.9 (2)	C15—C14—H14B	110.3
C2—C3—C16	121.7 (3)	H14A—C14—H14B	108.5
C3—C4—C5	121.26 (18)	C14—C15—H15A	109.5
C3—C4—H4	119.4	C14—C15—H15B	109.5
C5—C4—H4	119.4	H15A—C15—H15B	109.5
C4—C5—C6	119.43 (17)	C14—C15—H15C	109.5
C4—C5—H5	120.3	H15A—C15—H15C	109.5
C6—C5—H5	120.3	H15B—C15—H15C	109.5
C1—C6—C5	120.23 (15)	C3—C16—H16A	109.5
C1—C6—S1	120.31 (12)	C3—C16—H16B	109.5
C5—C6—S1	119.45 (12)	H16A—C16—H16B	109.5
C8—C7—C12	118.04 (12)	C3—C16—H16C	109.5
C8—C7—N1	122.39 (11)	H16A—C16—H16C	109.5

supplementary materials


C12—C7—N1	119.49 (11)	H16B—C16—H16C	109.5
C7—C8—C9	122.11 (12)		

Hydrogen-bond geometry (\AA , $^{\circ}$)

$D\cdots H$	$H\cdots A$	$D\cdots A$	$D\cdots H\cdots A$
C14—H14B \cdots O1 ⁱ	0.97	2.54	3.373 (3) 144
N1—H14 \cdots O2 ⁱⁱ	0.85	2.07	2.9159 (15) 172
N2—H2A \cdots N3 ⁱⁱⁱ	0.86	2.21	2.8974 (15) 136

Symmetry codes: (i) $x, -y+3/2, z+1/2$; (ii) $x, y-1, z$; (iii) $-x+1, -y+3, -z+1$.

Fig. 1



supplementary materials

Fig. 2

