

tert-Butyl N-[2-[bis(prop-2-yn-1-yl)-amino]phenyl]carbamate

Manavendra K. Singh,^a Alka Agarwal,^{a*} Charu Mahawar^a and Satish K. Awasthi^{b*}

^aDepartment of Medicinal Chemistry, Institute of Medical Sciences, Banaras Hindu University, Varanasi 225 001, U.P. India, and ^bChemical Biology Laboratory, Department of Chemistry, University of Delhi 110 007, Delhi, India
Correspondence e-mail: dralka@bhu.ac.in, awasthisatish@yahoo.com


Received 31 March 2011; accepted 4 May 2011

Key indicators: single-crystal X-ray study; $T = 293\text{ K}$; mean $\sigma(\text{C}-\text{C}) = 0.003\text{ \AA}$; R factor = 0.054; wR factor = 0.156; data-to-parameter ratio = 22.6.

In the crystal of the title compound, $\text{C}_{17}\text{H}_{20}\text{N}_2\text{O}_2$, the molecules are linked by $\text{C}-\text{H}\cdots\text{O}$ interactions. Intramolecular $\text{C}-\text{H}\cdots\text{O}$ and $\text{N}-\text{H}\cdots\text{N}$ hydrogen bonds also occur.

Related literature

For applications of alkyne scaffolds in biology, medicinal and materials chemistry, see: Diederich *et al.* (2005); Stang & Diederich (1995); Lam *et al.* (1988); Patai (1994). For background to click chemistry, which involves 1,3-dipolar cycloaddition of an alkyne with an azide and is an efficient and highly versatile tool that has allowed the preparation of a variety of macromolecule conjugates such as sugars, peptides or proteins and DNA, see: Rostovtsev *et al.* (2002). For the synthesis, see: Lilienkampf *et al.* (2009). For intermolecular interactions, see: Steiner & Desiraju (1998). For intramolecular $\text{C}-\text{H}\cdots\text{O}$ hydrogen bonds, see: Smith *et al.* (1993).



Experimental

Crystal data

$\text{C}_{17}\text{H}_{20}\text{N}_2\text{O}_2$

$M_r = 284.35$

Monoclinic, $C2/c$

$a = 19.1936(12)\text{ \AA}$

$b = 8.7181(4)\text{ \AA}$

$c = 19.7619(9)\text{ \AA}$

$\beta = 99.513(5)^\circ$

$V = 3261.3(3)\text{ \AA}^3$

$Z = 8$

Mo $K\alpha$ radiation

$\mu = 0.08\text{ mm}^{-1}$

$T = 293\text{ K}$

$0.40 \times 0.39 \times 0.38\text{ mm}$

Data collection

Oxford Diffraction Xcalibur Eos

diffractometer

Absorption correction: multi-scan
(*CrysAlis PRO*; Oxford

Diffraction, 2009)

$T_{\min} = 0.953$, $T_{\max} = 1.000$

6969 measured reflections

4389 independent reflections

2427 reflections with $I > 2\sigma(I)$

$R_{\text{int}} = 0.019$

Refinement

$R[F^2 > 2\sigma(F^2)] = 0.054$

$wR(F^2) = 0.156$

$S = 0.96$

4389 reflections

194 parameters

H atoms treated by a mixture of independent and constrained refinement

$\Delta\rho_{\max} = 0.19\text{ e \AA}^{-3}$

$\Delta\rho_{\min} = -0.19\text{ e \AA}^{-3}$

Table 1
Hydrogen-bond geometry (\AA , $^\circ$).

$D-\text{H}\cdots A$	$D-\text{H}$	$\text{H}\cdots A$	$D\cdots A$	$D-\text{H}\cdots A$
C10—H10B \cdots O1 ⁱ	0.97	2.55	3.512 (2)	171
C9—H9 \cdots O1 ⁱⁱ	0.93	2.28	3.194 (3)	166
C2—H2 \cdots O1	0.93	2.32	2.911 (3)	121
N1—H1N \cdots N2	0.810 (19)	2.28 (2)	2.703 (2)	114 (2)

Symmetry codes: (i) $-x + \frac{1}{2}, -y + \frac{3}{2}, -z + 1$; (ii) $x, -y + 2, z - \frac{1}{2}$.

Data collection: *CrysAlis PRO* (Oxford Diffraction, 2009); cell refinement: *CrysAlis PRO*; data reduction: *CrysAlis PRO*; program(s) used to solve structure: *SHELXS97* (Sheldrick, 2008); program(s) used to refine structure: *SHELXL97* (Sheldrick, 2008); molecular graphics: *Mercury* (Macrae *et al.*, 2006); software used to prepare material for publication: *publCIF* (Westrip, 2010).

AA, MKS and CM are thankful to the University Grant Commission (scheme No. 34–311/2008), New Delhi, and Banaras Hindu University, Varanasi, India, for financial assistance. SKA is thankful to the University of Delhi, India, for financial assistance.

Supplementary data and figures for this paper are available from the IUCr electronic archives (Reference: ZJ2008).

References

- Diederich, F., Stang, P. J. & Tykwinski, R. R. (2005). *Acetylene Chemistry: Chemistry, Biology, and Material Science*. Weinheim: Wiley-VCH.
- Lam, J., Bretel, H., Arnason, T. & Hansen, L. (1988). *Chemistry and Biology of Naturally-Occurring Acetylenes and Related Compounds*. Amsterdam: Elsevier.
- Lilienkampf, A., Mao, J., Wan, B., Wang, Y., Franzblau, S. G. & Kozikowski, A. P. (2009). *J. Med. Chem.* **52**, 2109–2118.
- Macrae, C. F., Edgington, P. R., McCabe, P., Pidcock, E., Shields, G. P., Taylor, R., Towler, M. & van de Streek, J. (2006). *J. Appl. Cryst.* **39**, 453–457.
- Oxford Diffraction (2009). *CrysAlis PRO*. Oxford Diffraction Ltd, Yarnton, England.
- Patai, S. (1994). *Chemistry of Triple-Bonded Functional Groups*. New York: Wiley.
- Rostovtsev, V. V., Green, L.-G., Fokin, V. V. & Sharpless, K. B. (2002). *Angew. Chem. Int. Ed.* **41**, 2596–2599.
- Sheldrick, G. M. (2008). *Acta Cryst. A* **64**, 112–122.
- Smith, B. D., Haller, K. J. & Shang, M. (1993). *J. Org. Chem.* **58**, 6905–6907.
- Stang, P. J. & Diederich, F. (1995). *Modern Acetylene Chemistry*. Weinheim: VCH.
- Steiner, T. & Desiraju, G. R. (1998). *Chem. Commun.* pp. 891–892.
- Westrip, S. P. (2010). *J. Appl. Cryst.* **43**, 920–925.

supplementary materials

Acta Cryst. (2011). E67, o1382 [doi:10.1107/S1600536811016862]

tert-Butyl N-[2-[bis(prop-2-yn-1-yl)amino]phenyl]carbamate

M. K. Singh, A. Agarwal, C. Mahawar and S. K. Awasthi

Comment

The carbon-carbon triple bond is an important and versatile functional group in organic chemistry. Alkynes are found in numerous natural products as well as in synthetic organic molecules. These alkyne scaffolds have various applications in biology, medicinal and material chemistry (Diederich *et al.*, 2005; Stang & Diederich 1995; Lam *et al.*, 1988; Patai 1994). Click chemistry developed by Sharpless (Rostovtsev *et al.*, 2002) involves 1,3-dipolar cycloaddition of alkyne with azide as an efficient and highly versatile tool that has allowed to prepare a variety of macromolecule conjugates such as sugars, peptides or proteins and DNA. As part of our ongoing work on antimicrobial studies on small molecules, we characterized and report here the crystal structure of [2-(di-prop-2-ynyl-amino)phenyl]carbamic acid *tert*-butyl ester (Figure1).

In the crystal structure, the compound is stabilized by intermolecular interaction between C10—H10B···O1 and C9—H9···O1 (Steiner & Desiraju, 1998) and intramolecular hydrogen bond between C2—H2···O1 (Smith *et al.*, 1993) and N1—HN1···N2 as seen in the crystal packing diagram along *b* axis (Table 1, Figure 2). Considering C1—C6 C13—C15 N1 N2 O1 O2 atom as plane A, C7 C8 C9 atom as plane B, C10 C11 C12 atom as plane C, the dihedral angels between planes A/B, A/C and B/C are 74.74°, 57.52°, 48.94° respectively, suggests that the molecule is not co-planar.

Experimental

The synthesis of the title compound was carried out according to the published procedure (Lilienkampf, *et al.*, 2009). Briefly, to a solution of (2-aminophenyl)carbamic acid *tert*-butyl ester (1.5 g, 7.2 mmol) in dry acetone was added anhydrous K₂CO₃ (7.95 g, 54.6 m mol) and reaction mixture was refluxed for 15–30 minutes. Subsequently, KI (0.60 g m, 3.6 mmol) and propargyl bromide (0.75 ml, 7.8 mmol) were added and further refluxed the reaction mixture for 18 hrs. The reaction mixture was cooled, filtered, and the filtrate was evaporated *in vacuo* to give the product. The crude product was purified by column chromatography using hexane and dichloromethane (65:35) as eluent. The purified product was recrystallized from hexane-dichloromethane (1:1). The colourless crystals were obtained by slow evaporation of solvent at room temperature in several days. Yield: 20%.

¹H NMR (CDCl₃): 8.11 (bs, 1H, NH), 7.56–7.55 (m, 1H, Ar—H), 7.35–7.32 (m, 1H, Ar—H), 7.19–7.14 (m, 1H, Ar—H), 6.99–6.94 (m, 1H, Ar—H), 3.83 (s, 4H, CH₂), 2.28 (s, 2H, CH), 1.51 (s, 9H, 3xCH₃).

Refinement

All H atoms were located from difference Fourier map (range of C—H = 0.93 - 0.97 Å, and N—H = 0.81 Å) and allowed to refine freely.

supplementary materials

Figures




Fig. 1. *ORTEP* diagram of molecule with thermal ellipsoids drawn at 50% probability level
Color code: White: C; red: O; blue: N; white: H




Fig. 2. Intermolecular interaction between C—H···O (blue line) and Intramolecular hydrogen bond (red line) showed in packing diagram of molecule along b-plane



Fig. 3. The formation of the title compound.

tert-Butyl N-[2-[bis(prop-2-yn-1-yl)amino]phenyl]carbamate

Crystal data

C ₁₇ H ₂₀ N ₂ O ₂	F(000) = 1216.0
M _r = 284.35	D _x = 1.158 Mg m ⁻³
Monoclinic, C2/c	Mo K α radiation, λ = 0.71073 Å
Hall symbol: -C 2yc	Cell parameters from 4389 reflections
a = 19.1936 (12) Å	θ = 3.2–29.0°
b = 8.7181 (4) Å	μ = 0.08 mm ⁻¹
c = 19.7619 (9) Å	T = 293 K
β = 99.513 (5)°	Block, colourless
V = 3261.3 (3) Å ³	0.40 × 0.39 × 0.38 mm
Z = 8	

Data collection

Oxford Diffraction Xcalibur Eos diffractometer	4389 independent reflections
Radiation source: fine-focus sealed tube graphite	2427 reflections with $I > 2\sigma(I)$
ω scans	$R_{\text{int}} = 0.019$
Absorption correction: multi-scan (<i>CrysAlis PRO</i> ; Oxford Diffraction, 2009)	$\theta_{\max} = 29.1^\circ$, $\theta_{\min} = 3.2^\circ$
$T_{\min} = 0.953$, $T_{\max} = 1.000$	$h = -24 \rightarrow 7$
6969 measured reflections	$k = -10 \rightarrow 10$
	$l = -23 \rightarrow 26$

Refinement

Refinement on F^2	Primary atom site location: structure-invariant direct methods
Least-squares matrix: full	Secondary atom site location: difference Fourier map
$R[F^2 > 2\sigma(F^2)] = 0.054$	Hydrogen site location: inferred from neighbouring sites
$wR(F^2) = 0.156$	H atoms treated by a mixture of independent and constrained refinement
$S = 0.96$	$w = 1/[\sigma^2(F_o^2) + (0.0754P)^2 + 1.325P]$ where $P = (F_o^2 + 2F_c^2)/3$
4389 reflections	$(\Delta/\sigma)_{\max} = 0.05$
194 parameters	$\Delta\rho_{\max} = 0.19 \text{ e \AA}^{-3}$
0 restraints	$\Delta\rho_{\min} = -0.19 \text{ e \AA}^{-3}$

Special details

Geometry. All e.s.d.'s (except the e.s.d. in the dihedral angle between two l.s. planes) are estimated using the full covariance matrix. The cell e.s.d.'s are taken into account individually in the estimation of e.s.d.'s in distances, angles and torsion angles; correlations between e.s.d.'s in cell parameters are only used when they are defined by crystal symmetry. An approximate (isotropic) treatment of cell e.s.d.'s is used for estimating e.s.d.'s involving l.s. planes.

Refinement. Refinement of F^2 against ALL reflections. The weighted R -factor wR and goodness of fit S are based on F^2 , conventional R -factors R are based on F , with F set to zero for negative F^2 . The threshold expression of $F^2 > \sigma(F^2)$ is used only for calculating R -factors(gt) etc. and is not relevant to the choice of reflections for refinement. R -factors based on F^2 are statistically about twice as large as those based on F , and R -factors based on ALL data will be even larger.

Fractional atomic coordinates and isotropic or equivalent isotropic displacement parameters (\AA^2)

	<i>x</i>	<i>y</i>	<i>z</i>	$U_{\text{iso}}^*/U_{\text{eq}}$
HN1	0.1292 (11)	0.706 (2)	0.4601 (9)	0.038 (5)*
N1	0.13858 (9)	0.78587 (19)	0.47962 (8)	0.0397 (4)
O2	0.05794 (7)	0.71231 (15)	0.53892 (6)	0.0505 (4)
N2	0.17096 (8)	0.68525 (17)	0.35922 (7)	0.0399 (4)
C1	0.18983 (9)	0.8707 (2)	0.45262 (8)	0.0362 (4)
O1	0.11848 (8)	0.93110 (16)	0.57014 (6)	0.0548 (4)
C13	0.10630 (10)	0.8202 (2)	0.53354 (8)	0.0380 (4)
C6	0.20658 (10)	0.8197 (2)	0.38966 (8)	0.0383 (4)
C2	0.22421 (11)	0.9975 (2)	0.48421 (10)	0.0454 (5)
H2	0.2130	1.0324	0.5256	0.055*
C11	0.19018 (12)	0.4288 (3)	0.31791 (9)	0.0510 (5)
C5	0.25792 (11)	0.8980 (2)	0.36113 (10)	0.0507 (5)
H5	0.2691	0.8655	0.3194	0.061*
C10	0.21577 (11)	0.5866 (2)	0.32379 (9)	0.0476 (5)
H10A	0.2168	0.6274	0.2783	0.057*
H10B	0.2637	0.5879	0.3489	0.057*
C8	0.10589 (12)	0.7963 (3)	0.25187 (10)	0.0586 (6)

supplementary materials

C3	0.27519 (12)	1.0729 (2)	0.45480 (11)	0.0544 (5)
H3	0.2980	1.1581	0.4765	0.065*
C7	0.10173 (11)	0.7167 (2)	0.31671 (9)	0.0491 (5)
H7A	0.0771	0.6202	0.3064	0.059*
H7B	0.0739	0.7786	0.3431	0.059*
C14	0.00843 (11)	0.7249 (2)	0.58831 (9)	0.0484 (5)
C4	0.29225 (12)	1.0225 (2)	0.39370 (12)	0.0576 (6)
H4	0.3270	1.0727	0.3744	0.069*
C12	0.17318 (14)	0.2999 (3)	0.31334 (12)	0.0681 (7)
H12	0.1597	0.1974	0.3097	0.082*
C15	-0.03735 (16)	0.5849 (3)	0.57144 (14)	0.0852 (9)
H15A	-0.0629	0.5935	0.5255	0.128*
H15B	-0.0701	0.5772	0.6031	0.128*
H15C	-0.0081	0.4949	0.5750	0.128*
C9	0.11141 (16)	0.8551 (4)	0.20009 (13)	0.0838 (8)
H9	0.1158	0.9020	0.1587	0.101*
C16	0.04878 (15)	0.7145 (4)	0.66033 (11)	0.0833 (9)
H16A	0.0750	0.6203	0.6656	0.125*
H16B	0.0163	0.7169	0.6924	0.125*
H16C	0.0808	0.7996	0.6689	0.125*
C17	-0.03353 (15)	0.8709 (3)	0.57635 (15)	0.0854 (9)
H17A	-0.0585	0.8727	0.5300	0.128*
H17B	-0.0021	0.9572	0.5837	0.128*
H17C	-0.0668	0.8759	0.6076	0.128*

Atomic displacement parameters (\AA^2)

	U^{11}	U^{22}	U^{33}	U^{12}	U^{13}	U^{23}
N1	0.0456 (10)	0.0393 (9)	0.0381 (8)	-0.0030 (7)	0.0187 (7)	-0.0042 (7)
O2	0.0517 (9)	0.0599 (8)	0.0465 (7)	-0.0109 (7)	0.0271 (6)	-0.0086 (6)
N2	0.0396 (9)	0.0496 (9)	0.0328 (7)	0.0035 (7)	0.0129 (6)	0.0014 (6)
C1	0.0340 (10)	0.0386 (9)	0.0379 (9)	0.0065 (8)	0.0118 (7)	0.0075 (7)
O1	0.0565 (9)	0.0634 (9)	0.0486 (7)	-0.0082 (7)	0.0212 (6)	-0.0186 (7)
C13	0.0358 (10)	0.0462 (10)	0.0332 (8)	0.0030 (8)	0.0094 (7)	0.0006 (7)
C6	0.0362 (10)	0.0424 (10)	0.0386 (9)	0.0062 (8)	0.0126 (7)	0.0083 (7)
C2	0.0431 (11)	0.0422 (10)	0.0530 (11)	0.0036 (9)	0.0133 (8)	-0.0008 (8)
C11	0.0530 (13)	0.0616 (14)	0.0412 (10)	0.0101 (11)	0.0160 (9)	-0.0043 (9)
C5	0.0506 (13)	0.0553 (12)	0.0518 (11)	0.0017 (10)	0.0250 (9)	0.0092 (9)
C10	0.0457 (12)	0.0610 (13)	0.0391 (9)	0.0088 (10)	0.0155 (8)	-0.0007 (8)
C8	0.0522 (14)	0.0731 (15)	0.0499 (12)	0.0075 (12)	0.0066 (9)	0.0173 (10)
C3	0.0455 (13)	0.0434 (11)	0.0756 (14)	-0.0007 (10)	0.0141 (10)	0.0011 (10)
C7	0.0418 (12)	0.0638 (13)	0.0432 (10)	0.0022 (10)	0.0113 (8)	0.0076 (9)
C14	0.0456 (12)	0.0621 (12)	0.0435 (10)	0.0018 (10)	0.0247 (8)	0.0024 (9)
C4	0.0466 (13)	0.0529 (12)	0.0794 (14)	-0.0020 (10)	0.0287 (11)	0.0122 (11)
C12	0.0762 (18)	0.0616 (15)	0.0727 (15)	0.0037 (13)	0.0307 (13)	-0.0087 (11)
C15	0.081 (2)	0.095 (2)	0.0936 (18)	-0.0271 (16)	0.0533 (15)	-0.0110 (15)
C9	0.081 (2)	0.108 (2)	0.0634 (15)	0.0147 (17)	0.0150 (13)	0.0378 (15)
C16	0.086 (2)	0.121 (2)	0.0472 (13)	0.0107 (18)	0.0253 (12)	0.0175 (13)

C17	0.0630 (17)	0.093 (2)	0.110 (2)	0.0268 (15)	0.0427 (15)	0.0300 (16)
-----	-------------	-----------	-----------	-------------	-------------	-------------

Geometric parameters (\AA , $^\circ$)

N1—C13	1.352 (2)	C8—C7	1.471 (3)
N1—C1	1.404 (2)	C3—C4	1.374 (3)
N1—HN1	0.801 (19)	C3—H3	0.9300
O2—C13	1.338 (2)	C7—H7A	0.9700
O2—C14	1.475 (2)	C7—H7B	0.9700
N2—C6	1.438 (2)	C14—C17	1.504 (3)
N2—C7	1.476 (2)	C14—C16	1.507 (3)
N2—C10	1.472 (2)	C14—C15	1.509 (3)
C1—C2	1.382 (3)	C4—H4	0.9300
C1—C6	1.407 (2)	C12—H12	0.9300
O1—C13	1.207 (2)	C15—H15A	0.9600
C6—C5	1.392 (3)	C15—H15B	0.9600
C2—C3	1.384 (3)	C15—H15C	0.9600
C2—H2	0.9300	C9—H9	0.9300
C11—C12	1.170 (3)	C16—H16A	0.9600
C11—C10	1.459 (3)	C16—H16B	0.9600
C5—C4	1.374 (3)	C16—H16C	0.9600
C5—H5	0.9300	C17—H17A	0.9600
C10—H10A	0.9700	C17—H17B	0.9600
C10—H10B	0.9700	C17—H17C	0.9600
C8—C9	1.164 (3)		
C13—N1—C1	128.60 (16)	C8—C7—H7A	108.7
C13—N1—HN1	118.4 (14)	N2—C7—H7A	108.7
C1—N1—HN1	113.0 (14)	C8—C7—H7B	108.7
C13—O2—C14	122.13 (14)	N2—C7—H7B	108.7
C6—N2—C7	114.07 (15)	H7A—C7—H7B	107.6
C6—N2—C10	113.60 (15)	O2—C14—C17	110.24 (16)
C7—N2—C10	112.31 (14)	O2—C14—C16	109.50 (18)
C2—C1—C6	119.35 (16)	C17—C14—C16	112.2 (2)
C2—C1—N1	124.15 (16)	O2—C14—C15	102.07 (16)
C6—C1—N1	116.50 (16)	C17—C14—C15	111.9 (2)
O1—C13—O2	125.71 (16)	C16—C14—C15	110.5 (2)
O1—C13—N1	125.54 (17)	C5—C4—C3	119.92 (19)
O2—C13—N1	108.74 (15)	C5—C4—H4	120.0
C5—C6—C1	118.91 (18)	C3—C4—H4	120.0
C5—C6—N2	123.34 (16)	C11—C12—H12	180.0
C1—C6—N2	117.72 (15)	C14—C15—H15A	109.5
C3—C2—C1	120.53 (18)	C14—C15—H15B	109.5
C3—C2—H2	119.7	H15A—C15—H15B	109.5
C1—C2—H2	119.7	C14—C15—H15C	109.5
C12—C11—C10	176.5 (2)	H15A—C15—H15C	109.5
C4—C5—C6	120.96 (19)	H15B—C15—H15C	109.5
C4—C5—H5	119.5	C8—C9—H9	180.0
C6—C5—H5	119.5	C14—C16—H16A	109.5
C11—C10—N2	111.90 (16)	C14—C16—H16B	109.5

supplementary materials


C11—C10—H10A	109.2	H16A—C16—H16B	109.5
N2—C10—H10A	109.2	C14—C16—H16C	109.5
C11—C10—H10B	109.2	H16A—C16—H16C	109.5
N2—C10—H10B	109.2	H16B—C16—H16C	109.5
H10A—C10—H10B	107.9	C14—C17—H17A	109.5
C9—C8—C7	177.1 (3)	C14—C17—H17B	109.5
C4—C3—C2	120.3 (2)	H17A—C17—H17B	109.5
C4—C3—H3	119.8	C14—C17—H17C	109.5
C2—C3—H3	119.8	H17A—C17—H17C	109.5
C8—C7—N2	114.21 (17)	H17B—C17—H17C	109.5
C13—N1—C1—C2	10.1 (3)	N1—C1—C2—C3	178.67 (18)
C13—N1—C1—C6	-170.50 (17)	C1—C6—C5—C4	0.4 (3)
C14—O2—C13—O1	5.5 (3)	N2—C6—C5—C4	-177.51 (18)
C14—O2—C13—N1	-173.54 (16)	C12—C11—C10—N2	-149 (4)
C1—N1—C13—O1	-2.8 (3)	C6—N2—C10—C11	157.11 (15)
C1—N1—C13—O2	176.26 (16)	C7—N2—C10—C11	-71.6 (2)
C2—C1—C6—C5	0.5 (3)	C1—C2—C3—C4	0.0 (3)
N1—C1—C6—C5	-178.92 (16)	C9—C8—C7—N2	46 (6)
C2—C1—C6—N2	178.52 (16)	C6—N2—C7—C8	70.5 (2)
N1—C1—C6—N2	-0.9 (2)	C10—N2—C7—C8	-60.6 (2)
C7—N2—C6—C5	-96.4 (2)	C13—O2—C14—C17	57.2 (3)
C10—N2—C6—C5	34.0 (2)	C13—O2—C14—C16	-66.8 (2)
C7—N2—C6—C1	85.68 (18)	C13—O2—C14—C15	176.18 (18)
C10—N2—C6—C1	-143.86 (16)	C6—C5—C4—C3	-1.1 (3)
C6—C1—C2—C3	-0.7 (3)	C2—C3—C4—C5	0.9 (3)

Hydrogen-bond geometry (\AA , $^\circ$)

$D—\text{H}\cdots A$	$D—\text{H}$	$\text{H}\cdots A$	$D\cdots A$	$D—\text{H}\cdots A$
C10—H10B \cdots O1 ⁱ	0.97	2.55	3.512 (2)	171
C9—H9 \cdots O1 ⁱⁱ	0.93	2.28	3.194 (3)	166
C2—H2 \cdots O1	0.93	2.32	2.911 (3)	121
N1—HN1 \cdots N2	0.810 (19)	2.28 (2)	2.703 (2)	114 (2)

Symmetry codes: (i) $-x+1/2, -y+3/2, -z+1$; (ii) $x, -y+2, z-1/2$.

Fig. 1



supplementary materials

Fig. 2




Fig. 3

