

# Methyl 3-(4-chlorophenyl)-2-(1,3-dimethyl-2,5-dioxo-4-phenylimidazolidin-4-yl)-3-oxopropanoate

Dongxue Zhang, Cong Deng and Yan Yang\*

Key Laboratory of Pesticide and Chemical Biology of the Ministry of Education, College of Chemistry, Central China Normal University, Wuhan 430079, People's Republic of China

Correspondence e-mail: hbdongxue@163.com


Received 13 April 2011; accepted 17 May 2011

Key indicators: single-crystal X-ray study;  $T = 298$  K; mean  $\sigma(\text{C}-\text{C}) = 0.004$  Å;  $R$  factor = 0.069;  $wR$  factor = 0.164; data-to-parameter ratio = 15.1.

The title compound,  $\text{C}_{21}\text{H}_{19}\text{ClN}_2\text{O}_5$ , is a tetrasubstituted hydantoin derivative which contains an imidazolidine-2,4-dione core. The dihedral angle between the aromatic rings is  $64.53(14)^\circ$ . In the crystal, weak intermolecular  $\text{C}-\text{H}\cdots\text{O}$  hydrogen bonding is found. An intramolecular  $\text{C}-\text{H}\cdots\text{O}$  interaction also occurs.

## Related literature

For the preparation of the title compound, see: Gao *et al.* (2010).



## Experimental

### Crystal data

$\text{C}_{21}\text{H}_{19}\text{ClN}_2\text{O}_5$	$V = 2040.4(3) \text{ \AA}^3$
$M_r = 414.83$	$Z = 4$
Monoclinic, $P2_1/n$	Mo $K\alpha$ radiation
$a = 11.4644(11) \text{ \AA}$	$\mu = 0.22 \text{ mm}^{-1}$
$b = 12.0231(12) \text{ \AA}$	$T = 298 \text{ K}$
$c = 15.1184(15) \text{ \AA}$	$0.40 \times 0.30 \times 0.20 \text{ mm}$
$\beta = 101.731(2)^\circ$	

### Data collection

Bruker SMART CCD area-detector diffractometer	4007 independent reflections
21013 measured reflections	3573 reflections with $I > 2\sigma(I)$
	$R_{\text{int}} = 0.059$

### Refinement

$R[F^2 > 2\sigma(F^2)] = 0.069$	265 parameters
$wR(F^2) = 0.164$	H-atom parameters constrained
$S = 1.17$	$\Delta\rho_{\text{max}} = 0.32 \text{ e \AA}^{-3}$
4007 reflections	$\Delta\rho_{\text{min}} = -0.22 \text{ e \AA}^{-3}$

Table 1

Hydrogen-bond geometry ( $\text{\AA}$ ,  $^\circ$ ).

$D-\text{H}\cdots A$	$D-\text{H}$	$\text{H}\cdots A$	$D\cdots A$	$D-\text{H}\cdots A$
$\text{C3}-\text{H3}\cdots\text{O3}^i$	0.93	2.49	3.168 (3)	129
$\text{C10}-\text{H10A}\cdots\text{O5}^{\text{ii}}$	0.96	2.51	3.235 (4)	132
$\text{C15}-\text{H15B}\cdots\text{O2}$	0.96	2.59	3.312 (4)	132

Symmetry codes: (i)  $x + \frac{1}{2}, -y + \frac{1}{2}, z + \frac{1}{2}$ ; (ii)  $-x + \frac{1}{2}, y + \frac{1}{2}, -z + \frac{3}{2}$ .

Data collection: *SMART* (Bruker, 1997); cell refinement: *SAINTE* (Bruker, 1999); data reduction: *SAINTE*; program(s) used to solve structure: *SHELXS97* (Sheldrick, 2008); program(s) used to refine structure: *SHELXL97* (Sheldrick, 2008); molecular graphics: *SHELXTL* (Sheldrick, 2008); software used to prepare material for publication: *SHELXTL*.

The authors are grateful to the Central China Normal University for financial support and thank Dr Xiang-Gao Meng for the X-ray data collection.

Supplementary data and figures for this paper are available from the IUCr electronic archives (Reference: NC2228).

## References

- Bruker (1997). *SMART*. Bruker AXS Inc., Madison, Wisconsin, USA.  
 Bruker (1999). *SAINTE*. Bruker AXS Inc., Madison, Wisconsin, USA.  
 Gao, M., Yang, Y., Wu, Y.-D., Deng, C., Shu, W.-M., Zhang, D.-X., Cao, L.-P., She, N.-F. & Wu, A.-X. (2010). *Org. Lett.* **12**, 4026–4029.  
 Sheldrick, G. M. (2008). *Acta Cryst.* **A64**, 112–122.

**supplementary materials**

*Acta Cryst.* (2011). E67, o1508 [ doi:10.1107/S1600536811018733 ]

**Methyl 3-(4-chlorophenyl)-2-(1,3-dimethyl-2,5-dioxo-4-phenylimidazolidin-4-yl)-3-oxopropanoate**

**D. Zhang, C. Deng and Y. Yang**

**Experimental**

The title compound was synthesized according to the reported literature (Gao *et al.*, 2010). The crystal was grown by slow evaporation of the solvent at room temperature from a chloroform-methanol(1:1) solution of the title compound.

**Refinement**

All H atoms were positioned in geometrically idealized positions and constrained to ride on their parent atoms, with C—H distances in the range of 0.93–0.98 Å and  $U_{\text{iso}}(\text{H}) = 1.2U_{\text{eq}}(\text{C})$  (1.5 for methyl H atoms).

**Figures**




Fig. 1. A view of (I), showing the atom-labelling scheme, with displacement ellipsoids drawn at the 30% probability level. H atoms omitted for clarity.

**Methyl 3-(4-chlorophenyl)-2-(1,3-dimethyl-2,5-dioxo-4-phenylimidazolidin-4-yl)-3-oxopropanoate**

*Crystal data*

$\text{C}_{21}\text{H}_{19}\text{ClN}_2\text{O}_5$

$M_r = 414.83$

Monoclinic,  $P2_1/n$

Hall symbol: -P 2yn

$a = 11.4644$  (11) Å

$b = 12.0231$  (12) Å

$c = 15.1184$  (15) Å

$\beta = 101.731$  (2)°

$V = 2040.4$  (3) Å<sup>3</sup>

$Z = 4$

$F(000) = 864$

$D_x = 1.350$  Mg m<sup>-3</sup>

Mo  $K\alpha$  radiation,  $\lambda = 0.71073$  Å

Cell parameters from 6439 reflections

$\theta = 2.2$ – $27.8^\circ$

$\mu = 0.22$  mm<sup>-1</sup>

$T = 298$  K

Block, colourless

$0.40 \times 0.30 \times 0.20$  mm

*Data collection*

Bruker SMART CCD area-detector diffractometer

Radiation source: fine-focus sealed tube graphite

3573 reflections with  $I > 2\sigma(I)$

$R_{\text{int}} = 0.059$

$\theta_{\text{max}} = 26.0^\circ$ ,  $\theta_{\text{min}} = 2.0^\circ$

# supplementary materials

---

$\varphi$ and $\omega$ scans	$h = -14 \rightarrow 14$
21013 measured reflections	$k = -14 \rightarrow 14$
4007 independent reflections	$l = -18 \rightarrow 18$

## Refinement

Refinement on $F^2$	Primary atom site location: structure-invariant direct methods
Least-squares matrix: full	Secondary atom site location: difference Fourier map
$R[F^2 > 2\sigma(F^2)] = 0.069$	Hydrogen site location: inferred from neighbouring sites
$wR(F^2) = 0.164$	H-atom parameters constrained
$S = 1.17$	$w = 1/[\sigma^2(F_o^2) + (0.0623P)^2 + 1.2007P]$
4007 reflections	where $P = (F_o^2 + 2F_c^2)/3$
265 parameters	$(\Delta/\sigma)_{\max} = 0.001$
0 restraints	$\Delta\rho_{\max} = 0.32 \text{ e } \text{\AA}^{-3}$
	$\Delta\rho_{\min} = -0.22 \text{ e } \text{\AA}^{-3}$

## Special details

**Geometry.** All e.s.d.'s (except the e.s.d. in the dihedral angle between two l.s. planes) are estimated using the full covariance matrix. The cell e.s.d.'s are taken into account individually in the estimation of e.s.d.'s in distances, angles and torsion angles; correlations between e.s.d.'s in cell parameters are only used when they are defined by crystal symmetry. An approximate (isotropic) treatment of cell e.s.d.'s is used for estimating e.s.d.'s involving l.s. planes.

**Refinement.** Refinement of  $F^2$  against ALL reflections. The weighted  $R$ -factor  $wR$  and goodness of fit  $S$  are based on  $F^2$ , conventional  $R$ -factors  $R$  are based on  $F$ , with  $F$  set to zero for negative  $F^2$ . The threshold expression of  $F^2 > \sigma(F^2)$  is used only for calculating  $R$ -factors(gt) *etc.* and is not relevant to the choice of reflections for refinement.  $R$ -factors based on  $F^2$  are statistically about twice as large as those based on  $F$ , and  $R$ -factors based on ALL data will be even larger.

## Fractional atomic coordinates and isotropic or equivalent isotropic displacement parameters ( $\text{\AA}^2$ )

	$x$	$y$	$z$	$U_{\text{iso}}^*/U_{\text{eq}}$
C1	0.6491 (2)	0.2716 (2)	0.94021 (16)	0.0355 (5)
C2	0.6948 (3)	0.3041 (2)	1.02851 (18)	0.0487 (7)
H2	0.7238	0.2505	1.0718	0.058*
C3	0.6979 (3)	0.4144 (3)	1.05292 (19)	0.0551 (8)
H3	0.7286	0.4356	1.1122	0.066*
C4	0.6548 (2)	0.4928 (2)	0.9885 (2)	0.0482 (7)
C5	0.6083 (3)	0.4641 (2)	0.90104 (19)	0.0493 (7)
H5	0.5786	0.5183	0.8585	0.059*
C6	0.6058 (2)	0.3535 (2)	0.87671 (17)	0.0443 (6)
H6	0.5750	0.3333	0.8172	0.053*
C7	0.6478 (2)	0.1512 (2)	0.91874 (15)	0.0350 (5)
C8	0.6010 (2)	0.1162 (2)	0.81994 (15)	0.0344 (5)
H8	0.6337	0.1691	0.7821	0.041*
C9	0.4663 (2)	0.1244 (2)	0.79297 (18)	0.0437 (6)
C10	0.2854 (3)	0.1228 (4)	0.8447 (3)	0.0785 (11)

H10A	0.2548	0.1872	0.8102	0.118*
H10B	0.2578	0.1223	0.9006	0.118*
H10C	0.2582	0.0569	0.8109	0.118*
C11	0.6383 (2)	-0.0019 (2)	0.79438 (15)	0.0347 (5)
C12	0.6180 (2)	-0.0061 (2)	0.69012 (16)	0.0378 (6)
C13	0.5134 (2)	-0.1486 (2)	0.7368 (2)	0.0483 (7)
C14	0.5159 (3)	-0.1377 (3)	0.5715 (2)	0.0702 (10)
H14A	0.4433	-0.1012	0.5435	0.105*
H14B	0.5033	-0.2166	0.5712	0.105*
H14C	0.5771	-0.1207	0.5386	0.105*
C15	0.5383 (3)	-0.1235 (3)	0.9003 (2)	0.0591 (8)
H15A	0.5019	-0.1958	0.8957	0.089*
H15B	0.4867	-0.0708	0.9207	0.089*
H15C	0.6130	-0.1263	0.9426	0.089*
C16	0.7700 (2)	-0.0284 (2)	0.82963 (15)	0.0352 (5)
C17	0.8069 (3)	-0.1364 (2)	0.85344 (17)	0.0442 (6)
H17	0.7505	-0.1925	0.8509	0.053*
C18	0.9261 (3)	-0.1613 (3)	0.88070 (19)	0.0548 (7)
H18	0.9493	-0.2339	0.8966	0.066*
C19	1.0107 (3)	-0.0798 (3)	0.8845 (2)	0.0615 (8)
H19	1.0910	-0.0964	0.9040	0.074*
C20	0.9752 (3)	0.0270 (3)	0.8592 (2)	0.0607 (8)
H20	1.0322	0.0823	0.8606	0.073*
C21	0.8562 (2)	0.0526 (2)	0.83176 (19)	0.0457 (6)
H21	0.8336	0.1250	0.8146	0.055*
Cl1	0.66072 (9)	0.63209 (7)	1.01981 (7)	0.0773 (3)
N1	0.5525 (2)	-0.0993 (2)	0.66416 (14)	0.0480 (6)
N2	0.55847 (19)	-0.08964 (18)	0.81188 (14)	0.0425 (5)
O1	0.68288 (17)	0.08220 (15)	0.97634 (11)	0.0474 (5)
O2	0.41472 (16)	0.12565 (17)	0.86363 (13)	0.0524 (5)
O3	0.41480 (19)	0.1265 (2)	0.71633 (14)	0.0767 (7)
O4	0.65545 (17)	0.05911 (16)	0.64252 (12)	0.0486 (5)
O5	0.4498 (2)	-0.22967 (19)	0.73042 (17)	0.0748 (7)

Atomic displacement parameters ( $\text{\AA}^2$ )

	$U^{11}$	$U^{22}$	$U^{33}$	$U^{12}$	$U^{13}$	$U^{23}$
C1	0.0306 (12)	0.0426 (13)	0.0341 (12)	-0.0007 (10)	0.0082 (9)	-0.0021 (10)
C2	0.0519 (16)	0.0500 (16)	0.0380 (14)	0.0056 (13)	-0.0052 (12)	-0.0034 (12)
C3	0.0582 (18)	0.0561 (17)	0.0433 (15)	0.0039 (14)	-0.0077 (13)	-0.0166 (13)
C4	0.0420 (14)	0.0442 (15)	0.0596 (17)	-0.0026 (12)	0.0135 (13)	-0.0134 (13)
C5	0.0577 (17)	0.0422 (15)	0.0504 (16)	0.0018 (13)	0.0165 (13)	0.0041 (12)
C6	0.0520 (15)	0.0484 (15)	0.0325 (13)	-0.0016 (12)	0.0086 (11)	-0.0028 (11)
C7	0.0290 (12)	0.0449 (13)	0.0312 (12)	0.0007 (10)	0.0064 (9)	-0.0012 (10)
C8	0.0341 (12)	0.0382 (13)	0.0310 (12)	-0.0004 (10)	0.0065 (9)	-0.0027 (10)
C9	0.0386 (14)	0.0498 (15)	0.0397 (14)	0.0030 (11)	0.0005 (11)	-0.0070 (12)
C10	0.0333 (16)	0.119 (3)	0.082 (2)	0.0004 (18)	0.0098 (15)	-0.023 (2)
C11	0.0357 (12)	0.0376 (13)	0.0316 (12)	-0.0044 (10)	0.0087 (10)	-0.0019 (10)

## supplementary materials

---

C12	0.0358 (13)	0.0432 (14)	0.0343 (13)	0.0058 (11)	0.0066 (10)	-0.0043 (11)
C13	0.0426 (15)	0.0441 (15)	0.0574 (17)	-0.0059 (12)	0.0084 (12)	-0.0077 (13)
C14	0.084 (2)	0.073 (2)	0.0471 (18)	-0.0079 (18)	-0.0021 (16)	-0.0221 (16)
C15	0.072 (2)	0.0542 (17)	0.0597 (19)	-0.0116 (15)	0.0324 (16)	0.0037 (14)
C16	0.0392 (13)	0.0381 (13)	0.0286 (11)	0.0018 (10)	0.0080 (10)	-0.0029 (10)
C17	0.0537 (16)	0.0414 (14)	0.0381 (14)	0.0002 (12)	0.0104 (12)	0.0021 (11)
C18	0.0620 (19)	0.0526 (17)	0.0472 (16)	0.0210 (15)	0.0047 (13)	0.0053 (13)
C19	0.0415 (16)	0.074 (2)	0.0642 (19)	0.0140 (15)	-0.0004 (14)	-0.0005 (16)
C20	0.0378 (15)	0.065 (2)	0.077 (2)	-0.0027 (14)	0.0069 (14)	-0.0041 (16)
C21	0.0388 (14)	0.0416 (14)	0.0568 (16)	0.0012 (11)	0.0102 (12)	0.0016 (12)
Cl1	0.0874 (7)	0.0451 (4)	0.0994 (7)	-0.0030 (4)	0.0187 (5)	-0.0212 (4)
N1	0.0512 (13)	0.0505 (13)	0.0394 (12)	-0.0044 (11)	0.0023 (10)	-0.0118 (10)
N2	0.0428 (12)	0.0443 (12)	0.0424 (12)	-0.0103 (9)	0.0132 (9)	-0.0039 (10)
O1	0.0588 (12)	0.0461 (10)	0.0353 (10)	0.0056 (9)	0.0046 (8)	0.0028 (8)
O2	0.0337 (10)	0.0734 (14)	0.0498 (11)	0.0006 (9)	0.0076 (8)	-0.0116 (10)
O3	0.0454 (12)	0.133 (2)	0.0455 (12)	0.0053 (13)	-0.0058 (9)	-0.0075 (13)
O4	0.0559 (11)	0.0557 (11)	0.0364 (10)	0.0022 (9)	0.0147 (8)	0.0041 (8)
O5	0.0732 (15)	0.0613 (14)	0.0883 (17)	-0.0323 (12)	0.0122 (13)	-0.0147 (12)

### *Geometric parameters (Å, °)*

C1—C2	1.387 (3)	C11—C12	1.547 (3)
C1—C6	1.394 (4)	C12—O4	1.201 (3)
C1—C7	1.483 (3)	C12—N1	1.361 (3)
C2—C3	1.375 (4)	C13—O5	1.209 (3)
C2—H2	0.9300	C13—N2	1.348 (3)
C3—C4	1.373 (4)	C13—N1	1.400 (4)
C3—H3	0.9300	C14—N1	1.453 (3)
C4—C5	1.366 (4)	C14—H14A	0.9600
C4—Cl1	1.738 (3)	C14—H14B	0.9600
C5—C6	1.378 (4)	C14—H14C	0.9600
C5—H5	0.9300	C15—N2	1.460 (3)
C6—H6	0.9300	C15—H15A	0.9600
C7—O1	1.211 (3)	C15—H15B	0.9600
C7—C8	1.540 (3)	C15—H15C	0.9600
C8—C9	1.518 (3)	C16—C21	1.384 (4)
C8—C11	1.554 (3)	C16—C17	1.390 (3)
C8—H8	0.9800	C17—C18	1.377 (4)
C9—O3	1.189 (3)	C17—H17	0.9300
C9—O2	1.322 (3)	C18—C19	1.372 (5)
C10—O2	1.452 (3)	C18—H18	0.9300
C10—H10A	0.9600	C19—C20	1.377 (5)
C10—H10B	0.9600	C19—H19	0.9300
C10—H10C	0.9600	C20—C21	1.378 (4)
C11—N2	1.456 (3)	C20—H20	0.9300
C11—C16	1.529 (3)	C21—H21	0.9300
C2—C1—C6	118.4 (2)	O4—C12—C11	126.2 (2)
C2—C1—C7	118.1 (2)	N1—C12—C11	106.3 (2)
C6—C1—C7	123.5 (2)	O5—C13—N2	127.8 (3)

C3—C2—C1	120.9 (3)	O5—C13—N1	124.2 (3)
C3—C2—H2	119.5	N2—C13—N1	108.0 (2)
C1—C2—H2	119.5	N1—C14—H14A	109.5
C4—C3—C2	119.1 (3)	N1—C14—H14B	109.5
C4—C3—H3	120.5	H14A—C14—H14B	109.5
C2—C3—H3	120.5	N1—C14—H14C	109.5
C5—C4—C3	121.8 (3)	H14A—C14—H14C	109.5
C5—C4—C11	119.5 (2)	H14B—C14—H14C	109.5
C3—C4—C11	118.7 (2)	N2—C15—H15A	109.5
C4—C5—C6	119.0 (3)	N2—C15—H15B	109.5
C4—C5—H5	120.5	H15A—C15—H15B	109.5
C6—C5—H5	120.5	N2—C15—H15C	109.5
C5—C6—C1	120.8 (2)	H15A—C15—H15C	109.5
C5—C6—H6	119.6	H15B—C15—H15C	109.5
C1—C6—H6	119.6	C21—C16—C17	118.3 (2)
O1—C7—C1	121.6 (2)	C21—C16—C11	120.7 (2)
O1—C7—C8	120.6 (2)	C17—C16—C11	120.8 (2)
C1—C7—C8	117.7 (2)	C18—C17—C16	120.8 (3)
C9—C8—C7	112.17 (19)	C18—C17—H17	119.6
C9—C8—C11	107.97 (19)	C16—C17—H17	119.6
C7—C8—C11	115.58 (19)	C19—C18—C17	120.5 (3)
C9—C8—H8	106.9	C19—C18—H18	119.8
C7—C8—H8	106.9	C17—C18—H18	119.8
C11—C8—H8	106.9	C18—C19—C20	119.2 (3)
O3—C9—O2	124.9 (3)	C18—C19—H19	120.4
O3—C9—C8	122.7 (2)	C20—C19—H19	120.4
O2—C9—C8	112.4 (2)	C19—C20—C21	120.7 (3)
O2—C10—H10A	109.5	C19—C20—H20	119.6
O2—C10—H10B	109.5	C21—C20—H20	119.6
H10A—C10—H10B	109.5	C20—C21—C16	120.5 (3)
O2—C10—H10C	109.5	C20—C21—H21	119.7
H10A—C10—H10C	109.5	C16—C21—H21	119.7
H10B—C10—H10C	109.5	C12—N1—C13	111.6 (2)
N2—C11—C16	113.5 (2)	C12—N1—C14	125.0 (3)
N2—C11—C12	101.05 (18)	C13—N1—C14	123.1 (2)
C16—C11—C12	106.35 (18)	C13—N2—C11	112.1 (2)
N2—C11—C8	113.72 (19)	C13—N2—C15	121.2 (2)
C16—C11—C8	113.95 (19)	C11—N2—C15	126.2 (2)
C12—C11—C8	106.94 (19)	C9—O2—C10	116.6 (2)
O4—C12—N1	127.4 (2)		
C6—C1—C2—C3	-0.2 (4)	N2—C11—C16—C21	-169.5 (2)
C7—C1—C2—C3	-179.6 (3)	C12—C11—C16—C21	80.4 (3)
C1—C2—C3—C4	0.0 (5)	C8—C11—C16—C21	-37.2 (3)
C2—C3—C4—C5	0.5 (5)	N2—C11—C16—C17	15.7 (3)
C2—C3—C4—C11	-179.2 (2)	C12—C11—C16—C17	-94.4 (3)
C3—C4—C5—C6	-0.8 (4)	C8—C11—C16—C17	148.0 (2)
C11—C4—C5—C6	179.0 (2)	C21—C16—C17—C18	1.7 (4)
C4—C5—C6—C1	0.6 (4)	C11—C16—C17—C18	176.6 (2)
C2—C1—C6—C5	-0.1 (4)	C16—C17—C18—C19	-0.2 (4)

## supplementary materials

C7—C1—C6—C5	179.2 (2)	C17—C18—C19—C20	-1.2 (5)
C2—C1—C7—O1	1.6 (4)	C18—C19—C20—C21	1.1 (5)
C6—C1—C7—O1	-177.7 (2)	C19—C20—C21—C16	0.3 (5)
C2—C1—C7—C8	-177.9 (2)	C17—C16—C21—C20	-1.7 (4)
C6—C1—C7—C8	2.8 (3)	C11—C16—C21—C20	-176.6 (3)
O1—C7—C8—C9	105.8 (3)	O4—C12—N1—C13	-173.5 (3)
C1—C7—C8—C9	-74.8 (3)	C11—C12—N1—C13	8.1 (3)
O1—C7—C8—C11	-18.6 (3)	O4—C12—N1—C14	0.5 (4)
C1—C7—C8—C11	160.9 (2)	C11—C12—N1—C14	-178.0 (3)
C7—C8—C9—O3	162.1 (3)	O5—C13—N1—C12	176.9 (3)
C11—C8—C9—O3	-69.4 (3)	N2—C13—N1—C12	-2.7 (3)
C7—C8—C9—O2	-19.6 (3)	O5—C13—N1—C14	2.9 (5)
C11—C8—C9—O2	108.9 (2)	N2—C13—N1—C14	-176.7 (3)
C9—C8—C11—N2	-39.6 (3)	O5—C13—N2—C11	176.0 (3)
C7—C8—C11—N2	86.9 (2)	N1—C13—N2—C11	-4.4 (3)
C9—C8—C11—C16	-171.7 (2)	O5—C13—N2—C15	3.3 (5)
C7—C8—C11—C16	-45.2 (3)	N1—C13—N2—C15	-177.1 (2)
C9—C8—C11—C12	71.1 (2)	C16—C11—N2—C13	-104.7 (2)
C7—C8—C11—C12	-162.42 (19)	C12—C11—N2—C13	8.7 (3)
N2—C11—C12—O4	171.7 (2)	C8—C11—N2—C13	122.9 (2)
C16—C11—C12—O4	-69.6 (3)	C16—C11—N2—C15	67.5 (3)
C8—C11—C12—O4	52.5 (3)	C12—C11—N2—C15	-179.1 (2)
N2—C11—C12—N1	-9.8 (2)	C8—C11—N2—C15	-64.9 (3)
C16—C11—C12—N1	108.9 (2)	O3—C9—O2—C10	3.8 (4)
C8—C11—C12—N1	-129.0 (2)	C8—C9—O2—C10	-174.4 (3)

### Hydrogen-bond geometry ( $\text{\AA}$ , $^\circ$ )

$D-H\cdots A$	$D-H$	$H\cdots A$	$D\cdots A$	$D-H\cdots A$
C3—H3 $\cdots$ O3 <sup>i</sup>	0.93	2.49	3.168 (3)	129
C10—H10A $\cdots$ O5 <sup>ii</sup>	0.96	2.51	3.235 (4)	132
C15—H15B $\cdots$ O2	0.96	2.59	3.312 (4)	132

Symmetry codes: (i)  $x+1/2, -y+1/2, z+1/2$ ; (ii)  $-x+1/2, y+1/2, -z+3/2$ .



Fig. 1

