

1,5-Bis(1-phenylethylidene)thiocarbonylhydrazide

Lei Feng,^{a*} Haiwei Ji,^b Renliang Wang,^b Haiyan Ge^b and Li Li^b

^aAtherosclerosis Institute, Taishan Medical University, 271000 Taian, People's Republic of China, and ^bCollege of Chemistry and Chemical Engineering, Taishan Medical University, 271016 Taian, People's Republic of China

Correspondence e-mail: jzpap@yahoo.com.cn


Received 23 April 2011; accepted 14 May 2011

Key indicators: single-crystal X-ray study; $T = 293$ K; mean $\sigma(\text{C}-\text{C}) = 0.004$ Å; R factor = 0.047; wR factor = 0.146; data-to-parameter ratio = 14.2.

The title molecule, $\text{C}_{17}\text{H}_{18}\text{N}_4\text{S}$, is not planar, as indicated by the dihedral angle of 27.24 (9)° between the two benzene rings. In the crystal, intermolecular $\text{N}-\text{H}\cdots\text{S}$ hydrogen bonds link pairs of molecules into inversion dimers.

Related literature

For the biological activity and catalytic abilities of Schiff base derivatives and complexes, see: Loncle *et al.* (2004); Camp *et al.* (2010). For a related structure, see: Meyers *et al.* (1995).



Experimental

Crystal data

$\text{C}_{17}\text{H}_{18}\text{N}_4\text{S}$
 $M_r = 310.41$
 Triclinic, $P\bar{1}$
 $a = 7.5947$ (15) Å
 $b = 8.8202$ (18) Å

$c = 13.084$ (3) Å
 $\alpha = 76.62$ (3)°
 $\beta = 76.39$ (3)°
 $\gamma = 82.35$ (3)°
 $V = 825.8$ (3) Å³

$Z = 2$
 Mo $K\alpha$ radiation
 $\mu = 0.20$ mm⁻¹

$T = 293$ K
 $0.20 \times 0.16 \times 0.12$ mm

Data collection

Bruker SMART APEX CCD area-detector diffractometer
 Absorption correction: multi-scan (SADABS; Sheldrick, 1996)
 $T_{\min} = 0.962$, $T_{\max} = 0.977$

4272 measured reflections
 2876 independent reflections
 2033 reflections with $I > 2\sigma(I)$
 $R_{\text{int}} = 0.022$

Refinement

$R[F^2 > 2\sigma(F^2)] = 0.047$
 $wR(F^2) = 0.146$
 $S = 1.02$
 2876 reflections
 202 parameters

1 restraint
 H-atom parameters constrained
 $\Delta\rho_{\text{max}} = 0.24$ e Å⁻³
 $\Delta\rho_{\text{min}} = -0.19$ e Å⁻³

Table 1

Hydrogen-bond geometry (Å, °).

$D-H\cdots A$	$D-H$	$H\cdots A$	$D\cdots A$	$D-H\cdots A$
$\text{N3}-\text{H3}\cdots\text{S1}^i$	0.86	2.67	3.515 (2)	169

Symmetry code: (i) $-x + 2, -y + 1, -z$.

Data collection: SMART (Bruker, 2007); cell refinement: SAINT (Bruker, 2007); data reduction: SAINT; program(s) used to solve structure: SHELXS97 (Sheldrick, 2008); program(s) used to refine structure: SHELXL97 (Sheldrick, 2008); molecular graphics: SHELXTL (Sheldrick, 2008); software used to prepare material for publication: SHELXTL.

The authors acknowledge financial support by Taishan Medical University.

Supplementary data and figures for this paper are available from the IUCr electronic archives (Reference: VM2088).

References

- Bruker (2007). SMART and SAINT. Bruker AXS Inc., Madison, Wisconsin, USA.
 Camp, C., Mougél, V., Horeglad, P., Pcaut, J. & Mazzanti, M. (2010). *J. Am. Chem. Soc.* **132**, 17374–17377.
 Loncle, C., Brunel, J. M., Vidal, N., Dherbomez, M. & Letourneux, Y. (2004). *Eur. J. Med. Chem.* **39**, 1067–1071.
 Meyers, C. Y., Kolb, V. M. & Robinson, P. D. (1995). *Acta Cryst.* **C51**, 775–777.
 Sheldrick, G. M. (1996). SADABS. University of Göttingen, Germany.
 Sheldrick, G. M. (2008). *Acta Cryst.* **A64**, 112–122.

supplementary materials

Acta Cryst. (2011). E67, o1514 [doi:10.1107/S1600536811018320]

1,5-Bis(1-phenylethylidene)thiocarbonohydrazide

L. Feng, H. Ji, R. Wang, H. Ge and L. Li

Comment

Much interest has recently been paid to the design of schiff base derivatives and complexes, owing to their wide range of biological activities and catalytical abilities (Loncle *et al.*, 2004; Camp *et al.*, 2010). We have synthesized title compound, (I), and report here its crystal structure (Fig. 1). The bond lengths and angles are normal and correspond to those observed in bis(3-fluorophenylmethine)carbonohydrazide (Meyers *et al.*, 1995). The planes of atoms N2/N1/C1 and benzene ring C4-C9 form a dihedral angle of 2.62 (23)°, indicating that this part of the molecule is nearly planar. However, benzene rings C4-C9 and C12-C17 form a dihedral angle of 27.24 (9)°. Intermolecular N—H···S hydrogen bonds link two molecules into one dimer (Table 1, Fig. 2).

Experimental

Acetophenone (10.0 mmol) and thiocarbonylhydrazide (5.0 mmol) were mixed in 50 ml flask under solvent-free conditions. After stirring 3 h at 373 K, the resulting mixture was cooled to room temperature, and recrystallized from ethanol, and afforded the title compound as a crystalline solid.

Refinement

All H atoms were placed in geometrically idealized positions (N—H 0.86 and C—H 0.93–0.96 Å) and treated as riding on their parent atoms, with $U_{\text{iso}}(\text{H}) = 1.2\text{--}1.5U_{\text{eq}}(\text{C}, \text{N})$.

Figures




Fig. 1. View of (I) showing the atomic numbering and 30% probability displacement ellipsoids.




Fig. 2. View along *a*-axis of the packing of (I). N—H...S interactions are represented by dashed lines.

1,5-Bis(1-phenylethylidene)thiocarbonohydrazide

Crystal data

$C_{17}H_{18}N_4S$	$Z = 2$
$M_r = 310.41$	$F(000) = 328$
Triclinic, $P\bar{1}$	$D_x = 1.248 \text{ Mg m}^{-3}$
Hall symbol: $-P 1$	Mo $K\alpha$ radiation, $\lambda = 0.71073 \text{ \AA}$
$a = 7.5947 (15) \text{ \AA}$	Cell parameters from 1351 reflections
$b = 8.8202 (18) \text{ \AA}$	$\theta = 2.6\text{--}25.1^\circ$
$c = 13.084 (3) \text{ \AA}$	$\mu = 0.20 \text{ mm}^{-1}$
$\alpha = 76.62 (3)^\circ$	$T = 293 \text{ K}$
$\beta = 76.39 (3)^\circ$	Block, colourless
$\gamma = 82.35 (3)^\circ$	$0.20 \times 0.16 \times 0.12 \text{ mm}$
$V = 825.8 (3) \text{ \AA}^3$	

Data collection

Bruker SMART APEX CCD area-detector diffractometer	2876 independent reflections
Radiation source: fine-focus sealed tube graphite	2033 reflections with $I > 2\sigma(I)$
φ and ω scans	$R_{\text{int}} = 0.022$
Absorption correction: multi-scan (SADABS; Sheldrick, 1996)	$\theta_{\text{max}} = 25.0^\circ$, $\theta_{\text{min}} = 2.6^\circ$
$T_{\text{min}} = 0.962$, $T_{\text{max}} = 0.977$	$h = -8 \rightarrow 9$
4272 measured reflections	$k = -10 \rightarrow 9$
	$l = -15 \rightarrow 15$

Refinement

Refinement on F^2	Secondary atom site location: difference Fourier map
Least-squares matrix: full	Hydrogen site location: inferred from neighbouring sites
$R[F^2 > 2\sigma(F^2)] = 0.047$	H-atom parameters constrained
$wR(F^2) = 0.146$	$w = 1/[\sigma^2(F_o^2) + (0.0718P)^2 + 0.2086P]$

$S = 1.02$	where $P = (F_o^2 + 2F_c^2)/3$
2876 reflections	$(\Delta/\sigma)_{\max} = 0.001$
202 parameters	$\Delta\rho_{\max} = 0.24 \text{ e } \text{\AA}^{-3}$
1 restraint	$\Delta\rho_{\min} = -0.19 \text{ e } \text{\AA}^{-3}$
Primary atom site location: structure-invariant direct methods	Extinction correction: <i>SHELXL97</i> (Sheldrick, 2008), $F_c^* = kFc[1+0.001xFc^2\lambda^3/\sin(2\theta)]^{-1/4}$ Extinction coefficient: 0.015 (4)

Special details

Geometry. All e.s.d.'s (except the e.s.d. in the dihedral angle between two l.s. planes) are estimated using the full covariance matrix. The cell e.s.d.'s are taken into account individually in the estimation of e.s.d.'s in distances, angles and torsion angles; correlations between e.s.d.'s in cell parameters are only used when they are defined by crystal symmetry. An approximate (isotropic) treatment of cell e.s.d.'s is used for estimating e.s.d.'s involving l.s. planes.

Refinement. Refinement of F^2 against ALL reflections. The weighted R -factor wR and goodness of fit S are based on F^2 , conventional R -factors R are based on F , with F set to zero for negative F^2 . The threshold expression of $F^2 > \sigma(F^2)$ is used only for calculating R -factors(gt) *etc.* and is not relevant to the choice of reflections for refinement. R -factors based on F^2 are statistically about twice as large as those based on F , and R -factors based on ALL data will be even larger.

Fractional atomic coordinates and isotropic or equivalent isotropic displacement parameters (\AA^2)

	x	y	z	$U_{\text{iso}}^*/U_{\text{eq}}$
S1	0.91194 (11)	0.69768 (7)	-0.10817 (5)	0.0688 (3)
N1	0.7946 (3)	0.9200 (2)	0.00704 (14)	0.0560 (6)
H1	0.7779	0.9500	0.0669	0.067*
N2	0.7386 (3)	1.0205 (2)	-0.07909 (14)	0.0521 (5)
N3	0.9241 (3)	0.6971 (2)	0.09311 (14)	0.0566 (6)
H3	0.9787	0.6046	0.0982	0.068*
N4	0.8830 (3)	0.7712 (2)	0.17832 (15)	0.0520 (5)
C1	0.8745 (3)	0.7765 (3)	-0.00005 (18)	0.0524 (6)
C2	0.6582 (3)	1.1530 (3)	-0.06107 (18)	0.0498 (6)
C3	0.6205 (4)	1.2028 (3)	0.0448 (2)	0.0677 (8)
H3A	0.6098	1.1120	0.1022	0.102*
H3B	0.5091	1.2689	0.0525	0.102*
H3C	0.7185	1.2592	0.0474	0.102*
C4	0.6017 (3)	1.2612 (2)	-0.15566 (18)	0.0514 (6)
C5	0.6339 (4)	1.2169 (3)	-0.25393 (19)	0.0617 (7)
H5	0.6893	1.1180	-0.2604	0.074*
C6	0.5846 (4)	1.3179 (3)	-0.3422 (2)	0.0701 (8)
H6	0.6078	1.2867	-0.4076	0.084*
C7	0.5015 (4)	1.4645 (3)	-0.3343 (2)	0.0704 (8)
H7	0.4694	1.5324	-0.3942	0.085*
C8	0.4666 (4)	1.5095 (3)	-0.2381 (2)	0.0748 (8)
H8	0.4092	1.6080	-0.2321	0.090*
C9	0.5164 (4)	1.4089 (3)	-0.1491 (2)	0.0661 (8)
H9	0.4921	1.4411	-0.0839	0.079*

supplementary materials

C10	0.9131 (3)	0.6990 (3)	0.27089 (18)	0.0516 (6)
C11	0.9913 (4)	0.5327 (3)	0.2956 (2)	0.0708 (8)
H11A	0.9160	0.4657	0.2797	0.106*
H11B	0.9961	0.5032	0.3703	0.106*
H11C	1.1118	0.5228	0.2525	0.106*
C12	0.8641 (3)	0.7926 (3)	0.35463 (18)	0.0528 (6)
C13	0.8581 (5)	0.9550 (3)	0.3268 (2)	0.0790 (9)
H13	0.8894	1.0049	0.2550	0.095*
C14	0.8063 (7)	1.0415 (4)	0.4049 (3)	0.1186 (16)
H14	0.8012	1.1501	0.3855	0.142*
C15	0.7621 (7)	0.9703 (4)	0.5109 (3)	0.1202 (16)
H15	0.7270	1.0304	0.5631	0.144*
C16	0.7691 (5)	0.8118 (4)	0.5402 (2)	0.0960 (12)
H16	0.7384	0.7633	0.6124	0.115*
C17	0.8217 (4)	0.7239 (3)	0.4628 (2)	0.0696 (8)
H17	0.8291	0.6154	0.4834	0.084*

Atomic displacement parameters (\AA^2)

	U^{11}	U^{22}	U^{33}	U^{12}	U^{13}	U^{23}
S1	0.1111 (7)	0.0488 (4)	0.0460 (4)	0.0237 (4)	-0.0270 (4)	-0.0155 (3)
N1	0.0858 (16)	0.0416 (10)	0.0403 (10)	0.0161 (10)	-0.0229 (10)	-0.0105 (8)
N2	0.0758 (14)	0.0391 (10)	0.0407 (10)	0.0090 (9)	-0.0189 (9)	-0.0079 (8)
N3	0.0823 (15)	0.0436 (11)	0.0424 (11)	0.0156 (10)	-0.0214 (10)	-0.0092 (9)
N4	0.0683 (14)	0.0459 (11)	0.0418 (10)	0.0067 (9)	-0.0171 (9)	-0.0102 (8)
C1	0.0691 (17)	0.0430 (12)	0.0427 (13)	0.0073 (11)	-0.0156 (11)	-0.0071 (10)
C2	0.0654 (16)	0.0407 (12)	0.0446 (13)	0.0064 (11)	-0.0165 (11)	-0.0128 (10)
C3	0.097 (2)	0.0580 (15)	0.0537 (15)	0.0202 (14)	-0.0303 (15)	-0.0234 (12)
C4	0.0660 (16)	0.0410 (12)	0.0474 (13)	0.0063 (11)	-0.0176 (12)	-0.0103 (10)
C5	0.086 (2)	0.0492 (14)	0.0503 (14)	0.0137 (13)	-0.0234 (13)	-0.0130 (11)
C6	0.096 (2)	0.0658 (17)	0.0490 (14)	0.0156 (15)	-0.0264 (14)	-0.0144 (13)
C7	0.086 (2)	0.0609 (16)	0.0582 (17)	0.0102 (14)	-0.0271 (15)	0.0019 (13)
C8	0.103 (2)	0.0476 (14)	0.0700 (18)	0.0218 (15)	-0.0289 (16)	-0.0092 (13)
C9	0.094 (2)	0.0482 (14)	0.0533 (15)	0.0171 (13)	-0.0203 (14)	-0.0140 (12)
C10	0.0650 (16)	0.0453 (13)	0.0444 (13)	0.0039 (11)	-0.0194 (11)	-0.0058 (10)
C11	0.101 (2)	0.0559 (15)	0.0552 (15)	0.0207 (15)	-0.0320 (15)	-0.0116 (12)
C12	0.0649 (16)	0.0496 (13)	0.0446 (13)	0.0031 (11)	-0.0183 (11)	-0.0088 (11)
C13	0.132 (3)	0.0535 (16)	0.0543 (16)	0.0003 (16)	-0.0293 (17)	-0.0106 (13)
C14	0.230 (5)	0.0540 (18)	0.078 (2)	0.015 (2)	-0.049 (3)	-0.0248 (17)
C15	0.219 (5)	0.078 (2)	0.065 (2)	0.028 (3)	-0.036 (3)	-0.0342 (18)
C16	0.156 (3)	0.080 (2)	0.0468 (16)	0.014 (2)	-0.0207 (19)	-0.0171 (15)
C17	0.104 (2)	0.0558 (15)	0.0473 (15)	0.0050 (15)	-0.0223 (14)	-0.0075 (12)

Geometric parameters (\AA , $^\circ$)

S1—C1	1.667 (2)	C7—H7	0.9300
N1—C1	1.343 (3)	C8—C9	1.388 (3)
N1—N2	1.378 (2)	C8—H8	0.9300
N1—H1	0.8600	C9—H9	0.9300

N2—C2	1.290 (3)	C10—C12	1.471 (3)
N3—C1	1.367 (3)	C10—C11	1.499 (3)
N3—N4	1.374 (3)	C11—H11A	0.9600
N3—H3	0.8600	C11—H11B	0.9600
N4—C10	1.288 (3)	C11—H11C	0.9600
C2—C4	1.490 (3)	C12—C17	1.383 (3)
C2—C3	1.502 (3)	C12—C13	1.392 (3)
C3—H3A	0.9600	C13—C14	1.368 (4)
C3—H3B	0.9600	C13—H13	0.9300
C3—H3C	0.9600	C14—C15	1.366 (5)
C4—C5	1.387 (3)	C14—H14	0.9300
C4—C9	1.388 (3)	C15—C16	1.358 (5)
C5—C6	1.380 (3)	C15—H15	0.9300
C5—H5	0.9300	C16—C17	1.370 (4)
C6—C7	1.376 (4)	C16—H16	0.9300
C6—H6	0.9300	C17—H17	0.9300
C7—C8	1.363 (4)		
C1—N1—N2	121.73 (18)	C7—C8—H8	119.9
C1—N1—H1	119.1	C9—C8—H8	119.9
N2—N1—H1	119.1	C8—C9—C4	121.1 (2)
C2—N2—N1	115.81 (18)	C8—C9—H9	119.5
C1—N3—N4	117.21 (18)	C4—C9—H9	119.5
C1—N3—H3	121.4	N4—C10—C12	114.8 (2)
N4—N3—H3	121.4	N4—C10—C11	124.5 (2)
C10—N4—N3	120.56 (19)	C12—C10—C11	120.7 (2)
N1—C1—N3	112.62 (19)	C10—C11—H11A	109.5
N1—C1—S1	125.52 (17)	C10—C11—H11B	109.5
N3—C1—S1	121.86 (17)	H11A—C11—H11B	109.5
N2—C2—C4	114.99 (19)	C10—C11—H11C	109.5
N2—C2—C3	125.2 (2)	H11A—C11—H11C	109.5
C4—C2—C3	119.78 (19)	H11B—C11—H11C	109.5
C2—C3—H3A	109.5	C17—C12—C13	117.6 (2)
C2—C3—H3B	109.5	C17—C12—C10	121.9 (2)
H3A—C3—H3B	109.5	C13—C12—C10	120.5 (2)
C2—C3—H3C	109.5	C14—C13—C12	120.2 (3)
H3A—C3—H3C	109.5	C14—C13—H13	119.9
H3B—C3—H3C	109.5	C12—C13—H13	119.9
C5—C4—C9	117.8 (2)	C15—C14—C13	120.7 (3)
C5—C4—C2	120.7 (2)	C15—C14—H14	119.6
C9—C4—C2	121.6 (2)	C13—C14—H14	119.6
C6—C5—C4	120.8 (2)	C16—C15—C14	120.2 (3)
C6—C5—H5	119.6	C16—C15—H15	119.9
C4—C5—H5	119.6	C14—C15—H15	119.9
C7—C6—C5	120.6 (2)	C15—C16—C17	119.6 (3)
C7—C6—H6	119.7	C15—C16—H16	120.2
C5—C6—H6	119.7	C17—C16—H16	120.2
C8—C7—C6	119.5 (2)	C16—C17—C12	121.6 (3)
C8—C7—H7	120.2	C16—C17—H17	119.2
C6—C7—H7	120.2	C12—C17—H17	119.2

supplementary materials

C7—C8—C9	120.2 (2)		
C1—N1—N2—C2	177.7 (2)	C7—C8—C9—C4	-0.2 (5)
C1—N3—N4—C10	174.7 (2)	C5—C4—C9—C8	-0.6 (4)
N2—N1—C1—N3	178.3 (2)	C2—C4—C9—C8	179.1 (3)
N2—N1—C1—S1	-1.9 (4)	N3—N4—C10—C12	179.5 (2)
N4—N3—C1—N1	1.4 (3)	N3—N4—C10—C11	-0.6 (4)
N4—N3—C1—S1	-178.43 (18)	N4—C10—C12—C17	155.1 (3)
N1—N2—C2—C4	179.0 (2)	C11—C10—C12—C17	-24.8 (4)
N1—N2—C2—C3	-0.8 (4)	N4—C10—C12—C13	-24.6 (4)
N2—C2—C4—C5	0.8 (4)	C11—C10—C12—C13	155.6 (3)
C3—C2—C4—C5	-179.5 (2)	C17—C12—C13—C14	-1.9 (5)
N2—C2—C4—C9	-178.9 (3)	C10—C12—C13—C14	177.7 (3)
C3—C2—C4—C9	0.8 (4)	C12—C13—C14—C15	0.8 (6)
C9—C4—C5—C6	0.9 (4)	C13—C14—C15—C16	0.0 (7)
C2—C4—C5—C6	-178.8 (3)	C14—C15—C16—C17	0.3 (7)
C4—C5—C6—C7	-0.4 (5)	C15—C16—C17—C12	-1.5 (6)
C5—C6—C7—C8	-0.5 (5)	C13—C12—C17—C16	2.3 (5)
C6—C7—C8—C9	0.7 (5)	C10—C12—C17—C16	-177.3 (3)

Hydrogen-bond geometry (\AA , $^\circ$)

$D-H\cdots A$	$D-H$	$H\cdots A$	$D\cdots A$	$D-H\cdots A$
N3—H3 \cdots S1 ⁱ	0.86	2.67	3.515 (2)	169

Symmetry codes: (i) $-x+2, -y+1, -z$.

Fig. 1




Fig. 2

