

N-Benzyl-6-deoxy-3,6-iminomethylene-1,2,3,5-O-tetraacetyl-*a*-D-1(S)-epiallo-furanose

Qiurong Zhang, Chunli Wu, Pengyun Li, Weiyang Cheng and Hongmin Liu*

New Drug Research & Development Center, Zhengzhou University, Zhengzhou 450001, People's Republic of China
Correspondence e-mail: zqr409@163.com


Received 6 May 2011; accepted 24 May 2011

Key indicators: single-crystal X-ray study; $T = 291\text{ K}$; mean $\sigma(\text{C}-\text{C}) = 0.005\text{ \AA}$; R factor = 0.043; wR factor = 0.107; data-to-parameter ratio = 8.6.

The molecule of the title compound, $C_{22}\text{H}_{27}\text{NO}_9$, an azasugar derivative, consists of one benzene ring and two fused rings, which have the *cis* arrangement at the ring junctions, and gives a V-shaped geometry. The interplanar angle between the five- and six-membered rings is $65.69(11)^\circ$. The crystal structure is stabilized by weak intermolecular C—H···O hydrogen bonds.

Related literature

For the one-pot reaction used to obtain the title compound, see: Saito *et al.* (2002); Deshpande *et al.* (2004). For the activity of azasugars, see: Compain *et al.* (2001, 2003). For their powerful inhibitory aptitude towards carbohydrate-processing enzymes, see: Guaragna *et al.* (2009).



Experimental

Crystal data

$C_{22}\text{H}_{27}\text{NO}_9$

$M_r = 449.45$

Monoclinic, $P2_1$
 $a = 8.1768(16)\text{ \AA}$
 $b = 9.0613(18)\text{ \AA}$
 $c = 15.591(3)\text{ \AA}$
 $\beta = 94.56(3)^\circ$
 $V = 1151.5(4)\text{ \AA}^3$

$Z = 2$
Mo $K\alpha$ radiation
 $\mu = 0.10\text{ mm}^{-1}$
 $T = 291\text{ K}$
 $0.20 \times 0.18 \times 0.16\text{ mm}$

Data collection

Rigaku R-AXIS-IV diffractometer
Absorption correction: multi-scan (*SADABS*; Sheldrick, 1996)
 $T_{\min} = 0.980$, $T_{\max} = 0.984$
4341 measured reflections
2503 independent reflections
2173 reflections with $I > 2\sigma(I)$
 $R_{\text{int}} = 0.033$

Refinement

$R[F^2 > 2\sigma(F^2)] = 0.043$
 $wR(F^2) = 0.107$
 $S = 1.09$
2503 reflections
290 parameters
1 restraint
H-atom parameters constrained
 $\Delta\rho_{\max} = 0.15\text{ e \AA}^{-3}$
 $\Delta\rho_{\min} = -0.13\text{ e \AA}^{-3}$

Table 1
Hydrogen-bond geometry (\AA , $^\circ$).

$D-\text{H}\cdots A$	$D-\text{H}$	$\text{H}\cdots A$	$D\cdots A$	$D-\text{H}\cdots A$
C15—H15C···O6 ⁱ	0.96	2.33	3.161 (5)	144

Symmetry code: (i) $x - 1, y, z$.

Data collection: *R-AXIS II Software* (Rigaku, 1997); cell refinement: *R-AXIS II Software*; data reduction: *R-AXIS II Software*; program(s) used to solve structure: *SHELXS97* (Sheldrick, 2008); program(s) used to refine structure: *SHELXL97* (Sheldrick, 2008); molecular graphics: *SHELXTL* (Sheldrick, 2008); software used to prepare material for publication: *SHELXL97*.

We gratefully acknowledge the financial support of the National Natural Science Foundation of China (20572103).

Supplementary data and figures for this paper are available from the IUCr electronic archives (Reference: ZK2010).

References

- Compain, P. & Martin, O. R. (2001). *Bioorg. Med. Chem.* **9**, 3077–3092.
- Compain, P. & Martin, O. R. (2003). *Curr. Top. Med. Chem.* **3**, 541–560.
- Deshpande, S. G. & Pathak, T. (2004). *Tetrahedron Lett.* **45**, 2255–2258.
- Guaragna, A., D'Alonzo, D., Paolella, C. & Palumbo, G. (2009). *Tetrahedron Lett.* **50**, 2045–2047.
- Rigaku (1997). *R-AXIS II Software*. Rigaku Corporation, Tokyo, Japan.
- Saito, Y., Zevacob, T. A. & Agrofoglio, L. A. (2002). *Tetrahedron*, **58**, 9593–9603.
- Sheldrick, G. M. (1996). *SADABS*. University of Göttingen, Germany.
- Sheldrick, G. M. (2008). *Acta Cryst. A* **64**, 112–122.

supplementary materials

Acta Cryst. (2011). E67, o1545 [doi:10.1107/S1600536811019702]

N-Benzyl-6-deoxy-3,6-iminomethylene-1,2,3,5-O-tetraacetyl- α -D-1(S)-epiallofuranose

Q. Zhang, C. Wu, P. Li, W. Cheng and H. Liu

Comment

Owing to their powerful inhibitory aptitude towards carbohydrate processing enzymes, azasugars undoubtedly represent one of the most attractive classes of carbohydrate mimetics (Guaragna *et al.*, 2009). As a contribution to the azasugars chemistry, we report here the crystal structure of the title compound, which was obtained under one-pot reaction (Sachin *et al.*, 2004 and Saito *et al.*, 2002) of *N*-benzyl-6-deoxy-3,6-imino-methylene-1,2-*O*-isopropylidene- α -*D*-allofuranose.

In the crystal structure of the title compound (I) (Fig. 1), there are two fused rings (tetrahydrofuran ring and piperidine ring) having the *cis* arrangement at the ring junctions, giving a V-shaped molecule. The interplanar angle between the five and six membered rings is 65.69 (11) $^{\circ}$. The torsion angles O1—C1—C2—C3, O1—C1—C2—O2 and C4—O4—C1—O1 around the carbon of hemiacetal group(C1) are 109.0 (2), -136.5 (2) and -137.1 (2) $^{\circ}$ respectively, which can confirm the hemiacetal group is β configuration. The molecules are linked into a framework by means of weak C—H···O hydrogen bonds (Table 1), one of whcih occurs between CH₃ of C5-acetoxy moiety and O atoms of carbonyl of C1-acetoxy moiety, and three of which occur among CH and CH₂groups of five and six membered rings and O atoms of carbonyl of three acetoxy moieties (Fig. 2).

Experimental

N-benzyl-6-deoxy-3,6-imino-methylene-1,2-*O*-isopropylidene- α -*D*-allofuranose (2.0 g, 6.2 mmol) was dissolved in 85% acetic acid (10 ml). A solution of 15% hydrochoric acid was added to this mixture. The resulting mixture was stirred for about 6 h at ambient temperature. After the material was consumed, the reaction mixture was evaporated under reduced pressure to dryness to yield yellow solid, which was directly used without purification. Acetic anhydride (5 ml) was added to the yellow solid in dry pyridine (5 ml). The mixture was stirred for about 5 h at ambient temperature. The reaction mixture was adjusted to neutral with saturated NaHCO₃ under ice bath and filtered. The filtrate was extracted with EtOAc, dried (Na₂SO₄), and evaporated to obtain colorless oily. The oily was recrystallized from methanol to give the title compound as a white crystal. Crystals suitable for X-ray analysis were in two weeks by slow evaporation of methanol solution of the title compound at room temperature. ¹H NMR (400 MHz, CDCl₃) σ : 7.25–7.33 (5 H, m), 6.03 (1 H, s), 5.32 (1 H, s), 5.21 (1 H, m), 4.40 (1 H, d, *J*=3.2 Hz), 3.62 (2 H, dd), 3.34 (1 H, d, *J*=13.6 Hz), 2.81 (1 H, m), 2.54 (1 H, d, *J*=13.6 Hz), 2.42 (1 H, m), 2.09 (3 H, s), 2.09(3 H, s), 2.08 (3 H, s), 2.07 (3 H, s), 2.04 (3 H, s); ¹³C NMR (100 MHz, CDCl₃) σ : 170.2, 169.3, 168.9, 168.6, 137.0, 128.7, 128.3, 127.4, 99.3, 80.2, 77.4, 76.5, 68.1, 61.4, 51.4, 49.9, 21.1, 21.1, 21.0, 20.5.

Refinement

All H atoms were placed geometrically and treated as riding on their parent atoms with C—H are 0.96 Å (methylene) or 0.93 Å (aromatic), 0.82 Å (hydroxyl)and *U*_{iso}(H)=1.2*U*_{eq}(C).

supplementary materials

Figures




Fig. 1. The molecular structure of the compound, with atom labels and 30% probability displacement ellipsoids for non-H atoms.




Fig. 2. Packing of the molecules crystal structure of title compound, the molecules are connected by the C—H···O hydrogen bonds.

N-Benzyl-6-deoxy-3,6-iminomethylene-1,2,3,5-O-tetraacetyl- α -D-1(S)-epiallofuranose

Crystal data

C ₂₂ H ₂₇ NO ₉	<i>F</i> (000) = 476
<i>M_r</i> = 449.45	<i>D_x</i> = 1.296 Mg m ⁻³
Monoclinic, <i>P</i> 2 ₁	Mo <i>K</i> α radiation, λ = 0.71073 Å
Hall symbol: P 2yb	Cell parameters from 399 reflections
<i>a</i> = 8.1768 (16) Å	θ = 2–25.1°
<i>b</i> = 9.0613 (18) Å	μ = 0.10 mm ⁻¹
<i>c</i> = 15.591 (3) Å	<i>T</i> = 291 K
β = 94.56 (3)°	PRISMATIC, colorless
<i>V</i> = 1151.5 (4) Å ³	0.20 × 0.18 × 0.16 mm
<i>Z</i> = 2	

Data collection

Rigaku R-AXIS-IV diffractometer	2503 independent reflections
Radiation source: fine-focus sealed tube graphite	2173 reflections with $I > 2\sigma(I)$
Detector resolution: 0 pixels mm ⁻¹	R_{int} = 0.033
Oscillation frames scans	$\theta_{\text{max}} = 26.5^\circ$, $\theta_{\text{min}} = 1.3^\circ$
Absorption correction: multi-scan (<i>SADABS</i> : Sheldrick, 1996)	h = 0 → 10
$T_{\text{min}} = 0.980$, $T_{\text{max}} = 0.984$	k = -11 → 11
4341 measured reflections	l = -19 → 19

Refinement

Refinement on F^2	Secondary atom site location: difference Fourier map
---------------------	--

Least-squares matrix: full	Hydrogen site location: inferred from neighbouring sites
$R[F^2 > 2\sigma(F^2)] = 0.043$	H-atom parameters constrained
$wR(F^2) = 0.107$	$w = 1/[\sigma^2(F_o^2) + (0.0659P)^2]$ where $P = (F_o^2 + 2F_c^2)/3$
$S = 1.09$	$(\Delta/\sigma)_{\max} < 0.001$
2503 reflections	$\Delta\rho_{\max} = 0.15 \text{ e } \text{\AA}^{-3}$
290 parameters	$\Delta\rho_{\min} = -0.13 \text{ e } \text{\AA}^{-3}$
1 restraint	Extinction correction: <i>SHELXL</i> , $F_c^* = k F_c [1 + 0.001 x F_c^2 \lambda^3 / \sin(2\theta)]^{1/4}$
Primary atom site location: structure-invariant direct methods	Extinction coefficient: 0.040 (5)

Special details

Geometry. All e.s.d.'s (except the e.s.d. in the dihedral angle between two l.s. planes) are estimated using the full covariance matrix. The cell e.s.d.'s are taken into account individually in the estimation of e.s.d.'s in distances, angles and torsion angles; correlations between e.s.d.'s in cell parameters are only used when they are defined by crystal symmetry. An approximate (isotropic) treatment of cell e.s.d.'s is used for estimating e.s.d.'s involving l.s. planes.

Refinement. Refinement of F^2 against ALL reflections. The weighted R -factor wR and goodness of fit S are based on F^2 , conventional R -factors R are based on F , with F set to zero for negative F^2 . The threshold expression of $F^2 > \sigma(F^2)$ is used only for calculating R -factors(gt) etc. and is not relevant to the choice of reflections for refinement. R -factors based on F^2 are statistically about twice as large as those based on F , and R -factors based on ALL data will be even larger.

Fractional atomic coordinates and isotropic or equivalent isotropic displacement parameters (\AA^2)

	x	y	z	$U_{\text{iso}}^*/U_{\text{eq}}$
N1	0.1159 (3)	0.9181 (3)	0.34251 (14)	0.0450 (6)
O1	0.5766 (2)	1.1411 (2)	0.25616 (13)	0.0515 (5)
O2	0.5092 (2)	0.8192 (2)	0.13055 (12)	0.0471 (5)
O3	0.2300 (2)	0.7356 (2)	0.19908 (12)	0.0439 (5)
O4	0.3209 (2)	1.1154 (2)	0.18522 (13)	0.0461 (5)
O5	-0.0199 (2)	1.1524 (2)	0.15811 (13)	0.0489 (5)
O6	0.5922 (3)	1.3316 (3)	0.16484 (19)	0.0771 (7)
O7	0.7528 (3)	0.7358 (3)	0.18663 (16)	0.0700 (7)
O8	0.4156 (3)	0.5948 (3)	0.27378 (14)	0.0589 (6)
O9	-0.1360 (4)	1.0104 (3)	0.05304 (17)	0.0886 (9)
C1	0.4843 (3)	1.0630 (3)	0.18939 (19)	0.0438 (6)
H1A	0.5311	1.0805	0.1343	0.053*
C2	0.4826 (3)	0.8971 (3)	0.20843 (16)	0.0393 (6)
H2A	0.5621	0.8696	0.2561	0.047*
C3	0.3042 (3)	0.8717 (3)	0.22978 (17)	0.0376 (6)
C4	0.2149 (3)	0.9901 (3)	0.17496 (18)	0.0391 (6)
H4A	0.2062	0.9591	0.1146	0.047*
C5	0.0450 (3)	1.0225 (3)	0.20286 (18)	0.0428 (6)
H5A	-0.0268	0.9385	0.1872	0.051*
C6	0.0463 (3)	1.0483 (4)	0.29899 (19)	0.0486 (7)

supplementary materials

H6A	0.1118	1.1345	0.3153	0.058*
H6B	-0.0645	1.0647	0.3149	0.058*
C7	0.2889 (3)	0.8982 (3)	0.32508 (16)	0.0429 (6)
H7A	0.3346	0.8149	0.3579	0.052*
H7B	0.3509	0.9855	0.3433	0.052*
C8	0.6243 (3)	1.2802 (4)	0.2351 (3)	0.0569 (8)
C9	0.7193 (4)	1.3524 (5)	0.3088 (3)	0.0813 (12)
H9A	0.7512	1.4497	0.2923	0.122*
H9B	0.8155	1.2951	0.3251	0.122*
H9C	0.6525	1.3590	0.3566	0.122*
C10	0.6505 (3)	0.7422 (3)	0.1276 (2)	0.0496 (7)
C11	0.6557 (5)	0.6680 (5)	0.0432 (2)	0.0733 (10)
H11A	0.7565	0.6139	0.0420	0.110*
H11B	0.6494	0.7407	-0.0018	0.110*
H11C	0.5646	0.6013	0.0345	0.110*
C12	0.2937 (3)	0.6055 (3)	0.2257 (2)	0.0473 (7)
C13	0.1927 (5)	0.4817 (4)	0.1878 (3)	0.0770 (11)
H13A	0.2399	0.3895	0.2075	0.115*
H13B	0.1899	0.4862	0.1262	0.115*
H13C	0.0831	0.4895	0.2054	0.115*
C14	-0.1022 (4)	1.1307 (4)	0.08049 (19)	0.0501 (7)
C15	-0.1374 (4)	1.2725 (4)	0.0348 (2)	0.0664 (10)
H15A	-0.1958	1.2534	-0.0199	0.100*
H15B	-0.0361	1.3219	0.0262	0.100*
H15C	-0.2031	1.3339	0.0687	0.100*
C16	0.0969 (4)	0.9206 (5)	0.43531 (19)	0.0587 (8)
H16A	0.1255	1.0180	0.4576	0.070*
H16B	0.1727	0.8504	0.4637	0.070*
C17	-0.0747 (3)	0.8835 (4)	0.45673 (18)	0.0511 (8)
C18	-0.1365 (5)	0.9410 (6)	0.5301 (2)	0.0842 (14)
H18A	-0.0748	1.0086	0.5639	0.101*
C19	-0.2898 (5)	0.8983 (8)	0.5533 (3)	0.107 (2)
H19A	-0.3307	0.9381	0.6023	0.128*
C20	-0.3806 (4)	0.7987 (7)	0.5048 (3)	0.0886 (15)
H20A	-0.4820	0.7684	0.5215	0.106*
C21	-0.3232 (4)	0.7438 (5)	0.4322 (2)	0.0690 (10)
H21A	-0.3867	0.6778	0.3981	0.083*
C22	-0.1714 (4)	0.7852 (4)	0.4087 (2)	0.0551 (8)
H22A	-0.1332	0.7456	0.3590	0.066*

Atomic displacement parameters (\AA^2)

	U^{11}	U^{22}	U^{33}	U^{12}	U^{13}	U^{23}
N1	0.0465 (12)	0.0503 (14)	0.0377 (12)	0.0078 (11)	0.0004 (9)	0.0007 (11)
O1	0.0520 (11)	0.0460 (12)	0.0553 (12)	-0.0058 (10)	-0.0042 (8)	-0.0094 (10)
O2	0.0529 (10)	0.0469 (12)	0.0407 (10)	0.0098 (10)	-0.0010 (8)	-0.0082 (9)
O3	0.0525 (10)	0.0281 (9)	0.0493 (11)	0.0011 (8)	-0.0072 (8)	-0.0006 (9)
O4	0.0463 (10)	0.0301 (10)	0.0609 (12)	-0.0010 (8)	-0.0033 (8)	0.0031 (9)

O5	0.0541 (11)	0.0351 (10)	0.0551 (12)	0.0073 (9)	-0.0101 (9)	0.0059 (9)
O6	0.0976 (18)	0.0516 (15)	0.0848 (19)	-0.0140 (14)	0.0235 (14)	-0.0001 (15)
O7	0.0537 (11)	0.0827 (18)	0.0724 (15)	0.0187 (13)	-0.0030 (11)	-0.0070 (14)
O8	0.0694 (13)	0.0403 (12)	0.0643 (14)	0.0104 (10)	-0.0114 (11)	0.0051 (10)
O9	0.132 (2)	0.0629 (19)	0.0632 (16)	-0.0026 (16)	-0.0376 (15)	0.0019 (14)
C1	0.0451 (14)	0.0417 (16)	0.0437 (15)	-0.0028 (12)	-0.0019 (11)	-0.0016 (12)
C2	0.0458 (13)	0.0380 (15)	0.0333 (13)	0.0064 (11)	-0.0017 (10)	-0.0032 (11)
C3	0.0440 (13)	0.0284 (13)	0.0389 (14)	0.0011 (11)	-0.0059 (10)	-0.0008 (11)
C4	0.0478 (14)	0.0299 (14)	0.0382 (14)	-0.0005 (11)	-0.0048 (11)	-0.0011 (11)
C5	0.0453 (14)	0.0304 (14)	0.0505 (16)	0.0041 (12)	-0.0097 (12)	0.0031 (12)
C6	0.0478 (14)	0.0480 (18)	0.0493 (17)	0.0100 (13)	-0.0006 (12)	-0.0023 (14)
C7	0.0462 (13)	0.0450 (16)	0.0362 (13)	0.0053 (12)	-0.0052 (10)	0.0024 (13)
C8	0.0457 (15)	0.0437 (18)	0.082 (2)	-0.0031 (13)	0.0118 (15)	-0.0152 (17)
C9	0.0606 (19)	0.062 (2)	0.119 (3)	-0.0016 (18)	-0.0048 (19)	-0.040 (2)
C10	0.0489 (14)	0.0401 (15)	0.0604 (19)	0.0019 (13)	0.0086 (13)	0.0001 (15)
C11	0.084 (2)	0.070 (2)	0.068 (2)	0.012 (2)	0.0200 (17)	-0.020 (2)
C12	0.0549 (16)	0.0341 (15)	0.0523 (17)	0.0037 (13)	0.0014 (13)	0.0062 (13)
C13	0.086 (2)	0.042 (2)	0.099 (3)	-0.0069 (18)	-0.015 (2)	0.003 (2)
C14	0.0511 (15)	0.054 (2)	0.0439 (16)	0.0042 (14)	-0.0049 (12)	0.0069 (15)
C15	0.0665 (18)	0.068 (2)	0.064 (2)	0.0137 (17)	-0.0023 (15)	0.0268 (18)
C16	0.0624 (18)	0.074 (2)	0.0394 (16)	0.0000 (17)	0.0001 (13)	-0.0014 (16)
C17	0.0490 (15)	0.064 (2)	0.0404 (15)	0.0121 (15)	0.0019 (12)	0.0024 (15)
C18	0.074 (2)	0.117 (4)	0.061 (2)	0.002 (2)	0.0043 (18)	-0.034 (2)
C19	0.072 (2)	0.188 (6)	0.064 (2)	0.006 (3)	0.0209 (19)	-0.044 (3)
C20	0.0533 (17)	0.147 (5)	0.066 (2)	0.004 (3)	0.0058 (16)	-0.003 (3)
C21	0.0557 (17)	0.090 (3)	0.060 (2)	0.0025 (19)	-0.0044 (14)	0.001 (2)
C22	0.0602 (16)	0.060 (2)	0.0438 (16)	0.0102 (16)	-0.0012 (13)	-0.0006 (14)

Geometric parameters (\AA , $^\circ$)

N1—C6	1.454 (4)	C8—C9	1.487 (5)
N1—C16	1.468 (4)	C9—H9A	0.9600
N1—C7	1.472 (3)	C9—H9B	0.9600
O1—C8	1.368 (4)	C9—H9C	0.9600
O1—C1	1.424 (3)	C10—C11	1.483 (5)
O2—C10	1.353 (3)	C11—H11A	0.9600
O2—C2	1.436 (3)	C11—H11B	0.9600
O3—C12	1.341 (3)	C11—H11C	0.9600
O3—C3	1.440 (3)	C12—C13	1.487 (5)
O4—C1	1.415 (3)	C13—H13A	0.9600
O4—C4	1.429 (3)	C13—H13B	0.9600
O5—C14	1.352 (4)	C13—H13C	0.9600
O5—C5	1.447 (3)	C14—C15	1.486 (5)
O6—C8	1.200 (4)	C15—H15A	0.9600
O7—C10	1.194 (4)	C15—H15B	0.9600
O8—C12	1.203 (3)	C15—H15C	0.9600
O9—C14	1.195 (4)	C16—C17	1.506 (4)
C1—C2	1.532 (4)	C16—H16A	0.9700
C1—H1A	0.9800	C16—H16B	0.9700

supplementary materials

C2—C3	1.539 (4)	C17—C22	1.374 (4)
C2—H2A	0.9800	C17—C18	1.389 (5)
C3—C7	1.520 (4)	C18—C19	1.387 (6)
C3—C4	1.522 (4)	C18—H18A	0.9300
C4—C5	1.517 (4)	C19—C20	1.360 (7)
C4—H4A	0.9800	C19—H19A	0.9300
C5—C6	1.516 (4)	C20—C21	1.355 (6)
C5—H5A	0.9800	C20—H20A	0.9300
C6—H6A	0.9700	C21—C22	1.374 (5)
C6—H6B	0.9700	C21—H21A	0.9300
C7—H7A	0.9700	C22—H22A	0.9300
C7—H7B	0.9700		
C6—N1—C16	112.3 (2)	H9A—C9—H9B	109.5
C6—N1—C7	111.0 (2)	C8—C9—H9C	109.5
C16—N1—C7	111.2 (2)	H9A—C9—H9C	109.5
C8—O1—C1	115.3 (3)	H9B—C9—H9C	109.5
C10—O2—C2	118.0 (2)	O7—C10—O2	123.1 (3)
C12—O3—C3	120.49 (18)	O7—C10—C11	126.4 (3)
C1—O4—C4	107.5 (2)	O2—C10—C11	110.4 (3)
C14—O5—C5	116.8 (2)	C10—C11—H11A	109.5
O4—C1—O1	108.1 (2)	C10—C11—H11B	109.5
O4—C1—C2	108.4 (2)	H11A—C11—H11B	109.5
O1—C1—C2	111.0 (2)	C10—C11—H11C	109.5
O4—C1—H1A	109.8	H11A—C11—H11C	109.5
O1—C1—H1A	109.8	H11B—C11—H11C	109.5
C2—C1—H1A	109.8	O8—C12—O3	123.1 (3)
O2—C2—C1	108.3 (2)	O8—C12—C13	126.4 (3)
O2—C2—C3	108.6 (2)	O3—C12—C13	110.5 (3)
C1—C2—C3	102.2 (2)	C12—C13—H13A	109.5
O2—C2—H2A	112.4	C12—C13—H13B	109.5
C1—C2—H2A	112.4	H13A—C13—H13B	109.5
C3—C2—H2A	112.4	C12—C13—H13C	109.5
O3—C3—C7	113.3 (2)	H13A—C13—H13C	109.5
O3—C3—C4	104.26 (19)	H13B—C13—H13C	109.5
C7—C3—C4	111.4 (2)	O9—C14—O5	122.5 (3)
O3—C3—C2	116.0 (2)	O9—C14—C15	125.9 (3)
C7—C3—C2	110.0 (2)	O5—C14—C15	111.5 (3)
C4—C3—C2	101.2 (2)	C14—C15—H15A	109.5
O4—C4—C5	112.0 (2)	C14—C15—H15B	109.5
O4—C4—C3	103.7 (2)	H15A—C15—H15B	109.5
C5—C4—C3	112.6 (2)	C14—C15—H15C	109.5
O4—C4—H4A	109.4	H15A—C15—H15C	109.5
C5—C4—H4A	109.4	H15B—C15—H15C	109.5
C3—C4—H4A	109.4	N1—C16—C17	112.9 (2)
O5—C5—C6	108.9 (2)	N1—C16—H16A	109.0
O5—C5—C4	109.2 (2)	C17—C16—H16A	109.0
C6—C5—C4	112.3 (2)	N1—C16—H16B	109.0
O5—C5—H5A	108.8	C17—C16—H16B	109.0
C6—C5—H5A	108.8	H16A—C16—H16B	107.8

C4—C5—H5A	108.8	C22—C17—C18	117.5 (3)
N1—C6—C5	107.9 (2)	C22—C17—C16	122.0 (3)
N1—C6—H6A	110.1	C18—C17—C16	120.4 (3)
C5—C6—H6A	110.1	C19—C18—C17	120.4 (4)
N1—C6—H6B	110.1	C19—C18—H18A	119.8
C5—C6—H6B	110.1	C17—C18—H18A	119.8
H6A—C6—H6B	108.4	C20—C19—C18	120.3 (3)
N1—C7—C3	110.8 (2)	C20—C19—H19A	119.8
N1—C7—H7A	109.5	C18—C19—H19A	119.8
C3—C7—H7A	109.5	C21—C20—C19	119.9 (4)
N1—C7—H7B	109.5	C21—C20—H20A	120.0
C3—C7—H7B	109.5	C19—C20—H20A	120.0
H7A—C7—H7B	108.1	C20—C21—C22	120.2 (4)
O6—C8—O1	122.1 (3)	C20—C21—H21A	119.9
O6—C8—C9	126.7 (4)	C22—C21—H21A	119.9
O1—C8—C9	111.2 (4)	C17—C22—C21	121.7 (3)
C8—C9—H9A	109.5	C17—C22—H22A	119.2
C8—C9—H9B	109.5	C21—C22—H22A	119.2
C4—O4—C1—O1	-137.1 (2)	O4—C4—C5—C6	-67.2 (3)
C4—O4—C1—C2	-16.8 (3)	C3—C4—C5—C6	49.2 (3)
C8—O1—C1—O4	-78.8 (3)	C16—N1—C6—C5	-170.1 (2)
C8—O1—C1—C2	162.5 (2)	C7—N1—C6—C5	64.6 (3)
C10—O2—C2—C1	113.2 (3)	O5—C5—C6—N1	-178.5 (2)
C10—O2—C2—C3	-136.5 (2)	C4—C5—C6—N1	-57.5 (3)
O4—C1—C2—O2	105.0 (2)	C6—N1—C7—C3	-63.0 (3)
O1—C1—C2—O2	-136.5 (2)	C16—N1—C7—C3	171.1 (3)
O4—C1—C2—C3	-9.6 (3)	O3—C3—C7—N1	-65.2 (3)
O1—C1—C2—C3	109.0 (2)	C4—C3—C7—N1	51.9 (3)
C12—O3—C3—C7	-67.5 (3)	C2—C3—C7—N1	163.2 (2)
C12—O3—C3—C4	171.3 (2)	C1—O1—C8—O6	-1.0 (4)
C12—O3—C3—C2	61.0 (3)	C1—O1—C8—C9	179.9 (2)
O2—C2—C3—O3	27.8 (3)	C2—O2—C10—O7	0.3 (4)
C1—C2—C3—O3	142.1 (2)	C2—O2—C10—C11	179.3 (3)
O2—C2—C3—C7	157.9 (2)	C3—O3—C12—O8	-1.8 (4)
C1—C2—C3—C7	-87.8 (3)	C3—O3—C12—C13	177.9 (3)
O2—C2—C3—C4	-84.3 (2)	C5—O5—C14—O9	7.6 (5)
C1—C2—C3—C4	30.0 (2)	C5—O5—C14—C15	-170.2 (2)
C1—O4—C4—C5	158.4 (2)	C6—N1—C16—C17	76.4 (4)
C1—O4—C4—C3	36.7 (3)	C7—N1—C16—C17	-158.4 (3)
O3—C3—C4—O4	-161.9 (2)	N1—C16—C17—C22	33.4 (5)
C7—C3—C4—O4	75.6 (3)	N1—C16—C17—C18	-151.1 (4)
C2—C3—C4—O4	-41.2 (2)	C22—C17—C18—C19	0.5 (6)
O3—C3—C4—C5	76.7 (3)	C16—C17—C18—C19	-175.2 (4)
C7—C3—C4—C5	-45.7 (3)	C17—C18—C19—C20	0.6 (8)
C2—C3—C4—C5	-162.5 (2)	C18—C19—C20—C21	-1.8 (8)
C14—O5—C5—C6	-149.2 (2)	C19—C20—C21—C22	1.9 (7)
C14—O5—C5—C4	87.9 (3)	C18—C17—C22—C21	-0.5 (5)
O4—C4—C5—O5	53.6 (3)	C16—C17—C22—C21	175.2 (3)
C3—C4—C5—O5	170.1 (2)	C20—C21—C22—C17	-0.7 (6)


supplementary materials

Hydrogen-bond geometry (Å, °)

$D-H\cdots A$	$D-H$	$H\cdots A$	$D\cdots A$	$D-H\cdots A$
C2—H2A···O7	0.98	2.31	2.693 (3)	102
C5—H5A···O9	0.98	2.30	2.666 (4)	101
C7—H7A···O8	0.97	2.50	3.067 (4)	117
C15—H15C···O6 ⁱ	0.96	2.33	3.161 (5)	144

Symmetry codes: (i) $x-1, y, z$.

Fig. 1



supplementary materials

Fig. 2

