

metal-organic compounds

- Férey, G., Mellot-Draznieks, C., Serre, C., Millange, F., Dutour, J., Surblé, S. & Margiolaki, I. (2005). *Science*, **309**, 2040–2042.
- Humphrey, S. M., Chang, J. S., Jhung, S. H., Yoon, J. W. & Wood, P. T. (2007). *Angew. Chem. Int. Ed.* **46**, 272–275.
- Long, J. R. & Yaghi, O. M. (2009). *Chem. Soc. Rev.* **38**, 1213–1214.
- Luo, F., Batten, S. R., Che, Y. X. & Zheng, J. M. (2007). *Chem. Eur. J.* **13**, 4948–5008.
- Luo, F., Che, Y. X. & Zheng, J. M. (2008). *Inorg. Chem. Commun.* **11**, 358–362.
- Rosi, N. L., Eckert, J., Eddaoudi, M., Vodak, D. T., Kim, J., O'Keeffe, M. & Yaghi, O. M. (2003). *Science*, **300**, 1127–1129.
- Rosi, N. L., Kim, J., Eddaoudi, M., Chen, B. L., O'Keeffe, M. & Yaghi, O. M. (2005). *J. Am. Chem. Soc.* **127**, 1504–1518.
- Sheldrick, G. M. (2008). *Acta Cryst. A* **64**, 112–122.

supplementary materials

Acta Cryst. (2011). E67, m841-m842 [doi:10.1107/S1600536811020009]

Poly[[aquatrism(μ -benzene-1,4-dicarboxylato)tricobalt(II)] methanol monosolvate monohydrate]

H. Zhou, C.-X. Chu and Y.-Z. Li

Comment

Recently, research on metal-organic frameworks (MOFs) has become of increasing interest (Long *et al.*, 2009). In this context compounds based on dicarboxylate ligands are of special importance (Férey *et al.*, 2005; Rosi *et al.*, 2003). In our own investigations we tried to prepare new MOFs based on benzene-1,4-dicarboxylic acid (H_2bdc) and amines as ligands. Within this project we have reacted $CoCl_2 \cdot 6H_2O$ with H_2bdc and *N,N'*-bis(4-pyridylmethylidine)-1,4-phenylenediamine and we have obtained crystals of the title compound by accident.

The title compound is a three-dimensional framework the form **hex** type (Rosi *et al.*, 2005) built from Co cations that are linked by bdc anions. From this arrangement cavities are formed, that are filled with additional uncoordinated methanol and water molecules. Altogether there are four crystallographically independent Co cations of which two (Co2 and Co4) are located on centers of inversion. Co1 is coordinated by three oxygen atoms from three bdc ligands and one oxygen atom from one coordinated water molecule within a strongly distorted tetrahedron. Co3 is also tetrahedrally coordinated but connected only to four oxygen atoms from four bdc ligands. In contrast Co2 and Co4 are 6-fold coordinated by six oxygen atoms from bdc anions within slightly distorted octahedra. Related complexes have been reported recently (Devic *et al.*, 2005; Humphrey *et al.*, 2007; Luo *et al.*, 2007; Luo *et al.*, 2008).

Experimental

A mixture of $CoCl_2 \cdot 6H_2O$ (0.0238 g, 0.1 mmol), benzene-1,4-dicarboxylic acid (0.0166 g, 0.1 mmol), *N,N'*-bis(4-pyridylmethylidine)-1,4-phenylenediamine (0.0290 g, 0.1 mmol) combined with 5 ml mixed solvent (DMF: CH_3OH = 1:1) was stirred for 20 min at room temperature. Afterwards the solution was heated in a 25 ml Teflon-lined stainless-steel vessel at 140 °C for 72 h under autogenous pressure. Slow cooling of the resulting solution to room temperature at the rate of 10 °C.h⁻¹ afforded purple block-shaped crystals suitable for single-crystal X-ray structure analysis. Yield based on $CoCl_2 \cdot 6H_2O$: 35%.

Refinement

All non-hydrogen atoms were refined with anisotropic thermal parameters. The C—H H atoms were calculated in idealized positions with C—H = 0.95 or 0.96 Å and included in the refinement in a riding mode with U_{iso} for H assigned as 1.2 or 1.5 times U_{eq} of the attached atoms. The H atoms bound to oxygen atoms from coordinated and crystallized water molecules were located from difference maps and refined as riding, with O - H = 0.96 or 0.85 Å, and with $U_{iso}(H)$ = 1.2 $U_{eq}(O)$.

supplementary materials

Figures




Fig. 1. ORTEP diagram of the title compound. Displacement ellipsoids are drawn at 30% probability level. Hydrogen atoms have been omitted for clarity. Symmetry codes: (i) $-x, -y + 1, -z + 1$; (ii) $-x + 3, -y + 2, -z$; (iii) $-x + 1, -y + 1, -z + 1$; (iv) $x, y + 1, z - 1$; (v) $-x + 2, -y + 2, -z + 1$; (vi) $-x + 2, -y + 1, -z + 1$; (vii) $-x + 2, -y + 2, -z$.

Poly[[aquatrism(μ-benzene-1,4-dicarboxylato)tricobalt(II)] methanol monosolvate monohydrate]

Crystal data

$[Co_3(C_8H_4O_4)_3(H_2O)] \cdot CH_4O \cdot H_2O$	$Z = 2$
$M_r = 737.20$	$F(000) = 742$
Triclinic, $P\bar{1}$	$D_x = 1.652 \text{ Mg m}^{-3}$
Hall symbol: $-P\bar{1}$	Mo $K\alpha$ radiation, $\lambda = 0.71073 \text{ \AA}$
$a = 9.8456 (11) \text{ \AA}$	Cell parameters from 3571 reflections
$b = 12.0753 (15) \text{ \AA}$	$\theta = 2.2\text{--}24.1^\circ$
$c = 13.0039 (16) \text{ \AA}$	$\mu = 1.73 \text{ mm}^{-1}$
$\alpha = 91.842 (2)^\circ$	$T = 291 \text{ K}$
$\beta = 100.925 (1)^\circ$	Block, purple
$\gamma = 101.745 (1)^\circ$	$0.30 \times 0.26 \times 0.24 \text{ mm}$
$V = 1482.2 (3) \text{ \AA}^3$	

Data collection

Bruker SMART APEX CCD diffractometer	5751 independent reflections
Radiation source: sealed tube graphite	4241 reflections with $I > 2\sigma(I)$
φ and ω scans	$R_{\text{int}} = 0.031$
Absorption correction: multi-scan (<i>SADABS</i> ; Bruker, 2004)	$\theta_{\max} = 26.0^\circ, \theta_{\min} = 1.6^\circ$
$T_{\min} = 0.682, T_{\max} = 0.746$	$h = -11 \rightarrow 12$
11744 measured reflections	$k = -14 \rightarrow 14$
	$l = -15 \rightarrow 16$

Refinement

Refinement on F^2	Primary atom site location: structure-invariant direct methods
Least-squares matrix: full	Secondary atom site location: difference Fourier map
$R[F^2 > 2\sigma(F^2)] = 0.051$	Hydrogen site location: inferred from neighbouring sites
$wR(F^2) = 0.110$	H-atom parameters constrained
$S = 1.04$	$w = 1/[\sigma^2(F_o^2) + (0.05P)^2 + 1.22P]$ where $P = (F_o^2 + 2F_c^2)/3$

5751 reflections	$(\Delta/\sigma)_{\max} < 0.001$
392 parameters	$\Delta\rho_{\max} = 0.36 \text{ e Å}^{-3}$
0 restraints	$\Delta\rho_{\min} = -0.70 \text{ e Å}^{-3}$

Special details

Geometry. All e.s.d.'s (except the e.s.d. in the dihedral angle between two l.s. planes) are estimated using the full covariance matrix. The cell e.s.d.'s are taken into account individually in the estimation of e.s.d.'s in distances, angles and torsion angles; correlations between e.s.d.'s in cell parameters are only used when they are defined by crystal symmetry. An approximate (isotropic) treatment of cell e.s.d.'s is used for estimating e.s.d.'s involving l.s. planes.

Refinement. Refinement of F^2 against ALL reflections. The weighted R-factor wR and goodness of fit S are based on F^2 , conventional R-factors R are based on F , with F set to zero for negative F^2 . The threshold expression of $F^2 > \sigma(F^2)$ is used only for calculating R-factors(gt) etc. and is not relevant to the choice of reflections for refinement. R-factors based on F^2 are statistically about twice as large as those based on F , and R-factors based on ALL data will be even larger.

Fractional atomic coordinates and isotropic or equivalent isotropic displacement parameters (\AA^2)

	x	y	z	$U_{\text{iso}}^*/U_{\text{eq}}$
C1	0.7655 (4)	0.4662 (3)	0.7910 (3)	0.0345 (9)
C2	0.7907 (5)	0.3680 (4)	0.7456 (4)	0.0454 (11)
H2	0.7628	0.3508	0.6719	0.054*
C3	0.8574 (5)	0.2946 (4)	0.8090 (4)	0.0419 (10)
H3	0.8717	0.2262	0.7790	0.050*
C4	0.9021 (5)	0.3234 (3)	0.9161 (3)	0.0398 (10)
C5	0.8818 (5)	0.4247 (4)	0.9588 (4)	0.0485 (12)
H5	0.9157	0.4461	1.0315	0.058*
C6	0.8129 (5)	0.4934 (4)	0.8958 (3)	0.0426 (10)
H6	0.7980	0.5615	0.9261	0.051*
C7	0.6950 (5)	0.5449 (4)	0.7247 (4)	0.0412 (10)
C8	0.9731 (5)	0.2503 (3)	0.9811 (4)	0.0422 (11)
C9	0.7203 (4)	0.8267 (3)	0.4169 (3)	0.0301 (8)
C10	0.6962 (5)	0.7946 (4)	0.3063 (4)	0.0505 (12)
H10	0.6323	0.7266	0.2758	0.061*
C11	0.7716 (5)	0.8688 (4)	0.2457 (4)	0.0429 (10)
H11	0.7599	0.8486	0.1728	0.052*
C12	0.8616 (5)	0.9694 (3)	0.2872 (3)	0.0400 (10)
C13	0.8770 (5)	1.0004 (4)	0.3927 (4)	0.0501 (12)
H13	0.9324	1.0721	0.4222	0.060*
C14	0.8078 (5)	0.9223 (4)	0.4566 (3)	0.0421 (11)
H14	0.8257	0.9401	0.5302	0.051*
C15	0.6447 (4)	0.7483 (3)	0.4814 (3)	0.0317 (8)
C16	0.9373 (4)	1.0442 (4)	0.2203 (3)	0.0375 (9)
C17	0.2859 (4)	0.6297 (4)	0.5711 (3)	0.0371 (10)
C18	0.1060 (5)	0.4418 (4)	0.5392 (4)	0.0426 (10)
H18	0.1803	0.4042	0.5644	0.051*
C19	0.1364 (4)	0.5612 (4)	0.5314 (3)	0.0356 (9)
C20	0.0241 (4)	0.6157 (4)	0.4882 (3)	0.0388 (10)

supplementary materials

C6—H6	0.9500	C25—H25C	0.9800
C7—O1	1.095 (5)	C25—H25D	0.9800
C7—O2	1.383 (5)	Co1—O5	1.845 (3)
C8—O4	1.262 (5)	Co1—O9	2.062 (3)
C8—O3	1.305 (5)	Co1—O1	2.072 (3)
C9—C14	1.315 (6)	Co1—O1W	2.304 (3)
C9—C10	1.440 (6)	Co2—O2	1.952 (3)
C9—C15	1.468 (5)	Co2—O2 ⁱⁱⁱ	1.952 (3)
C10—C11	1.404 (6)	Co2—O6	2.024 (3)
C10—H10	0.9500	Co2—O6 ⁱⁱⁱ	2.024 (3)
C11—C12	1.376 (6)	Co2—O9	2.230 (3)
C11—H11	0.9500	Co2—O9 ⁱⁱⁱ	2.230 (3)
C12—C13	1.381 (6)	Co3—O4 ^{iv}	1.984 (3)
C12—C16	1.467 (6)	Co3—O8	1.990 (3)
C13—C14	1.431 (6)	Co3—O11	2.107 (3)
C13—H13	0.9500	Co3—O1 ^v	2.222 (3)
C14—H14	0.9500	Co4—O3 ^{iv}	2.030 (3)
C15—O6	1.289 (5)	Co4—O3 ^{vi}	2.030 (3)
C15—O5	1.437 (5)	Co4—O7	2.030 (3)
C16—O8	1.260 (5)	Co4—O7 ^{vii}	2.030 (3)
C16—O7	1.271 (5)	Co4—O11	2.201 (3)
C17—O9	1.226 (5)	Co4—O11 ^{vii}	2.201 (3)
C17—O10	1.258 (5)	O1—Co3 ^v	2.222 (3)
C17—C19	1.519 (6)	O3—Co4 ^{viii}	2.030 (3)
C18—C20 ⁱ	1.304 (6)	O4—Co3 ^{viii}	1.984 (3)
C18—C19	1.422 (6)	O13—H15B	0.9600
C18—H18	0.9500	O1W—H1X	0.9601
C19—C20	1.434 (5)	O1W—H1Y	0.9600
C20—C18 ⁱ	1.304 (6)	O2W—H2X	0.8501
C20—H20	0.9500	O2W—H2Y	0.8500
C6—C1—C2	119.5 (4)	C23—C24—H24	119.4
C6—C1—C7	119.8 (4)	O13—C25—H25A	109.5
C2—C1—C7	120.6 (4)	O13—C25—H25C	109.5
C1—C2—C3	120.0 (4)	H25A—C25—H25C	109.5
C1—C2—H2	120.0	O13—C25—H25D	109.5
C3—C2—H2	120.0	H25A—C25—H25D	109.5
C4—C3—C2	119.1 (4)	H25C—C25—H25D	109.5
C4—C3—H3	120.5	O5—Co1—O9	110.87 (13)
C2—C3—H3	120.5	O5—Co1—O1	96.16 (16)
C3—C4—C5	119.8 (4)	O9—Co1—O1	99.86 (11)
C3—C4—C8	119.5 (4)	O5—Co1—O1W	140.91 (13)
C5—C4—C8	120.6 (4)	O9—Co1—O1W	93.65 (11)
C6—C5—C4	120.0 (4)	O1—Co1—O1W	109.49 (14)
C6—C5—H5	120.0	O2—Co2—O2 ⁱⁱⁱ	180.000 (1)
C4—C5—H5	120.0	O2—Co2—O6	96.10 (12)
C1—C6—C5	121.5 (4)	O2 ⁱⁱⁱ —Co2—O6	83.90 (12)

supplementary materials

C1—C6—H6	119.3	O2—Co2—O6 ⁱⁱⁱ	83.90 (12)
C5—C6—H6	119.3	O2 ⁱⁱⁱ —Co2—O6 ⁱⁱⁱ	96.10 (12)
O1—C7—O2	120.2 (4)	O6—Co2—O6 ⁱⁱⁱ	180.0
O1—C7—C1	124.2 (4)	O2—Co2—O9	90.15 (11)
O2—C7—C1	115.2 (4)	O2 ⁱⁱⁱ —Co2—O9	89.85 (11)
O4—C8—O3	122.1 (4)	O6—Co2—O9	91.77 (11)
O4—C8—C4	119.7 (4)	O6 ⁱⁱⁱ —Co2—O9	88.23 (11)
O3—C8—C4	118.2 (4)	O2—Co2—O9 ⁱⁱⁱ	89.85 (11)
C14—C9—C10	120.2 (4)	O2 ⁱⁱⁱ —Co2—O9 ⁱⁱⁱ	90.15 (11)
C14—C9—C15	122.6 (4)	O6—Co2—O9 ⁱⁱⁱ	88.23 (11)
C10—C9—C15	117.2 (4)	O6 ⁱⁱⁱ —Co2—O9 ⁱⁱⁱ	91.77 (11)
C11—C10—C9	116.8 (4)	O9—Co2—O9 ⁱⁱⁱ	180.000 (1)
C11—C10—H10	121.6	O4 ^{iv} —Co3—O8	106.09 (14)
C9—C10—H10	121.6	O4 ^{iv} —Co3—O11	106.51 (11)
C12—C11—C10	122.8 (4)	O8—Co3—O11	97.95 (13)
C12—C11—H11	118.6	O4 ^{iv} —Co3—O1 ^v	98.73 (11)
C10—C11—H11	118.6	O8—Co3—O1 ^v	121.70 (13)
C11—C12—C13	118.9 (4)	O11—Co3—O1 ^v	124.18 (12)
C11—C12—C16	120.6 (4)	O3 ^{iv} —Co4—O3 ^{vi}	180.00 (6)
C13—C12—C16	120.4 (4)	O3 ^{iv} —Co4—O7	93.55 (11)
C12—C13—C14	118.7 (4)	O3 ^{vi} —Co4—O7	86.45 (11)
C12—C13—H13	120.7	O3 ^{iv} —Co4—O7 ^{vii}	86.45 (11)
C14—C13—H13	120.7	O3 ^{vi} —Co4—O7 ^{vii}	93.55 (11)
C9—C14—C13	122.3 (4)	O7—Co4—O7 ^{vii}	180.000 (1)
C9—C14—H14	118.8	O3 ^{iv} —Co4—O11	92.68 (12)
C13—C14—H14	118.8	O3 ^{vi} —Co4—O11	87.32 (12)
O6—C15—O5	119.8 (3)	O7—Co4—O11	91.18 (11)
O6—C15—C9	120.9 (4)	O7 ^{vii} —Co4—O11	88.82 (11)
O5—C15—C9	119.3 (3)	O3 ^{iv} —Co4—O11 ^{vii}	87.32 (12)
O8—C16—O7	124.3 (4)	O3 ^{vi} —Co4—O11 ^{vii}	92.68 (12)
O8—C16—C12	118.2 (4)	O7—Co4—O11 ^{vii}	88.82 (11)
O7—C16—C12	117.5 (4)	O7 ^{vii} —Co4—O11 ^{vii}	91.18 (10)
O9—C17—O10	122.1 (4)	O11—Co4—O11 ^{vii}	180.0
O9—C17—C19	120.4 (4)	C7—O1—Co1	124.3 (3)
O10—C17—C19	117.4 (4)	C7—O1—Co3 ^v	132.2 (3)
C20 ⁱ —C18—C19	119.8 (4)	Co1—O1—Co3 ^v	93.67 (12)
C20 ⁱ —C18—H18	120.1	C7—O2—Co2	140.1 (3)
C19—C18—H18	120.1	C8—O3—Co4 ^{viii}	135.7 (3)
C18—C19—C20	119.3 (4)	C8—O4—Co3 ^{viii}	130.1 (3)
C18—C19—C17	119.9 (4)	C15—O5—Co1	128.3 (3)
C20—C19—C17	120.8 (4)	C15—O6—Co2	136.0 (3)
C18 ⁱ —C20—C19	120.8 (4)	C16—O7—Co4	138.1 (3)

supplementary materials

C18 ⁱ —C20—H20	119.6	C16—O8—Co3	119.8 (3)
C19—C20—H20	119.6	C17—O9—Co1	100.9 (3)
O11—C21—O12	124.7 (4)	C17—O9—Co2	126.7 (3)
O11—C21—C23	122.0 (4)	Co1—O9—Co2	102.39 (11)
O12—C21—C23	113.0 (4)	C21—O11—Co3	98.8 (3)
C24 ⁱⁱ —C22—C23	123.1 (4)	C21—O11—Co4	127.7 (3)
C24 ⁱⁱ —C22—H22	118.5	Co3—O11—Co4	101.90 (12)
C23—C22—H22	118.5	C25—O13—H15B	109.3
C22—C23—C24	115.3 (4)	Co1—O1W—H1X	109.6
C22—C23—C21	131.7 (4)	Co1—O1W—H1Y	109.4
C24—C23—C21	112.7 (4)	H1X—O1W—H1Y	109.5
C22 ⁱⁱ —C24—C23	121.1 (4)	H2X—O2W—H2Y	109.5
C22 ⁱⁱ —C24—H24	119.4		
C6—C1—C2—C3	3.6 (7)	O6 ⁱⁱⁱ —Co2—O2—C7	-82.0 (4)
C7—C1—C2—C3	179.9 (4)	O9—Co2—O2—C7	6.3 (4)
C1—C2—C3—C4	-2.4 (7)	O9 ⁱⁱⁱ —Co2—O2—C7	-173.7 (4)
C2—C3—C4—C5	-0.7 (7)	O4—C8—O3—Co4 ^{viii}	17.9 (7)
C2—C3—C4—C8	-178.6 (4)	C4—C8—O3—Co4 ^{viii}	-160.1 (3)
C3—C4—C5—C6	2.5 (7)	O3—C8—O4—Co3 ^{viii}	11.3 (7)
C8—C4—C5—C6	-179.6 (5)	C4—C8—O4—Co3 ^{viii}	-170.8 (3)
C2—C1—C6—C5	-1.8 (7)	O6—C15—O5—Co1	-11.2 (5)
C7—C1—C6—C5	-178.1 (4)	C9—C15—O5—Co1	167.2 (3)
C4—C5—C6—C1	-1.3 (8)	O9—Co1—O5—C15	-5.8 (4)
C6—C1—C7—O1	3.7 (8)	O1—Co1—O5—C15	97.2 (3)
C2—C1—C7—O1	-172.6 (5)	O1W—Co1—O5—C15	-131.1 (3)
C6—C1—C7—O2	176.1 (4)	O5—C15—O6—Co2	-22.8 (6)
C2—C1—C7—O2	-0.2 (6)	C9—C15—O6—Co2	158.9 (3)
C3—C4—C8—O4	169.3 (4)	O2—Co2—O6—C15	-33.7 (4)
C5—C4—C8—O4	-8.6 (7)	O2 ⁱⁱⁱ —Co2—O6—C15	146.3 (4)
C3—C4—C8—O3	-12.7 (7)	O9—Co2—O6—C15	56.6 (4)
C5—C4—C8—O3	169.4 (4)	O9 ⁱⁱⁱ —Co2—O6—C15	-123.4 (4)
C14—C9—C10—C11	1.0 (7)	O8—C16—O7—Co4	46.4 (7)
C15—C9—C10—C11	-178.5 (4)	C12—C16—O7—Co4	-129.9 (4)
C9—C10—C11—C12	-1.9 (7)	O3 ^{iv} —Co4—O7—C16	-106.4 (4)
C10—C11—C12—C13	-1.2 (7)	O3 ^{vi} —Co4—O7—C16	73.6 (4)
C10—C11—C12—C16	-179.7 (4)	O11—Co4—O7—C16	-13.6 (4)
C11—C12—C13—C14	5.1 (7)	O11 ^{vii} —Co4—O7—C16	166.4 (4)
C16—C12—C13—C14	-176.4 (4)	O7—C16—O8—Co3	-4.6 (6)
C10—C9—C14—C13	3.0 (7)	C12—C16—O8—Co3	171.7 (3)
C15—C9—C14—C13	-177.5 (4)	O4 ^{iv} —Co3—O8—C16	60.0 (4)
C12—C13—C14—C9	-6.1 (8)	O11—Co3—O8—C16	-49.8 (4)
C14—C9—C15—O6	-171.9 (4)	O1 ^v —Co3—O8—C16	171.3 (3)
C10—C9—C15—O6	7.7 (6)	O10—C17—O9—Co1	14.7 (5)
C14—C9—C15—O5	9.8 (6)	C19—C17—O9—Co1	-165.7 (3)
C10—C9—C15—O5	-170.7 (4)	O10—C17—O9—Co2	-99.9 (4)

C11—C12—C16—O8	−167.5 (4)	C19—C17—O9—Co2	79.7 (5)
C13—C12—C16—O8	14.1 (7)	O5—Co1—O9—C17	−90.8 (3)
C11—C12—C16—O7	9.0 (6)	O1—Co1—O9—C17	168.6 (3)
C13—C12—C16—O7	−169.4 (4)	O1W—Co1—O9—C17	58.1 (3)
C20 ⁱ —C18—C19—C20	−3.2 (7)	O5—Co1—O9—Co2	40.87 (17)
C20 ⁱ —C18—C19—C17	175.3 (4)	O1—Co1—O9—Co2	−59.66 (15)
O9—C17—C19—C18	6.7 (6)	O1W—Co1—O9—Co2	−170.18 (12)
O10—C17—C19—C18	−173.7 (4)	O2—Co2—O9—C17	158.8 (4)
O9—C17—C19—C20	−174.8 (4)	O2 ⁱⁱⁱ —Co2—O9—C17	−21.2 (4)
O10—C17—C19—C20	4.8 (6)	O6—Co2—O9—C17	62.6 (4)
C18—C19—C20—C18 ⁱ	3.2 (7)	O6 ⁱⁱⁱ —Co2—O9—C17	−117.4 (4)
C17—C19—C20—C18 ⁱ	−175.3 (4)	O2—Co2—O9—Co1	44.79 (13)
C24 ⁱⁱ —C22—C23—C24	7.7 (8)	O2 ⁱⁱⁱ —Co2—O9—Co1	−135.21 (13)
C24 ⁱⁱ —C22—C23—C21	−165.6 (5)	O6—Co2—O9—Co1	−51.32 (12)
O11—C21—C23—C22	172.5 (5)	O6 ⁱⁱⁱ —Co2—O9—Co1	128.68 (12)
O12—C21—C23—C22	−13.3 (7)	O12—C21—O11—Co3	−15.7 (5)
O11—C21—C23—C24	−0.9 (6)	C23—C21—O11—Co3	157.8 (4)
O12—C21—C23—C24	173.4 (4)	O12—C21—O11—Co4	96.9 (5)
C22—C23—C24—C22 ⁱⁱ	−7.5 (7)	C23—C21—O11—Co4	−89.6 (5)
C21—C23—C24—C22 ⁱⁱ	167.0 (4)	O4 ^{iv} —Co3—O11—C21	88.6 (3)
O2—C7—O1—Co1	27.2 (7)	O8—Co3—O11—C21	−161.9 (3)
C1—C7—O1—Co1	−160.8 (3)	O1 ^v —Co3—O11—C21	−24.5 (3)
O2—C7—O1—Co3 ^v	−109.1 (4)	O4 ^{iv} —Co3—O11—Co4	−43.08 (15)
C1—C7—O1—Co3 ^v	63.0 (7)	O8—Co3—O11—Co4	66.39 (14)
O5—Co1—O1—C7	−84.9 (5)	O1 ^v —Co3—O11—Co4	−156.19 (11)
O9—Co1—O1—C7	27.6 (5)	O3 ^{iv} —Co4—O11—C21	−58.9 (4)
O1W—Co1—O1—C7	125.1 (4)	O3 ^{vi} —Co4—O11—C21	121.1 (4)
O5—Co1—O1—Co3 ^v	64.22 (14)	O7—Co4—O11—C21	−152.6 (4)
O9—Co1—O1—Co3 ^v	176.70 (12)	O7 ^{vii} —Co4—O11—C21	27.4 (4)
O1W—Co1—O1—Co3 ^v	−85.82 (14)	O3 ^{iv} —Co4—O11—Co3	52.20 (13)
O1—C7—O2—Co2	−53.0 (7)	O3 ^{vi} —Co4—O11—Co3	−127.80 (13)
C1—C7—O2—Co2	134.3 (4)	O7—Co4—O11—Co3	−41.41 (13)
O6—Co2—O2—C7	98.0 (4)	O7 ^{vii} —Co4—O11—Co3	138.59 (13)

Symmetry codes: (i) $-x, -y+1, -z+1$; (ii) $-x+3, -y+2, -z$; (iii) $-x+1, -y+1, -z+1$; (iv) $x, y+1, z-1$; (v) $-x+2, -y+2, -z+1$; (vi) $-x+2, -y+1, -z+1$; (vii) $-x+2, -y+2, -z$; (viii) $x, y-1, z+1$.

supplementary materials

Fig. 1

