

catena-Poly[[[(1,10-phenanthroline- κ^2N,N')zinc]- μ -pyridine-2,3-dicarboxylato- $\kappa^4N,O^2:O^2',O^3$] monohydrate]

Zhen-Hua Qin,* Shu-Fang Lou, Na Wang and Fu-Yu Wang

Shangqiu Medical College, Shangqiu, Henan 476100, People's Republic of China

Correspondence e-mail: qinzhenhua1969@163.com


Received 14 April 2011; accepted 27 May 2011

 Key indicators: single-crystal X-ray study; $T = 293$ K; mean $\sigma(C-C) = 0.005$ Å; R factor = 0.051; wR factor = 0.175; data-to-parameter ratio = 15.0.

In the title complex, $\{[Zn(C_7H_3NO_4)(C_{12}H_8N_2)] \cdot H_2O\}_n$, the Zn^{II} ion is in a distorted octahedral environment, defined by two N atoms from a chelating 1,10-phenanthroline (phen) ligand and one N atom and three O atoms from two pyridine-2,3-dicarboxylate (2,3-pydc) ligands. The bridging 2,3-pydc ligands connect the Zn^{II} ions into a chain extending along [010]. O—H...O hydrogen bonds between the uncoordinated water molecules and the uncoordinated carboxylate O atoms, as well as π – π interactions between the pyridine rings of the phen ligands [centroid–centroid distance = 3.557 (2) Å], are observed.

Related literature

For complexes based on pyridine-2,3-dicarboxylic acid, see: Du *et al.* (2008); Han *et al.* (2006); Li & Li (2004); Patrick *et al.* (2003). For a description of the Cambridge Structural Database, see: Allen (2002). For a related structure, see: Shit *et al.* (2008).



Experimental

Crystal data

$[Zn(C_7H_3NO_4)(C_{12}H_8N_2)] \cdot H_2O$
 $M_r = 428.69$
 Orthorhombic, $Pbca$
 $a = 12.2099$ (8) Å
 $b = 11.6922$ (8) Å
 $c = 23.2513$ (15) Å
 $V = 3319.4$ (4) Å³
 $Z = 8$
 Mo $K\alpha$ radiation
 $\mu = 1.52$ mm⁻¹
 $T = 293$ K
 $0.23 \times 0.19 \times 0.12$ mm

Data collection

Bruker APEXII CCD diffractometer
 Absorption correction: multi-scan (SADABS; Sheldrick, 1996)
 $T_{min} = 0.721$, $T_{max} = 0.839$
 30212 measured reflections
 3802 independent reflections
 3010 reflections with $I > 2\sigma(I)$
 $R_{int} = 0.040$

Refinement

$R[F^2 > 2\sigma(F^2)] = 0.051$
 $wR(F^2) = 0.175$
 $S = 1.06$
 3802 reflections
 253 parameters
 H-atom parameters constrained
 $\Delta\rho_{max} = 0.76$ e Å⁻³
 $\Delta\rho_{min} = -1.02$ e Å⁻³

Table 1

Selected bond lengths (Å).

Zn1—O1	2.102 (2)	Zn1—N1	2.142 (3)
Zn1—O2 ⁱ	2.110 (2)	Zn1—N2	2.139 (3)
Zn1—O4 ⁱ	2.062 (2)	Zn1—N3	2.131 (3)

 Symmetry code: (i) $-x + \frac{1}{2}, y - \frac{1}{2}, z$.

Table 2

Hydrogen-bond geometry (Å, °).

$D-H \cdots A$	$D-H$	$H \cdots A$	$D \cdots A$	$D-H \cdots A$
O1W—H1WA...O3	0.85	2.22	3.022 (5)	157
O1W—H1WB...O3 ⁱⁱ	0.85	2.42	3.215 (5)	157

 Symmetry code: (ii) $-x + \frac{1}{2}, y + \frac{1}{2}, z$.

Data collection: APEX2 (Bruker, 2007); cell refinement: SAINT (Bruker, 2007); data reduction: SAINT; program(s) used to solve structure: SHELXS97 (Sheldrick, 2008); program(s) used to refine structure: SHELXL97 (Sheldrick, 2008); molecular graphics: SHELXTL (Sheldrick, 2008); software used to prepare material for publication: SHELXTL.

Supplementary data and figures for this paper are available from the IUCr electronic archives (Reference: HY2421).

References

- Allen, F. H. (2002). *Acta Cryst.* **B58**, 380–388.
 Bruker (2007). APEX2 and SAINT. Bruker AXS Inc., Madison, Wisconsin, USA.
 Du, L., Fang, R.-B. & Zhao, Q.-H. (2008). *Chin. J. Chem.* **26**, 957–961.
 Han, Z.-B., Ma, Y., Sun, Z.-G. & Yu, W.-S. (2006). *Inorg. Chem. Commun.* **9**, 844–847.
 Li, L. & Li, Y. (2004). *J. Mol. Struct.* **694**, 199–203.
 Patrick, B. O., Stevens, C. L., Storr, A. & Thompson, R. C. (2003). *Polyhedron*, **22**, 3025–3035.
 Sheldrick, G. M. (1996). SADABS. University of Göttingen, Germany.
 Sheldrick, G. M. (2008). *Acta Cryst.* **A64**, 112–122.
 Shit, S., Charkraborty, J., Howard, J. A. K., Spencer, E. C., Desplanches, C. & Mitra, S. (2008). *Struct. Chem.* **19**, 553–558.

supplementary materials

Acta Cryst. (2011). E67, m864 [doi:10.1107/S1600536811020344]

***catena*-Poly[[[(1,10-phenanthroline- κ^2N,N')zinc]- μ -pyridine-2,3-dicarboxylato- $\kappa^4N,O^2:O^2',O^3$] monohydrate]**

Z.-H. Qin, S.-F. Lou, N. Wang and F.-Y. Wang

Comment

Pyridine-carboxylic acids are widely used to form supramolecular architectures due to their diverse coordination modes such as monodentate terminal, monodentate bridging and bidentate chelating. Recently, pyridine-2,3-dicarboxylic acid (2,3- H_2pydc) acts as a bidentate and bridging ligand to construct one-dimensional chain, two-dimensional layer and three-dimensional structures (Du *et al.*, 2008; Han *et al.*, 2006; Li & Li, 2004; Patrick *et al.*, 2003). Herein, a one-dimensional polymeric complex based on 2,3- $pydc$ was investigated.

The title compound is composed of polymeric chains and uncoordinated water molecules. The asymmetric unit contains one Zn^{II} ion, one uncoordinated water molecule, one 2,3- $pydc$ ligand and one 1,10-phenanthroline (1,10-phen) ligand. The coordination geometry of the Zn^{II} ion can be described as distorted octahedral, which is defined by three N and three O atoms from one 1,10-phen and two 2,3- $pydc$ ligands, with $Zn-N$ and $Zn-O$ distances of 2.131 (3)–2.142 (3) and 2.062 (2)–2.110 (2) Å, respectively (Fig. 1, Table 1). As shown in Fig. 2, adjacent Zn^{II} ions are bridged by 2,3- $pydc$ ligands, which adopt a *syn-anti* configuration, with an internuclear $Zn\cdots Zn$ distance of 5.9206 (6) Å, building an infinite one-dimensional architecture. The 1,10-phen molecule serves as a bidentate chelating ligand to coordinate to the Zn^{II} ion. In the chain, the dihedral angle between the adjacent 1,10-phen ligands is 65.24 (3)°.

According to the search results in CSD database (Conquest version 1.12) (Allen, 2002), an isomorphous complex $\{[Mn(C_7H_3NO_4)(C_{12}H_8N_2)]\cdot H_2O\}_n$ is reported (Shit *et al.*, 2008).

Experimental

A mixture of pyridine-2,3-dicarboxylic acid (0.2 mmol), $Zn(CH_3COOH)_2\cdot 2H_2O$ (0.2 mmol), 1,10-phenanthroline (0.1 mmol) and 10 ml methanol/distilled water (v/v 9:2) sealed in a 25 ml Teflon-lined stainless steel autoclave was kept at 393 K for three days and then cooled to room temperature. Colorless crystals suitable for X-ray analysis were obtained.

Refinement

H atoms of aromatic rings were positioned geometrically and refined as riding atoms, with $C-H = 0.93$ Å and $U_{iso}(H) = 1.2U_{eq}(C)$. H atoms of water molecule were found in a difference Fourier map and refined as riding atoms, with $O-H = 0.85$ Å and $U_{iso}(H) = 1.5U_{eq}(O)$.

Figures




Fig. 1. Part of the chain structure in the title complex. Displacement ellipsoids are drawn at the 30% probability level. Uncoordinated water molecule has been omitted. [Symmetry code: (i) $1/2 - x, -1/2 + y, z.$]




Fig. 2. The packing diagram of the title complex. Dashed lines denote hydrogen bonds. H atoms not involving in hydrogen bonds have been omitted.

catena-Poly[[[(1,10-phenanthroline- κ^2N,N')zinc]- μ -pyridine-2,3-dicarboxylato- $\kappa^4N,O^2:O^2',O^3$] monohydrate]

Crystal data

$[\text{Zn}(\text{C}_7\text{H}_3\text{NO}_4)(\text{C}_{12}\text{H}_8\text{N}_2)] \cdot \text{H}_2\text{O}$

$M_r = 428.69$

Orthorhombic, *Pbca*

Hall symbol: -P 2ac 2ab

$a = 12.2099$ (8) Å

$b = 11.6922$ (8) Å

$c = 23.2513$ (15) Å

$V = 3319.4$ (4) Å³

$Z = 8$

$F(000) = 1744$

$D_x = 1.716$ Mg m⁻³

Mo $K\alpha$ radiation, $\lambda = 0.71073$ Å

Cell parameters from 4015 reflections

$\theta = 1.8$ – 28.3°

$\mu = 1.52$ mm⁻¹

$T = 293$ K

Block, colorless

$0.23 \times 0.19 \times 0.12$ mm

Data collection

Bruker APEXII CCD
diffractometer

Radiation source: fine-focus sealed tube
graphite

ϕ and ω scans

Absorption correction: multi-scan
(*SADABS*; Sheldrick, 1996)

$T_{\min} = 0.721$, $T_{\max} = 0.839$

30212 measured reflections

3802 independent reflections

3010 reflections with $I > 2\sigma(I)$

$R_{\text{int}} = 0.040$

$\theta_{\text{max}} = 27.5^\circ$, $\theta_{\text{min}} = 1.8^\circ$

$h = -15 \rightarrow 15$

$k = -15 \rightarrow 15$

$l = -29 \rightarrow 29$

Refinement

Refinement on F^2

Primary atom site location: structure-invariant direct
methods

Least-squares matrix: full

$$R[F^2 > 2\sigma(F^2)] = 0.051$$

$$wR(F^2) = 0.175$$

$$S = 1.06$$

3802 reflections

253 parameters

0 restraints

Secondary atom site location: difference Fourier map

Hydrogen site location: inferred from neighbouring sites

H-atom parameters constrained

$$w = 1/[\sigma^2(F_o^2) + (0.1103P)^2 + 4.4449P]$$

$$\text{where } P = (F_o^2 + 2F_c^2)/3$$

$$(\Delta/\sigma)_{\max} = 0.001$$

$$\Delta\rho_{\max} = 0.76 \text{ e } \text{\AA}^{-3}$$

$$\Delta\rho_{\min} = -1.02 \text{ e } \text{\AA}^{-3}$$

Fractional atomic coordinates and isotropic or equivalent isotropic displacement parameters (\AA^2)

	x	y	z	$U_{\text{iso}}^*/U_{\text{eq}}$
Zn1	0.21166 (4)	0.05837 (3)	0.378838 (18)	0.02796 (19)
O1	0.1506 (2)	0.22624 (19)	0.37472 (10)	0.0250 (5)
O2	0.19575 (19)	0.4075 (2)	0.38954 (11)	0.0248 (5)
O3	0.2708 (3)	0.4531 (3)	0.53194 (15)	0.0586 (10)
O4	0.3580 (2)	0.53247 (19)	0.45856 (10)	0.0288 (5)
N1	0.0847 (2)	-0.0270 (2)	0.33197 (12)	0.0230 (6)
N2	0.2625 (2)	0.0835 (2)	0.29172 (12)	0.0242 (6)
N3	0.3382 (2)	0.1532 (2)	0.42019 (11)	0.0217 (5)
C1	-0.0016 (3)	-0.0834 (3)	0.35158 (15)	0.0286 (7)
H1A	-0.0132	-0.0853	0.3911	0.034*
C2	-0.0768 (3)	-0.1405 (3)	0.31582 (17)	0.0345 (8)
H2A	-0.1351	-0.1809	0.3317	0.041*
C3	-0.0631 (3)	-0.1360 (3)	0.25741 (17)	0.0326 (8)
H3A	-0.1124	-0.1727	0.2332	0.039*
C4	0.0263 (3)	-0.0752 (3)	0.23440 (16)	0.0277 (7)
C5	0.0465 (3)	-0.0641 (3)	0.17420 (17)	0.0368 (9)
H5A	-0.0009	-0.0988	0.1483	0.044*
C6	0.1321 (3)	-0.0049 (4)	0.15423 (15)	0.0384 (9)
H6A	0.1427	0.0009	0.1147	0.046*
C7	0.2082 (3)	0.0498 (3)	0.19250 (16)	0.0288 (8)
C8	0.3005 (3)	0.1110 (3)	0.17423 (17)	0.0342 (8)
H8A	0.3138	0.1208	0.1352	0.041*
C9	0.3705 (3)	0.1557 (3)	0.21347 (16)	0.0349 (8)
H9A	0.4322	0.1958	0.2015	0.042*
C10	0.3491 (3)	0.1410 (3)	0.27237 (16)	0.0307 (8)
H10A	0.3973	0.1726	0.2989	0.037*
C11	0.1920 (3)	0.0382 (3)	0.25240 (15)	0.0231 (7)
C12	0.0994 (3)	-0.0230 (3)	0.27382 (14)	0.0225 (6)
C13	0.3092 (2)	0.2624 (3)	0.43099 (13)	0.0187 (6)
C14	0.3641 (3)	0.3299 (3)	0.47056 (14)	0.0219 (6)
C15	0.4553 (3)	0.2838 (3)	0.49769 (16)	0.0327 (8)
H15A	0.4938	0.3266	0.5247	0.039*
C16	0.4885 (3)	0.1736 (3)	0.48438 (18)	0.0379 (9)
H16A	0.5515	0.1428	0.5008	0.045*
C17	0.4264 (3)	0.1104 (3)	0.44629 (16)	0.0306 (8)

supplementary materials

H17A	0.4467	0.0353	0.4386	0.037*
C18	0.3260 (3)	0.4482 (3)	0.48823 (15)	0.0268 (7)
C19	0.2113 (2)	0.3025 (3)	0.39622 (14)	0.0184 (6)
O1W	0.2841 (4)	0.6974 (4)	0.57381 (19)	0.0825 (14)
H1WA	0.2747	0.6382	0.5532	0.124*
H1WB	0.2737	0.7562	0.5531	0.124*

Atomic displacement parameters (\AA^2)

	U^{11}	U^{22}	U^{33}	U^{12}	U^{13}	U^{23}
Zn1	0.0332 (3)	0.0191 (3)	0.0316 (3)	-0.00121 (15)	-0.00367 (16)	-0.00071 (14)
O1	0.0252 (12)	0.0133 (11)	0.0364 (13)	-0.0015 (9)	-0.0122 (10)	-0.0010 (9)
O2	0.0259 (12)	0.0095 (11)	0.0390 (13)	0.0011 (9)	-0.0060 (10)	-0.0012 (9)
O3	0.089 (3)	0.049 (2)	0.0371 (18)	0.0081 (17)	0.0260 (17)	-0.0036 (14)
O4	0.0411 (14)	0.0166 (11)	0.0286 (12)	-0.0053 (10)	-0.0077 (10)	-0.0003 (9)
N1	0.0236 (14)	0.0206 (13)	0.0247 (13)	-0.0018 (11)	-0.0032 (11)	0.0009 (11)
N2	0.0279 (14)	0.0189 (13)	0.0257 (14)	-0.0032 (11)	-0.0027 (11)	0.0006 (11)
N3	0.0252 (14)	0.0138 (12)	0.0260 (13)	-0.0005 (10)	-0.0049 (11)	0.0020 (10)
C1	0.0278 (17)	0.0292 (18)	0.0288 (18)	-0.0025 (14)	-0.0008 (14)	0.0031 (14)
C2	0.0253 (17)	0.0304 (19)	0.048 (2)	-0.0060 (14)	-0.0013 (16)	0.0016 (16)
C3	0.0268 (17)	0.0309 (19)	0.040 (2)	-0.0039 (15)	-0.0074 (15)	-0.0068 (15)
C4	0.0266 (17)	0.0252 (17)	0.0314 (18)	0.0019 (13)	-0.0067 (14)	-0.0047 (13)
C5	0.037 (2)	0.046 (2)	0.0278 (18)	0.0025 (17)	-0.0112 (16)	-0.0066 (15)
C6	0.043 (2)	0.051 (2)	0.0213 (17)	0.0007 (18)	-0.0068 (15)	-0.0019 (16)
C7	0.0320 (19)	0.031 (2)	0.0229 (17)	0.0047 (14)	-0.0040 (13)	0.0016 (13)
C8	0.043 (2)	0.033 (2)	0.0273 (18)	0.0008 (16)	0.0061 (15)	0.0068 (15)
C9	0.039 (2)	0.0272 (18)	0.039 (2)	-0.0074 (15)	0.0070 (16)	0.0053 (15)
C10	0.0335 (18)	0.0228 (17)	0.0358 (19)	-0.0083 (14)	-0.0003 (15)	0.0011 (14)
C11	0.0262 (16)	0.0192 (15)	0.0238 (16)	0.0014 (12)	-0.0047 (13)	0.0005 (12)
C12	0.0229 (16)	0.0193 (15)	0.0253 (16)	0.0003 (12)	-0.0051 (12)	0.0002 (12)
C13	0.0217 (14)	0.0157 (14)	0.0188 (14)	-0.0006 (11)	-0.0015 (11)	0.0016 (11)
C14	0.0279 (16)	0.0152 (14)	0.0227 (15)	-0.0050 (12)	-0.0033 (12)	0.0014 (11)
C15	0.036 (2)	0.0255 (18)	0.0363 (19)	-0.0051 (14)	-0.0167 (16)	0.0013 (14)
C16	0.034 (2)	0.0265 (19)	0.053 (2)	0.0010 (15)	-0.0228 (18)	0.0048 (16)
C17	0.0296 (18)	0.0182 (16)	0.044 (2)	0.0040 (13)	-0.0109 (15)	0.0016 (14)
C18	0.0351 (19)	0.0217 (17)	0.0237 (16)	-0.0009 (13)	-0.0046 (14)	-0.0049 (12)
C19	0.0218 (15)	0.0156 (14)	0.0178 (14)	0.0004 (11)	-0.0007 (11)	-0.0003 (11)
O1W	0.121 (4)	0.069 (3)	0.057 (3)	0.019 (2)	-0.026 (2)	-0.006 (2)

Geometric parameters (\AA , $^\circ$)

Zn1—O1	2.102 (2)	C5—C6	1.336 (6)
Zn1—O2 ⁱ	2.110 (2)	C5—H5A	0.9300
Zn1—O4 ⁱ	2.062 (2)	C6—C7	1.436 (5)
Zn1—N1	2.142 (3)	C6—H6A	0.9300
Zn1—N2	2.139 (3)	C7—C8	1.401 (5)
Zn1—N3	2.131 (3)	C7—C11	1.413 (5)
O1—C19	1.262 (4)	C8—C9	1.355 (6)

O2—C19	1.252 (4)	C8—H8A	0.9300
O3—C18	1.221 (5)	C9—C10	1.405 (5)
O4—C18	1.265 (4)	C9—H9A	0.9300
N1—C1	1.324 (4)	C10—H10A	0.9300
N1—C12	1.365 (4)	C11—C12	1.429 (5)
N2—C10	1.331 (5)	C13—C14	1.386 (4)
N2—C11	1.362 (4)	C13—C19	1.518 (4)
N3—C17	1.333 (4)	C14—C15	1.389 (5)
N3—C13	1.349 (4)	C14—C18	1.516 (4)
C1—C2	1.407 (5)	C15—C16	1.386 (5)
C1—H1A	0.9300	C15—H15A	0.9300
C2—C3	1.369 (6)	C16—C17	1.381 (5)
C2—H2A	0.9300	C16—H16A	0.9300
C3—C4	1.408 (5)	C17—H17A	0.9300
C3—H3A	0.9300	O1W—H1WA	0.85
C4—C12	1.417 (5)	O1W—H1WB	0.85
C4—C5	1.427 (5)		
O4 ⁱ —Zn1—O1	91.81 (9)	C5—C6—H6A	119.3
O4 ⁱ —Zn1—O2 ⁱ	89.56 (9)	C7—C6—H6A	119.3
O1—Zn1—O2 ⁱ	167.49 (9)	C8—C7—C11	117.4 (3)
O4 ⁱ —Zn1—N3	88.28 (10)	C8—C7—C6	124.0 (4)
O1—Zn1—N3	78.00 (9)	C11—C7—C6	118.6 (3)
O2 ⁱ —Zn1—N3	89.61 (10)	C9—C8—C7	120.0 (3)
O4 ⁱ —Zn1—N2	172.42 (11)	C9—C8—H8A	120.0
O1—Zn1—N2	86.07 (10)	C7—C8—H8A	120.0
O2 ⁱ —Zn1—N2	94.09 (11)	C8—C9—C10	119.4 (3)
N3—Zn1—N2	98.36 (11)	C8—C9—H9A	120.3
O4 ⁱ —Zn1—N1	95.17 (11)	C10—C9—H9A	120.3
O1—Zn1—N1	98.93 (10)	N2—C10—C9	122.6 (3)
O2 ⁱ —Zn1—N1	93.33 (10)	N2—C10—H10A	118.7
N3—Zn1—N1	175.47 (10)	C9—C10—H10A	118.7
N2—Zn1—N1	78.01 (11)	N2—C11—C7	122.4 (3)
C19—O1—Zn1	115.6 (2)	N2—C11—C12	117.5 (3)
C19—O2—Zn1 ⁱⁱ	138.9 (2)	C7—C11—C12	120.1 (3)
C18—O4—Zn1 ⁱⁱ	118.5 (2)	N1—C12—C4	122.9 (3)
C1—N1—C12	117.6 (3)	N1—C12—C11	117.8 (3)
C1—N1—Zn1	129.3 (2)	C4—C12—C11	119.3 (3)
C12—N1—Zn1	113.1 (2)	N3—C13—C14	122.4 (3)
C10—N2—C11	118.1 (3)	N3—C13—C19	113.6 (3)
C10—N2—Zn1	128.3 (2)	C14—C13—C19	124.0 (3)
C11—N2—Zn1	113.5 (2)	C13—C14—C15	117.9 (3)
C17—N3—C13	118.8 (3)	C13—C14—C18	123.5 (3)
C17—N3—Zn1	126.6 (2)	C15—C14—C18	118.5 (3)
C13—N3—Zn1	112.7 (2)	C16—C15—C14	119.7 (3)
N1—C1—C2	123.5 (3)	C16—C15—H15A	120.2
N1—C1—H1A	118.3	C14—C15—H15A	120.2

supplementary materials

C2—C1—H1A	118.3	C17—C16—C15	118.7 (3)
C3—C2—C1	119.2 (3)	C17—C16—H16A	120.7
C3—C2—H2A	120.4	C15—C16—H16A	120.7
C1—C2—H2A	120.4	N3—C17—C16	122.4 (3)
C2—C3—C4	119.4 (3)	N3—C17—H17A	118.8
C2—C3—H3A	120.3	C16—C17—H17A	118.8
C4—C3—H3A	120.3	O3—C18—O4	126.0 (3)
C3—C4—C12	117.4 (3)	O3—C18—C14	116.0 (3)
C3—C4—C5	123.6 (3)	O4—C18—C14	117.9 (3)
C12—C4—C5	119.1 (3)	O2—C19—O1	123.7 (3)
C6—C5—C4	121.6 (3)	O2—C19—C13	119.2 (3)
C6—C5—H5A	119.2	O1—C19—C13	117.1 (3)
C4—C5—H5A	119.2	H1WA—O1W—H1WB	108.7
C5—C6—C7	121.4 (3)		
O4 ⁱ —Zn1—O1—C19	88.5 (2)	Zn1—N2—C11—C7	177.2 (3)
O2 ⁱ —Zn1—O1—C19	-7.6 (6)	C10—N2—C11—C12	-179.9 (3)
N3—Zn1—O1—C19	0.6 (2)	Zn1—N2—C11—C12	-2.9 (4)
N2—Zn1—O1—C19	-98.8 (2)	C8—C7—C11—N2	-0.2 (5)
N1—Zn1—O1—C19	-176.0 (2)	C6—C7—C11—N2	177.4 (3)
O4 ⁱ —Zn1—N1—C1	-4.8 (3)	C8—C7—C11—C12	179.9 (3)
O1—Zn1—N1—C1	-97.5 (3)	C6—C7—C11—C12	-2.4 (5)
O2 ⁱ —Zn1—N1—C1	85.0 (3)	C1—N1—C12—C4	0.6 (5)
N2—Zn1—N1—C1	178.5 (3)	Zn1—N1—C12—C4	179.1 (3)
O4 ⁱ —Zn1—N1—C12	176.9 (2)	C1—N1—C12—C11	179.6 (3)
O1—Zn1—N1—C12	84.2 (2)	Zn1—N1—C12—C11	-1.9 (4)
O2 ⁱ —Zn1—N1—C12	-93.3 (2)	C3—C4—C12—N1	-1.8 (5)
N2—Zn1—N1—C12	0.2 (2)	C5—C4—C12—N1	178.4 (3)
O1—Zn1—N2—C10	78.1 (3)	C3—C4—C12—C11	179.2 (3)
O2 ⁱ —Zn1—N2—C10	-89.4 (3)	C5—C4—C12—C11	-0.7 (5)
N3—Zn1—N2—C10	0.8 (3)	N2—C11—C12—N1	3.3 (4)
N1—Zn1—N2—C10	178.1 (3)	C7—C11—C12—N1	-176.9 (3)
O1—Zn1—N2—C11	-98.5 (2)	N2—C11—C12—C4	-177.7 (3)
O2 ⁱ —Zn1—N2—C11	94.0 (2)	C7—C11—C12—C4	2.2 (5)
N3—Zn1—N2—C11	-175.8 (2)	C17—N3—C13—C14	-3.6 (5)
N1—Zn1—N2—C11	1.5 (2)	Zn1—N3—C13—C14	161.9 (2)
O4 ⁱ —Zn1—N3—C17	81.9 (3)	C17—N3—C13—C19	176.9 (3)
O1—Zn1—N3—C17	174.1 (3)	Zn1—N3—C13—C19	-17.7 (3)
O2 ⁱ —Zn1—N3—C17	-7.7 (3)	N3—C13—C14—C15	3.1 (5)
N2—Zn1—N3—C17	-101.8 (3)	C19—C13—C14—C15	-177.4 (3)
O4 ⁱ —Zn1—N3—C13	-82.2 (2)	N3—C13—C14—C18	-173.2 (3)
O1—Zn1—N3—C13	10.0 (2)	C19—C13—C14—C18	6.3 (5)
O2 ⁱ —Zn1—N3—C13	-171.8 (2)	C13—C14—C15—C16	0.4 (5)
N2—Zn1—N3—C13	94.2 (2)	C18—C14—C15—C16	176.8 (4)
C12—N1—C1—C2	1.3 (5)	C14—C15—C16—C17	-3.1 (6)
Zn1—N1—C1—C2	-176.9 (3)	C13—N3—C17—C16	0.6 (5)
N1—C1—C2—C3	-1.9 (6)	Zn1—N3—C17—C16	-162.6 (3)

C1—C2—C3—C4	0.5 (6)	C15—C16—C17—N3	2.7 (6)
C2—C3—C4—C12	1.2 (5)	Zn1 ⁱⁱ —O4—C18—O3	-109.1 (4)
C2—C3—C4—C5	-179.0 (4)	Zn1 ⁱⁱ —O4—C18—C14	74.6 (4)
C3—C4—C5—C6	179.6 (4)	C13—C14—C18—O3	95.4 (5)
C12—C4—C5—C6	-0.6 (6)	C15—C14—C18—O3	-80.8 (5)
C4—C5—C6—C7	0.3 (6)	C13—C14—C18—O4	-87.9 (4)
C5—C6—C7—C8	178.7 (4)	C15—C14—C18—O4	95.8 (4)
C5—C6—C7—C11	1.2 (6)	Zn1 ⁱⁱ —O2—C19—O1	-143.6 (3)
C11—C7—C8—C9	0.4 (5)	Zn1 ⁱⁱ —O2—C19—C13	35.2 (5)
C6—C7—C8—C9	-177.1 (4)	Zn1—O1—C19—O2	168.5 (3)
C7—C8—C9—C10	-0.5 (6)	Zn1—O1—C19—C13	-10.3 (4)
C11—N2—C10—C9	-0.4 (5)	N3—C13—C19—O2	-159.6 (3)
Zn1—N2—C10—C9	-176.9 (3)	C14—C13—C19—O2	20.9 (5)
C8—C9—C10—N2	0.5 (6)	N3—C13—C19—O1	19.2 (4)
C10—N2—C11—C7	0.3 (5)	C14—C13—C19—O1	-160.3 (3)

Symmetry codes: (i) $-x+1/2, y-1/2, z$; (ii) $-x+1/2, y+1/2, z$.

Hydrogen-bond geometry ($\text{\AA}, ^\circ$)

<i>D</i> —H \cdots <i>A</i>	<i>D</i> —H	H \cdots <i>A</i>	<i>D</i> \cdots <i>A</i>	<i>D</i> —H \cdots <i>A</i>
O1W—H1WA \cdots O3	0.85	2.22	3.022 (5)	157
O1W—H1WB \cdots O3 ⁱⁱ	0.85	2.42	3.215 (5)	157

Symmetry codes: (ii) $-x+1/2, y+1/2, z$.

Fig. 1




Fig. 2

