

## Bis( $\mu$ -methanolato- $\kappa^2$ O:O)bis{[4-bromo- $N'$ -(1-methyl-3-oxidobut-2-en-1-ylidene- $\kappa$ O)benzohydrazidato- $\kappa^2$ $N',O$ ]oxido-vanadium(V)}

Hon Wee Wong, Kong Mun Lo and Seik Weng Ng\*

 Department of Chemistry, University of Malaya, 50603 Kuala Lumpur, Malaysia  
 Correspondence e-mail: seikweng@um.edu.my


Received 30 May 2011; accepted 5 June 2011

 Key indicators: single-crystal X-ray study;  $T = 100$  K; mean  $\sigma(\text{C}-\text{C}) = 0.003$  Å; disorder in main residue;  $R$  factor = 0.023;  $wR$  factor = 0.060; data-to-parameter ratio = 15.4.

The dinuclear compound,  $[\text{V}_2(\text{C}_{12}\text{H}_{11}\text{BrN}_2\text{O}_2)_2(\text{CH}_3\text{O})_2\text{O}_2]$ , lies on a center of inversion. The doubly-deprotonated Schiff base  $O,N,O'$ -chelates to the  $\text{V}^{\text{V}}$  atom; two metal atoms are bridged by the methoxide units. The coordination geometry is a distorted octahedron. Weak intermolecular  $\text{C}-\text{H}\cdots\text{N}$  hydrogen bonding is present in the crystal structure. The bromophenyl unit is disordered over two positions, with the major component being in a 0.909 (6) proportion.

### Related literature

For the isotypic compound that has chlorine in place of bromine, see: Sarkar & Pal (2009).



### Experimental

#### Crystal data

$[\text{V}_2(\text{C}_{12}\text{H}_{11}\text{BrN}_2\text{O}_2)_2(\text{CH}_3\text{O})_2\text{O}_2]$	$a = 8.7517$ (5) Å
$M_r = 786.22$	$b = 12.3409$ (7) Å
Monoclinic, $P2_1/c$	$c = 13.9952$ (8) Å

$\beta = 101.8782$ (7)°
$V = 1479.17$ (15) Å <sup>3</sup>
$Z = 2$
Mo $K\alpha$ radiation

$\mu = 3.39$ mm <sup>-1</sup>
$T = 100$ K
$0.30 \times 0.25 \times 0.20$ mm

#### Data collection

Bruker SMART APEX diffractometer	18572 measured reflections
Absorption correction: multi-scan (SADABS; Sheldrick, 1996)	3400 independent reflections
$T_{\text{min}} = 0.430$ , $T_{\text{max}} = 0.551$	3106 reflections with $I > 2\sigma(I)$
	$R_{\text{int}} = 0.027$

#### Refinement

$R[F^2 > 2\sigma(F^2)] = 0.023$	44 restraints
$wR(F^2) = 0.060$	H-atom parameters constrained
$S = 0.99$	$\Delta\rho_{\text{max}} = 0.96$ e Å <sup>-3</sup>
3400 reflections	$\Delta\rho_{\text{min}} = -0.32$ e Å <sup>-3</sup>
221 parameters	

**Table 1**

Selected bond lengths (Å).

V1—O1	1.9314 (13)	V1—O4	2.3459 (13)
V1—O2	1.8607 (13)	V1—O4 <sup>i</sup>	1.8289 (13)
V1—O3	1.5896 (14)	V1—N2	2.0770 (15)

 Symmetry code: (i)  $-x, -y + 1, -z + 1$ .

**Table 2**

Hydrogen-bond geometry (Å, °).

$D-\text{H}\cdots A$	$D-\text{H}$	$\text{H}\cdots A$	$D\cdots A$	$D-\text{H}\cdots A$
$\text{C}12-\text{H}12\text{C}\cdots\text{N}1^{\text{ii}}$	0.96	2.59	3.541 (5)	169

 Symmetry code: (ii)  $x, -y + \frac{3}{2}, z + \frac{1}{2}$ .

Data collection: *APEX2* (Bruker, 2009); cell refinement: *SAINT* (Bruker, 2009); data reduction: *SAINT*; program(s) used to solve structure: *SHELXS97* (Sheldrick, 2008); program(s) used to refine structure: *SHELXL97* (Sheldrick, 2008); molecular graphics: *X-SEED* (Barbour, 2001); software used to prepare material for publication: *pubCIF* (Westrip, 2010).

We thank the University of Malaya (RG020/09AFR) for supporting this study.

Supplementary data and figures for this paper are available from the IUCr electronic archives (Reference: XU5231).

### References

- Barbour, L. J. (2001). *J. Supramol. Chem.* **1**, 189–191.  
 Bruker (2009). *APEX2* and *SAINT*. Bruker AXS Inc., Madison, Wisconsin, USA.  
 Sarkar, A. & Pal, S. (2009). *Inorg. Chim. Acta*, **362**, 3807–3812.  
 Sheldrick, G. M. (1996). *SADABS*. University of Göttingen, Germany.  
 Sheldrick, G. M. (2008). *Acta Cryst.* **A64**, 112–122.  
 Westrip, S. P. (2010). *J. Appl. Cryst.* **43**, 920–925.

**supplementary materials**

*Acta Cryst.* (2011). E67, m902 [ doi:10.1107/S1600536811021647 ]

**Bis( $\mu$ -methanolato- $\kappa^2 O:O$ )bis{[4-bromo- $N'$ -(1-methyl-3-oxidobut-2-en-1-ylidene- $\kappa O$ )benzohydrazidato- $\kappa^2 N',O$ ]oxidovanadium(V)}**

**H. W. Wong, K. M. Lo and S. W. Ng**

**Comment**

A recent study detailed the crystal structure of  $[\text{VO}(\text{C}_{12}\text{H}_{11}\text{ClN}_2\text{O}_2)(\text{CH}_3\text{O})_2]$  (Sarkar & Pal, 2009). The title bromo analog (Scheme I) is isostructural, the two compounds crystallizing with matching cell dimensions. However, the title compound shows some disorder in the halophenyl portion (Fig. 1). Dinuclear  $[\text{VO}(\text{C}_{12}\text{H}_{11}\text{BrN}_2\text{O}_2)(\text{CH}_3\text{O})_2]$  lies on a center of inversion. The doubly-deprotonated Schiff base  $O,N,O'$ -chelates to the  $\text{V}^{\text{V}}$  atom; two metal centers are bridged by the methoxide unit. The geometry is an octahedron.

**Experimental**

Bis(acetylacetonato)oxovanadium (0.67 g, 2.5 mmol) was heated with 4-bromobenzoic acid hydrazide (0.54 g, 2.5 mmol) in methanol (50 ml) for 1 h. The solution mixture was then filtered and upon slow cooling of the filtrate, dark brown crystals separated out.

**Refinement**

Carbon-bound H-atoms were placed in calculated positions (C–H 0.95 to 0.98 Å) and were included in the refinement in the riding model approximation, with  $U(\text{H})$  set to 1.2–1.5 $U(\text{C})$ .

The bromophenyl portion of the ligand is disordered over two positions. The pair of carbon–bromine distances were restrained to within  $\pm 0.01$  Å each other. For the ring, the 1,2-related distances were restrained to  $1.39 \pm 0.01$  Å, the 1,3-related distances to  $2.41 \pm 0.01$  Å and the 1,4-related distances to  $2.78 \pm 0.01$  Å. Each seven-atom component was restrained to near planarity. Owing to the low degree of disorder, additional restraints were imposed on the primed bromine atom as well as the primed *para*-carbon atom. Distance restraints were applied so that the angle at the carbon atoms was approximately  $120^\circ$ . The isotropic temperature factors of the primed atoms were set to the anisotropic temperature factors of the unprimed ones. The disorder refined to 0.909 (1): 0.091.

**Figures**




Fig. 1. Thermal ellipsoid plot (Barbour, 2001) of  $[\text{VO}(\text{C}_{12}\text{H}_{11}\text{BrN}_2\text{O}_2)(\text{CH}_3\text{O})_2]$  at the 70% probability level; hydrogen atoms are drawn as spheres of arbitrary radius. The disorder in bromophenyl ring is not shown.

# supplementary materials

---

## Bis( $\mu$ -methanolato- $\kappa^2$ O:O)bis{[4-bromo-*N*'-(1-methyl-3-oxidobut-2-en-1-ylidene- $\kappa$ O)benzohydrazidato- $\kappa^2$ N',O]oxidovanadium(V)}

### Crystal data

$[\text{V}_2(\text{C}_{12}\text{H}_{11}\text{BrN}_2\text{O}_2)_2(\text{CH}_3\text{O})_2\text{O}_2]$	$F(000) = 784$
$M_r = 786.22$	$D_x = 1.765 \text{ Mg m}^{-3}$
Monoclinic, $P2_1/c$	Mo $K\alpha$ radiation, $\lambda = 0.71073 \text{ \AA}$
Hall symbol: -P 2ybc	Cell parameters from 8361 reflections
$a = 8.7517 (5) \text{ \AA}$	$\theta = 2.2\text{--}28.2^\circ$
$b = 12.3409 (7) \text{ \AA}$	$\mu = 3.39 \text{ mm}^{-1}$
$c = 13.9952 (8) \text{ \AA}$	$T = 100 \text{ K}$
$\beta = 101.8782 (7)^\circ$	Prism, brown
$V = 1479.17 (15) \text{ \AA}^3$	$0.30 \times 0.25 \times 0.20 \text{ mm}$
$Z = 2$	

### Data collection

Bruker SMART APEX diffractometer	3400 independent reflections
Radiation source: fine-focus sealed tube graphite	3106 reflections with $I > 2\sigma(I)$
$\omega$ scans	$R_{\text{int}} = 0.027$
Absorption correction: multi-scan (SADABS; Sheldrick, 1996)	$\theta_{\text{max}} = 27.5^\circ$ , $\theta_{\text{min}} = 2.2^\circ$
$T_{\text{min}} = 0.430$ , $T_{\text{max}} = 0.551$	$h = -11 \rightarrow 11$
18572 measured reflections	$k = -16 \rightarrow 16$
	$l = -18 \rightarrow 18$

### Refinement

Refinement on $F^2$	Primary atom site location: structure-invariant direct methods
Least-squares matrix: full	Secondary atom site location: difference Fourier map
$R[F^2 > 2\sigma(F^2)] = 0.023$	Hydrogen site location: inferred from neighbouring sites
$wR(F^2) = 0.060$	H-atom parameters constrained
$S = 0.99$	$w = 1/[\sigma^2(F_o^2) + (0.0308P)^2 + 1.3638P]$
3400 reflections	where $P = (F_o^2 + 2F_c^2)/3$
221 parameters	$(\Delta/\sigma)_{\text{max}} = 0.001$
44 restraints	$\Delta\rho_{\text{max}} = 0.96 \text{ e \AA}^{-3}$
	$\Delta\rho_{\text{min}} = -0.32 \text{ e \AA}^{-3}$

### Fractional atomic coordinates and isotropic or equivalent isotropic displacement parameters ( $\text{\AA}^2$ )

	<i>x</i>	<i>y</i>	<i>z</i>	$U_{\text{iso}}^*/U_{\text{eq}}$	Occ. ( $<1$ )
Br1	0.53762 (15)	0.29274 (14)	0.06235 (3)	0.0224 (2)	0.909 (6)

Br1'	0.5773 (11)	0.3313 (11)	0.0656 (3)	0.0164 (15)	0.091 (6)
V1	0.11414 (4)	0.60960 (2)	0.50419 (2)	0.01169 (8)	
O1	0.21661 (15)	0.52544 (10)	0.41937 (9)	0.0152 (3)	
O2	-0.04425 (16)	0.69868 (10)	0.52973 (9)	0.0149 (3)	
O3	0.25986 (16)	0.67280 (11)	0.56588 (10)	0.0179 (3)	
O4	-0.10415 (15)	0.51204 (10)	0.42270 (9)	0.0126 (2)	
N1	0.15083 (18)	0.65573 (13)	0.30152 (11)	0.0142 (3)	
N2	0.08999 (18)	0.70085 (12)	0.37726 (11)	0.0125 (3)	
C1	0.4474 (2)	0.3788 (2)	0.14888 (15)	0.0186 (5)	0.909 (6)
C1'	0.4680 (11)	0.4032 (11)	0.1483 (7)	0.019*	0.091 (6)
C2	0.4295 (3)	0.3360 (2)	0.23695 (18)	0.0202 (6)	0.909 (6)
H2	0.4659	0.2650	0.2558	0.024*	0.909 (6)
C2'	0.455 (3)	0.3560 (15)	0.2361 (13)	0.020*	0.091 (6)
H2'	0.5014	0.2875	0.2544	0.024*	0.091 (6)
C3	0.3573 (3)	0.39817 (19)	0.29774 (18)	0.0181 (5)	0.909 (6)
H3	0.3448	0.3697	0.3587	0.022*	0.909 (6)
C3'	0.373 (3)	0.4097 (14)	0.2974 (13)	0.018*	0.091 (6)
H3'	0.3632	0.3780	0.3577	0.022*	0.091 (6)
C4	0.3030 (2)	0.50212 (18)	0.26988 (16)	0.0154 (4)	0.909 (6)
C4'	0.3054 (10)	0.5095 (9)	0.2697 (8)	0.015*	0.091 (6)
C5	0.3265 (4)	0.5453 (2)	0.18142 (19)	0.0177 (4)	0.909 (6)
H5	0.2927	0.6168	0.1628	0.021*	0.909 (6)
C5'	0.319 (3)	0.5568 (15)	0.1819 (13)	0.018*	0.091 (6)
H5'	0.2728	0.6254	0.1635	0.021*	0.091 (6)
C6	0.3994 (3)	0.4833 (2)	0.12096 (18)	0.0194 (5)	0.909 (6)
H6	0.4161	0.5122	0.0610	0.023*	0.909 (6)
C6'	0.401 (3)	0.5032 (15)	0.1208 (13)	0.019*	0.091 (6)
H6'	0.4110	0.5349	0.0606	0.023*	0.091 (6)
C7	0.2196 (2)	0.56521 (15)	0.33309 (13)	0.0136 (3)	
C8	0.0615 (2)	0.86357 (16)	0.27592 (14)	0.0179 (4)	
H8A	0.1688	0.8559	0.2664	0.027*	
H8B	0.0386	0.9404	0.2839	0.027*	
H8C	-0.0112	0.8345	0.2189	0.027*	
C9	0.0435 (2)	0.80226 (14)	0.36548 (13)	0.0136 (3)	
C10	-0.0241 (2)	0.85486 (15)	0.43769 (13)	0.0149 (4)	
H10	-0.0407	0.9309	0.4319	0.018*	
C11	-0.0663 (2)	0.80336 (14)	0.51441 (13)	0.0138 (3)	
C12	-0.1443 (2)	0.85908 (15)	0.58647 (13)	0.0166 (4)	
H12A	-0.2375	0.8183	0.5932	0.025*	
H12B	-0.1745	0.9325	0.5635	0.025*	
H12C	-0.0718	0.8629	0.6499	0.025*	
C13	-0.2359 (2)	0.56198 (17)	0.36097 (15)	0.0209 (4)	
H13A	-0.2989	0.5065	0.3209	0.031*	
H13B	-0.1997	0.6151	0.3185	0.031*	
H13C	-0.2994	0.5986	0.4013	0.031*	

## supplementary materials

---

### Atomic displacement parameters ( $\text{\AA}^2$ )

	$U^{11}$	$U^{22}$	$U^{33}$	$U^{12}$	$U^{13}$	$U^{23}$
Br1	0.0193 (3)	0.0288 (5)	0.01969 (13)	0.0092 (3)	0.00564 (12)	-0.00508 (14)
Br1'	0.0125 (19)	0.018 (3)	0.0200 (12)	-0.001 (2)	0.0061 (10)	-0.0048 (12)
V1	0.01496 (16)	0.01018 (15)	0.01058 (15)	-0.00098 (11)	0.00417 (11)	0.00064 (11)
O1	0.0180 (6)	0.0146 (6)	0.0144 (6)	0.0015 (5)	0.0069 (5)	0.0018 (5)
O2	0.0203 (7)	0.0113 (6)	0.0148 (6)	-0.0004 (5)	0.0073 (5)	0.0003 (5)
O3	0.0210 (7)	0.0173 (7)	0.0152 (6)	-0.0046 (5)	0.0033 (5)	0.0012 (5)
O4	0.0145 (6)	0.0121 (6)	0.0114 (6)	0.0006 (5)	0.0030 (5)	0.0018 (5)
N1	0.0173 (8)	0.0144 (7)	0.0119 (7)	-0.0009 (6)	0.0056 (6)	-0.0015 (6)
N2	0.0140 (7)	0.0128 (7)	0.0114 (7)	-0.0009 (6)	0.0045 (6)	-0.0006 (6)
C1	0.0122 (9)	0.0263 (12)	0.0180 (10)	0.0034 (8)	0.0048 (8)	-0.0065 (9)
C2	0.0191 (13)	0.0208 (12)	0.0203 (10)	0.0059 (10)	0.0031 (9)	-0.0011 (9)
C3	0.0184 (12)	0.0198 (11)	0.0170 (10)	0.0021 (9)	0.0058 (8)	-0.0004 (8)
C4	0.0132 (9)	0.0176 (10)	0.0160 (9)	-0.0016 (8)	0.0046 (8)	-0.0020 (8)
C5	0.0184 (11)	0.0178 (11)	0.0180 (10)	0.0001 (9)	0.0061 (8)	-0.0006 (8)
C6	0.0174 (10)	0.0263 (13)	0.0160 (10)	0.0001 (10)	0.0067 (8)	-0.0012 (9)
C7	0.0131 (8)	0.0147 (8)	0.0135 (8)	-0.0036 (7)	0.0033 (7)	-0.0018 (7)
C8	0.0236 (10)	0.0159 (9)	0.0150 (9)	0.0020 (7)	0.0058 (7)	0.0037 (7)
C9	0.0126 (8)	0.0149 (9)	0.0130 (8)	-0.0012 (7)	0.0016 (7)	0.0010 (7)
C10	0.0189 (9)	0.0102 (8)	0.0154 (9)	0.0004 (7)	0.0033 (7)	-0.0002 (7)
C11	0.0141 (8)	0.0124 (8)	0.0142 (8)	0.0003 (6)	0.0011 (7)	-0.0023 (7)
C12	0.0204 (9)	0.0155 (9)	0.0146 (9)	0.0019 (7)	0.0053 (7)	-0.0020 (7)
C13	0.0184 (9)	0.0194 (10)	0.0218 (10)	0.0005 (7)	-0.0032 (8)	0.0065 (8)

### Geometric parameters ( $\text{\AA}$ , $^\circ$ )

Br1—C1	1.9002 (19)	C3'—H3'	0.9500
Br1'—C1'	1.870 (8)	C4—C5	1.401 (3)
V1—O1	1.9314 (13)	C4—C7	1.479 (3)
V1—O2	1.8607 (13)	C4'—C5'	1.389 (9)
V1—O3	1.5896 (14)	C4'—C7	1.448 (8)
V1—O4	2.3459 (13)	C5—C6	1.389 (3)
V1—O4 <sup>i</sup>	1.8289 (13)	C5—H5	0.9500
V1—N2	2.0770 (15)	C5'—C6'	1.390 (9)
O1—C7	1.309 (2)	C5'—H5'	0.9500
O2—C11	1.317 (2)	C6—H6	0.9500
O4—C13	1.430 (2)	C6'—H6'	0.9500
O4—V1 <sup>i</sup>	1.8289 (13)	C8—C9	1.500 (2)
N1—C7	1.302 (2)	C8—H8A	0.9800
N1—N2	1.396 (2)	C8—H8B	0.9800
N2—C9	1.316 (2)	C8—H8C	0.9800
C1—C2	1.380 (3)	C9—C10	1.428 (3)
C1—C6	1.387 (3)	C10—C11	1.363 (3)
C1'—C6'	1.385 (9)	C10—H10	0.9500
C1'—C2'	1.386 (9)	C11—C12	1.496 (2)

C2—C3	1.391 (3)	C12—H12A	0.9800
C2—H2	0.9500	C12—H12B	0.9800
C2'—C3'	1.394 (9)	C12—H12C	0.9800
C2'—H2'	0.9500	C13—H13A	0.9800
C3—C4	1.395 (3)	C13—H13B	0.9800
C3—H3	0.9500	C13—H13C	0.9800
C3'—C4'	1.387 (9)		
O3—V1—O4 <sup>i</sup>	102.96 (6)	C3'—C4'—C7	119.7 (7)
O3—V1—O2	98.74 (7)	C5'—C4'—C7	119.6 (7)
O4 <sup>i</sup> —V1—O2	104.72 (6)	C6—C5—C4	119.90 (19)
O3—V1—O1	100.11 (7)	C6—C5—H5	120.1
O4 <sup>i</sup> —V1—O1	89.10 (6)	C4—C5—H5	120.1
O2—V1—O1	153.42 (6)	C4'—C5'—C6'	119.5 (6)
O3—V1—N2	97.33 (6)	C4'—C5'—H5'	120.2
O4 <sup>i</sup> —V1—N2	156.26 (6)	C6'—C5'—H5'	120.2
O2—V1—N2	83.99 (6)	C1—C6—C5	119.38 (19)
O1—V1—N2	75.18 (6)	C1—C6—H6	120.3
O3—V1—O4	176.30 (6)	C5—C6—H6	120.3
O4 <sup>i</sup> —V1—O4	73.91 (6)	C1'—C6'—C5'	119.7 (6)
O2—V1—O4	80.35 (5)	C1'—C6'—H6'	120.1
O1—V1—O4	81.95 (5)	C5'—C6'—H6'	120.1
N2—V1—O4	86.17 (5)	N1—C7—O1	122.62 (16)
C7—O1—V1	117.75 (12)	N1—C7—C4'	117.5 (5)
C11—O2—V1	129.67 (12)	O1—C7—C4'	119.9 (5)
C13—O4—V1 <sup>i</sup>	124.48 (12)	N1—C7—C4	119.95 (17)
C13—O4—V1	123.17 (11)	O1—C7—C4	117.43 (17)
V1 <sup>i</sup> —O4—V1	106.09 (6)	C9—C8—H8A	109.5
C7—N1—N2	107.84 (14)	C9—C8—H8B	109.5
C9—N2—N1	116.14 (15)	H8A—C8—H8B	109.5
C9—N2—V1	126.68 (12)	C9—C8—H8C	109.5
N1—N2—V1	116.45 (11)	H8A—C8—H8C	109.5
C2—C1—C6	121.62 (19)	H8B—C8—H8C	109.5
C2—C1—Br1	119.58 (16)	N2—C9—C10	120.41 (16)
C6—C1—Br1	118.79 (16)	N2—C9—C8	120.12 (17)
C6'—C1'—C2'	120.9 (6)	C10—C9—C8	119.48 (16)
C6'—C1'—Br1'	119.5 (6)	C11—C10—C9	124.39 (17)
C2'—C1'—Br1'	119.7 (6)	C11—C10—H10	117.8
C1—C2—C3	118.99 (19)	C9—C10—H10	117.8
C1—C2—H2	120.5	O2—C11—C10	122.07 (17)
C3—C2—H2	120.5	O2—C11—C12	114.39 (16)
C1'—C2'—C3'	119.5 (6)	C10—C11—C12	123.52 (17)
C1'—C2'—H2'	120.3	C11—C12—H12A	109.5
C3'—C2'—H2'	120.3	C11—C12—H12B	109.5
C2—C3—C4	120.53 (19)	H12A—C12—H12B	109.5
C2—C3—H3	119.7	C11—C12—H12C	109.5
C4—C3—H3	119.7	H12A—C12—H12C	109.5
C4'—C3'—C2'	119.6 (6)	H12B—C12—H12C	109.5

## supplementary materials

---

C4'—C3'—H3'	120.2	O4—C13—H13A	109.5
C2'—C3'—H3'	120.2	O4—C13—H13B	109.5
C3—C4—C5	119.52 (18)	H13A—C13—H13B	109.5
C3—C4—C7	119.89 (19)	O4—C13—H13C	109.5
C5—C4—C7	120.59 (19)	H13A—C13—H13C	109.5
C3'—C4'—C5'	120.7 (6)	H13B—C13—H13C	109.5
O3—V1—O1—C7	-94.55 (13)	C2'—C3'—C4'—C7	179.7 (4)
O4 <sup>i</sup> —V1—O1—C7	162.44 (13)	C3—C4—C5—C6	1.9 (3)
O2—V1—O1—C7	40.0 (2)	C7—C4—C5—C6	-177.37 (19)
N2—V1—O1—C7	0.43 (12)	C3'—C4'—C5'—C6'	0.2 (9)
O4—V1—O1—C7	88.57 (13)	C7—C4'—C5'—C6'	-179.8 (6)
O3—V1—O2—C11	56.92 (16)	C2—C1—C6—C5	-2.2 (3)
O4 <sup>i</sup> —V1—O2—C11	162.88 (15)	Br1—C1—C6—C5	176.95 (15)
O1—V1—O2—C11	-77.8 (2)	C4—C5—C6—C1	0.3 (3)
N2—V1—O2—C11	-39.60 (16)	C2'—C1'—C6'—C5'	-0.2 (7)
O4—V1—O2—C11	-126.72 (16)	Br1'—C1'—C6'—C5'	179.7 (5)
O4 <sup>i</sup> —V1—O4—C13	153.18 (16)	C4'—C5'—C6'—C1'	0.0 (9)
O2—V1—O4—C13	44.67 (14)	N2—N1—C7—O1	4.5 (2)
O1—V1—O4—C13	-115.43 (14)	N2—N1—C7—C4'	-173.8 (4)
N2—V1—O4—C13	-39.87 (14)	N2—N1—C7—C4	-176.61 (16)
O4 <sup>i</sup> —V1—O4—V1 <sup>i</sup>	0.0	V1—O1—C7—N1	-3.1 (2)
O2—V1—O4—V1 <sup>i</sup>	-108.51 (7)	V1—O1—C7—C4'	175.2 (4)
O1—V1—O4—V1 <sup>i</sup>	91.39 (6)	V1—O1—C7—C4	177.94 (13)
N2—V1—O4—V1 <sup>i</sup>	166.95 (7)	C3'—C4'—C7—N1	-176.1 (15)
C7—N1—N2—C9	167.03 (16)	C5'—C4'—C7—N1	3.8 (16)
C7—N1—N2—V1	-3.84 (18)	C3'—C4'—C7—O1	5.5 (16)
O3—V1—N2—C9	-69.23 (16)	C5'—C4'—C7—O1	-174.6 (15)
O4 <sup>i</sup> —V1—N2—C9	142.13 (16)	C3'—C4'—C7—C4	-40 (8)
O2—V1—N2—C9	28.85 (16)	C5'—C4'—C7—C4	140 (8)
O1—V1—N2—C9	-167.80 (16)	C3—C4—C7—N1	-170.83 (19)
O4—V1—N2—C9	109.53 (15)	C5—C4—C7—N1	8.4 (3)
O3—V1—N2—N1	100.55 (13)	C3—C4—C7—O1	8.1 (3)
O4 <sup>i</sup> —V1—N2—N1	-48.1 (2)	C5—C4—C7—O1	-172.6 (2)
O2—V1—N2—N1	-161.38 (13)	C3—C4—C7—C4'	144 (8)
O1—V1—N2—N1	1.97 (11)	C5—C4—C7—C4'	-37 (8)
O4—V1—N2—N1	-80.70 (12)	N1—N2—C9—C10	178.40 (16)
C6—C1—C2—C3	1.91 (17)	V1—N2—C9—C10	-11.8 (3)
Br1—C1—C2—C3	-177.28 (11)	N1—N2—C9—C8	-1.7 (2)
C6'—C1'—C2'—C3'	0.2 (3)	V1—N2—C9—C8	168.13 (13)
Br1'—C1'—C2'—C3'	-179.8 (2)	N2—C9—C10—C11	-9.9 (3)
C1—C2—C3—C4	0.34 (15)	C8—C9—C10—C11	170.21 (18)
C1'—C2'—C3'—C4'	0.0 (3)	V1—O2—C11—C10	32.7 (3)
C2—C3—C4—C5	-2.2 (2)	V1—O2—C11—C12	-148.61 (13)
C2—C3—C4—C7	177.03 (14)	C9—C10—C11—O2	1.7 (3)
C2'—C3'—C4'—C5'	-0.2 (7)	C9—C10—C11—C12	-176.86 (18)

Symmetry codes: (i)  $-x, -y+1, -z+1$ .



*Hydrogen-bond geometry* (Å, °)

<i>D</i> —H··· <i>A</i>	<i>D</i> —H	H··· <i>A</i>	<i>D</i> ··· <i>A</i>	<i>D</i> —H··· <i>A</i>
C12—H12C···N1 <sup>ii</sup>	0.96	2.59	3.541 (5)	169

Symmetry codes: (ii)  $x, -y+3/2, z+1/2$ .

Fig. 1

