

4-[(*E*)-(4-Fluorobenzylidene)amino]-phenol

Li-Xia Sun, Yun-Dan Yu and Guo-Ying Wei*

College of Materials Science & Engineering, China Jiliang University, Hangzhou 310053, People's Republic of China
Correspondence e-mail: nanocrystal11@163.com


Received 25 May 2011; accepted 25 May 2011

Key indicators: single-crystal X-ray study; $T = 296\text{ K}$; mean $\sigma(\text{C}-\text{C}) = 0.003\text{ \AA}$; R factor = 0.040; wR factor = 0.116; data-to-parameter ratio = 13.2.

In the title compound, $\text{C}_{13}\text{H}_{10}\text{FNO}$, the dihedral angle between the aromatic rings is $55.60(8)^\circ$. In the crystal, molecules are linked by $\text{O}-\text{H}-\text{N}$ hydrogen bonds, forming zigzag $C(7)$ chains propagating in [101].

Related literature

For a related structure and background references, see: Sun *et al.* (2011). For related structures, see: Nie *et al.* (2008); Fun *et al.* (2008); Alhadi *et al.* (2008).



Experimental

Crystal data

$\text{C}_{13}\text{H}_{10}\text{FNO}$

$M_r = 215.22$

Monoclinic, $P2_1/n$

$a = 9.400(5)\text{ \AA}$

$b = 12.213(7)\text{ \AA}$

$c = 9.450(5)\text{ \AA}$

$\beta = 104.666(5)^\circ$
 $V = 1049.5(10)\text{ \AA}^3$
 $Z = 4$
Mo $K\alpha$ radiation

$\mu = 0.10\text{ mm}^{-1}$
 $T = 296\text{ K}$
 $0.25 \times 0.23 \times 0.22\text{ mm}$

Data collection

Bruker APEXII CCD
diffractometer
Absorption correction: multi-scan
(*SADABS*; Bruker, 2004)
 $T_{\min} = 0.976$, $T_{\max} = 0.979$

5880 measured reflections
1935 independent reflections
1474 reflections with $I > 2\sigma(I)$
 $R_{\text{int}} = 0.036$

Refinement

$R[F^2 > 2\sigma(F^2)] = 0.040$
 $wR(F^2) = 0.116$
 $S = 1.09$
1935 reflections

147 parameters
H-atom parameters constrained
 $\Delta\rho_{\max} = 0.18\text{ e \AA}^{-3}$
 $\Delta\rho_{\min} = -0.19\text{ e \AA}^{-3}$

Table 1
Hydrogen-bond geometry (\AA , $^\circ$).

$D-\text{H}\cdots A$	$D-\text{H}$	$\text{H}\cdots A$	$D\cdots A$	$D-\text{H}\cdots A$
O1—H1A \cdots N1 ⁱ	0.82	2.09	2.885 (2)	163
Symmetry code: (i) $x - \frac{1}{2}, -y + \frac{3}{2}, z - \frac{1}{2}$.				

Data collection: *APEX2* (Bruker, 2004); cell refinement: *SAINT* (Bruker, 2004); data reduction: *SAINT*; program(s) used to solve structure: *SHELXS97* (Sheldrick, 2008); program(s) used to refine structure: *SHELXL97* (Sheldrick, 2008); molecular graphics: *SHELXTL* (Sheldrick, 2008); software used to prepare material for publication: *SHELXTL*.

This project was supported by Zhejiang Provincial Natural Science Foundation of China (grant No. Y4110290).

Supplementary data and figures for this paper are available from the IUCr electronic archives (Reference: HB5896).

References

- Alhadi, A. A., Ali, H. M., Puvaneswary, S., Robinson, W. T. & Ng, S. W. (2008). *Acta Cryst. E64*, o1584.
Bruker (2004). *APEX2*, *SAINT* and *SADABS*. Bruker AXS Inc., Madison, Wisconsin, USA.
Fun, H.-K., Patil, P. S., Jebas, S. R., Sujith, K. V. & Kalluraya, B. (2008). *Acta Cryst. E64*, o1594–o1595.
Nie, Y. (2008). *Acta Cryst. E64*, o471.
Sheldrick, G. M. (2008). *Acta Cryst. A64*, 112–122.
Sun, L.-X., Yu, Y.-D. & Wei, G.-Y. (2011). *Acta Cryst. E67*, o1564.

supplementary materials

Acta Cryst. (2011). E67, o1578 [doi:10.1107/S1600536811019921]

4-[*(E*)-(4-Fluorobenzylidene)amino]phenol

L.-X. Sun, Y.-D. Yu and G.-Y. Wei

Comment

As part of our ongoing studies of Schiff bases (Sun *et al.*, 2011), we report here the crystal structure of the title compound, (I). In (I) (Fig. 1), the bond lengths and angles are normal and comparable to the values observed in similar compounds (Nie *et al.*, 2008; Fun *et al.*, 2008; Alhadi *et al.*, 2008). The dihedral angle between the two aromatic rings in the Schiff base molecule is 55.6° , indicating that two these rings are not coplanar. Intermolecular O—H—N hydrogen bonds (Table 1) link the molecules along *a* axis (Fig. 2).

Experimental

A mixture of 4-chlorobenzaldehyde (5 mmol), 4-aminophenol (5 mmol) and methanol (40 ml) was refluxed for 2 h. It was then allowed to cool and filtered. Recrystallization of the crude product from methanol yielded yellow blocks of (I).

Refinement

H atoms were positioned geometrically and refined using the riding-model approximation, with C—H = 0.93–0.97 Å, O—H = 0.82 Å, and $U_{\text{iso}}(\text{H}) = 1.2U_{\text{eq}}(\text{C})$ or $U_{\text{iso}}(\text{H}) = 1.5U_{\text{eq}}(\text{O})$. In the absence of significant anomalous dispersion effects, Freidel pairs were merged.

Figures




Fig. 1. The molecular structure of the title compounds with 50% probability displacement ellipsoids for non-hydrogen atoms.




Fig. 2. Molecular packing of the title compound, viewed along the *a* axis. Hydrogen bonds are shown as dashed lines.

supplementary materials

4-[(*E*)-(4-Fluorobenzylidene)amino]phenol

Crystal data

C ₁₃ H ₁₀ FNO	<i>F</i> (000) = 448
<i>M_r</i> = 215.22	<i>D_x</i> = 1.362 Mg m ⁻³
Monoclinic, <i>P2₁/n</i>	Mo <i>Kα</i> radiation, λ = 0.71073 Å
Hall symbol: -P 2yn	Cell parameters from 2754 reflections
<i>a</i> = 9.400 (5) Å	θ = 2.7–28.2°
<i>b</i> = 12.213 (7) Å	μ = 0.10 mm ⁻¹
<i>c</i> = 9.450 (5) Å	<i>T</i> = 296 K
β = 104.666 (5)°	Block, yellow
<i>V</i> = 1049.5 (10) Å ³	0.25 × 0.23 × 0.22 mm
<i>Z</i> = 4	

Data collection

Bruker APEXII CCD diffractometer	1935 independent reflections
Radiation source: fine-focus sealed tube graphite	1474 reflections with $I > 2\sigma(I)$
φ and ω scans	$R_{\text{int}} = 0.036$
Absorption correction: multi-scan (<i>SADABS</i> ; Bruker, 2004)	$\theta_{\text{max}} = 25.5^\circ$, $\theta_{\text{min}} = 2.7^\circ$
$T_{\text{min}} = 0.976$, $T_{\text{max}} = 0.979$	$h = -10 \rightarrow 11$
5880 measured reflections	$k = -11 \rightarrow 14$
	$l = -11 \rightarrow 11$

Refinement

Refinement on F^2	Secondary atom site location: difference Fourier map
Least-squares matrix: full	Hydrogen site location: inferred from neighbouring sites
$R[F^2 > 2\sigma(F^2)] = 0.040$	H-atom parameters constrained
$wR(F^2) = 0.116$	$w = 1/[\sigma^2(F_o^2) + (0.0422P)^2 + 0.3813P]$
$S = 1.09$	where $P = (F_o^2 + 2F_c^2)/3$
1935 reflections	$(\Delta/\sigma)_{\text{max}} < 0.001$
147 parameters	$\Delta\rho_{\text{max}} = 0.18 \text{ e \AA}^{-3}$
0 restraints	$\Delta\rho_{\text{min}} = -0.19 \text{ e \AA}^{-3}$
Primary atom site location: structure-invariant direct methods	Extinction correction: <i>SHELXL97</i> (Sheldrick, 2008), $F_c^* = kF_c[1 + 0.001xF_c^2\lambda^3/\sin(2\theta)]^{1/4}$
	Extinction coefficient: 0.077 (5)

Special details

Geometry. All e.s.d.'s (except the e.s.d. in the dihedral angle between two l.s. planes) are estimated using the full covariance matrix. The cell e.s.d.'s are taken into account individually in the estimation of e.s.d.'s in distances, angles and torsion angles; correlations between e.s.d.'s in cell parameters are only used when they are defined by crystal symmetry. An approximate (isotropic) treatment of cell e.s.d.'s is used for estimating e.s.d.'s involving l.s. planes.

Refinement. Refinement of F^2 against ALL reflections. The weighted R -factor wR and goodness of fit S are based on F^2 , conventional R -factors R are based on F , with F set to zero for negative F^2 . The threshold expression of $F^2 > \sigma(F^2)$ is used only for calculating R -factors(gt) etc. and is not relevant to the choice of reflections for refinement. R -factors based on F^2 are statistically about twice as large as those based on F , and R -factors based on ALL data will be even larger.

Fractional atomic coordinates and isotropic or equivalent isotropic displacement parameters (\AA^2)

	x	y	z	$U_{\text{iso}}^*/U_{\text{eq}}$
C1	0.3906 (2)	0.14260 (14)	0.6005 (2)	0.0440 (5)
C2	0.4887 (2)	0.22747 (14)	0.63956 (18)	0.0409 (4)
H2	0.5876	0.2139	0.6826	0.049*
C3	0.43732 (19)	0.33301 (14)	0.61347 (18)	0.0368 (4)
H3	0.5019	0.3915	0.6400	0.044*
C4	0.28916 (18)	0.35297 (12)	0.54763 (17)	0.0327 (4)
C5	0.1950 (2)	0.26412 (14)	0.50840 (18)	0.0383 (4)
H5	0.0962	0.2765	0.4636	0.046*
C6	0.2447 (2)	0.15804 (14)	0.5344 (2)	0.0451 (5)
H6	0.1812	0.0989	0.5079	0.054*
C7	0.22867 (19)	0.46193 (13)	0.50988 (18)	0.0355 (4)
H7	0.1338	0.4662	0.4497	0.043*
C8	0.22355 (17)	0.65015 (12)	0.48809 (17)	0.0317 (4)
C9	0.22777 (18)	0.74242 (13)	0.57483 (17)	0.0343 (4)
H9	0.2750	0.7392	0.6739	0.041*
C10	0.16262 (19)	0.83889 (13)	0.51566 (18)	0.0373 (4)
H10	0.1647	0.8998	0.5752	0.045*
C11	0.09399 (18)	0.84530 (13)	0.36764 (18)	0.0354 (4)
C12	0.09253 (18)	0.75496 (14)	0.27921 (18)	0.0363 (4)
H12	0.0489	0.7595	0.1794	0.044*
C13	0.15602 (19)	0.65777 (13)	0.33919 (18)	0.0362 (4)
H13	0.1536	0.5969	0.2794	0.043*
N1	0.29403 (15)	0.55196 (11)	0.55190 (14)	0.0344 (4)
F1	0.44178 (15)	0.03938 (9)	0.62945 (16)	0.0739 (4)
O1	0.03080 (16)	0.94277 (10)	0.31471 (14)	0.0520 (4)
H1A	-0.0279	0.9327	0.2353	0.078*

Atomic displacement parameters (\AA^2)

	U^{11}	U^{22}	U^{33}	U^{12}	U^{13}	U^{23}
C1	0.0595 (12)	0.0274 (9)	0.0470 (10)	0.0055 (8)	0.0167 (9)	0.0046 (7)
C2	0.0407 (10)	0.0414 (10)	0.0392 (9)	0.0064 (8)	0.0077 (8)	0.0005 (7)

supplementary materials

C3	0.0408 (10)	0.0324 (9)	0.0371 (9)	-0.0032 (7)	0.0096 (7)	-0.0048 (7)
C4	0.0393 (9)	0.0291 (9)	0.0298 (8)	-0.0013 (7)	0.0089 (7)	0.0004 (6)
C5	0.0402 (10)	0.0359 (10)	0.0371 (9)	-0.0045 (7)	0.0064 (7)	0.0029 (7)
C6	0.0550 (12)	0.0302 (10)	0.0494 (10)	-0.0093 (8)	0.0117 (9)	-0.0002 (8)
C7	0.0350 (9)	0.0327 (9)	0.0371 (9)	-0.0010 (7)	0.0061 (7)	0.0017 (7)
C8	0.0314 (9)	0.0279 (9)	0.0346 (8)	-0.0006 (6)	0.0065 (7)	0.0036 (6)
C9	0.0371 (9)	0.0333 (9)	0.0295 (8)	-0.0027 (7)	0.0031 (7)	0.0009 (7)
C10	0.0427 (10)	0.0278 (9)	0.0390 (9)	-0.0022 (7)	0.0060 (7)	-0.0026 (7)
C11	0.0352 (9)	0.0279 (9)	0.0405 (9)	-0.0013 (7)	0.0049 (7)	0.0058 (7)
C12	0.0394 (10)	0.0374 (10)	0.0292 (8)	-0.0013 (7)	0.0033 (7)	0.0025 (7)
C13	0.0410 (10)	0.0318 (9)	0.0344 (9)	-0.0011 (7)	0.0070 (7)	-0.0035 (7)
N1	0.0380 (8)	0.0291 (8)	0.0345 (7)	0.0009 (6)	0.0064 (6)	0.0020 (6)
F1	0.0840 (10)	0.0317 (7)	0.1022 (10)	0.0132 (6)	0.0167 (8)	0.0105 (6)
O1	0.0641 (9)	0.0294 (7)	0.0508 (8)	0.0039 (6)	-0.0071 (6)	0.0060 (5)

Geometric parameters (\AA , $^\circ$)

C1—F1	1.352 (2)	C8—C9	1.388 (2)
C1—C6	1.368 (3)	C8—C13	1.392 (2)
C1—C2	1.374 (3)	C8—N1	1.428 (2)
C2—C3	1.377 (2)	C9—C10	1.379 (2)
C2—H2	0.9300	C9—H9	0.9300
C3—C4	1.395 (2)	C10—C11	1.386 (2)
C3—H3	0.9300	C10—H10	0.9300
C4—C5	1.390 (2)	C11—O1	1.367 (2)
C4—C7	1.456 (2)	C11—C12	1.382 (2)
C5—C6	1.378 (2)	C12—C13	1.384 (2)
C5—H5	0.9300	C12—H12	0.9300
C6—H6	0.9300	C13—H13	0.9300
C7—N1	1.273 (2)	O1—H1A	0.8200
C7—H7	0.9300		
F1—C1—C6	118.99 (17)	C9—C8—C13	118.68 (14)
F1—C1—C2	117.95 (17)	C9—C8—N1	119.45 (14)
C6—C1—C2	123.06 (16)	C13—C8—N1	121.80 (14)
C1—C2—C3	118.47 (17)	C10—C9—C8	120.71 (15)
C1—C2—H2	120.8	C10—C9—H9	119.6
C3—C2—H2	120.8	C8—C9—H9	119.6
C2—C3—C4	120.59 (16)	C9—C10—C11	120.18 (15)
C2—C3—H3	119.7	C9—C10—H10	119.9
C4—C3—H3	119.7	C11—C10—H10	119.9
C5—C4—C3	118.60 (15)	O1—C11—C12	122.48 (15)
C5—C4—C7	117.74 (15)	O1—C11—C10	117.79 (15)
C3—C4—C7	123.56 (15)	C12—C11—C10	119.73 (15)
C6—C5—C4	121.46 (17)	C13—C12—C11	119.99 (15)
C6—C5—H5	119.3	C13—C12—H12	120.0
C4—C5—H5	119.3	C11—C12—H12	120.0
C1—C6—C5	117.80 (16)	C12—C13—C8	120.68 (15)
C1—C6—H6	121.1	C12—C13—H13	119.7
C5—C6—H6	121.1	C8—C13—H13	119.7

N1—C7—C4	125.84 (15)	C7—N1—C8	117.26 (14)
N1—C7—H7	117.1	C11—O1—H1A	109.5
C4—C7—H7	117.1		
F1—C1—C2—C3	−178.66 (16)	N1—C8—C9—C10	178.98 (15)
C6—C1—C2—C3	1.3 (3)	C8—C9—C10—C11	−1.1 (3)
C1—C2—C3—C4	−0.6 (3)	C9—C10—C11—O1	179.82 (16)
C2—C3—C4—C5	−0.3 (2)	C9—C10—C11—C12	−0.7 (3)
C2—C3—C4—C7	−176.69 (15)	O1—C11—C12—C13	−178.88 (16)
C3—C4—C5—C6	0.6 (3)	C10—C11—C12—C13	1.7 (3)
C7—C4—C5—C6	177.20 (16)	C11—C12—C13—C8	−0.9 (3)
F1—C1—C6—C5	178.95 (16)	C9—C8—C13—C12	−0.9 (3)
C2—C1—C6—C5	−1.0 (3)	N1—C8—C13—C12	−177.93 (16)
C4—C5—C6—C1	0.0 (3)	C4—C7—N1—C8	172.19 (15)
C5—C4—C7—N1	171.35 (17)	C9—C8—N1—C7	140.76 (16)
C3—C4—C7—N1	−12.2 (3)	C13—C8—N1—C7	−42.2 (2)
C13—C8—C9—C10	1.9 (3)		

Hydrogen-bond geometry (Å, °)

D—H···A	D—H	H···A	D···A	D—H···A
O1—H1A···N1 ⁱ	0.82	2.09	2.885 (2)	163

Symmetry codes: (i) $x-1/2, -y+3/2, z-1/2$.

supplementary materials

Fig. 1




Fig. 2

