

Ethyl 5-methyl-1*H*-pyrrole-2-carboxylate

Zhao-Po Zhang,^a Yuan Wang^{a*} and Xiao-Xia Li^b

^aDepartment of Physics and Chemistry, Henan Polytechnic University, Jiaozuo 454000, People's Republic of China, and ^bInstitute of Functional Materials, Jiangxi University of Finance and Economics, Nanchang 330013, People's Republic of China
Correspondence e-mail: wangyuan08@hpu.edu.cn


Received 1 June 2011; accepted 1 June 2011

Key indicators: single-crystal X-ray study; $T = 296\text{ K}$; mean $\sigma(\text{C}-\text{C}) = 0.003\text{ \AA}$; R factor = 0.045; wR factor = 0.143; data-to-parameter ratio = 19.5.

In the title molecule, $\text{C}_8\text{H}_{11}\text{NO}_2$, the r.m.s. deviation of non-H atoms from their best plane is 0.031 \AA . Molecules are connected via a pair of $\text{N}-\text{H}\cdots\text{O}$ hydrogen bonds into a centrosymmetric dimer.

Related literature

For the crystal structure of the 5-cyano analogue of the title compound, see: Zhang *et al.* (2003). For the synthesis of the title compound, see: Motekaitis *et al.* (1970).



Experimental

Crystal data

$\text{C}_8\text{H}_{11}\text{NO}_2$
 $M_r = 153.18$
Monoclinic, $P2_1/c$
 $a = 7.0759 (3)\text{ \AA}$
 $b = 18.0705 (9)\text{ \AA}$
 $c = 6.6606 (3)\text{ \AA}$
 $\beta = 101.349 (3)^\circ$
 $V = 835.01 (7)\text{ \AA}^3$

$Z = 4$
Mo $K\alpha$ radiation
 $\mu = 0.09\text{ mm}^{-1}$

$T = 296\text{ K}$
 $0.20 \times 0.15 \times 0.13\text{ mm}$

Data collection

Bruker APEXII CCD diffractometer
Absorption correction: multi-scan (*SADABS*; Bruker, 2007)
 $T_{\min} = 0.984$, $T_{\max} = 0.989$

7496 measured reflections
1991 independent reflections
1157 reflections with $I > 2\sigma(I)$
 $R_{\text{int}} = 0.029$

Refinement

$R[F^2 > 2\sigma(F^2)] = 0.045$
 $wR(F^2) = 0.143$
 $S = 1.02$
1991 reflections

102 parameters
H-atom parameters constrained
 $\Delta\rho_{\max} = 0.16\text{ e \AA}^{-3}$
 $\Delta\rho_{\min} = -0.16\text{ e \AA}^{-3}$

Table 1
Hydrogen-bond geometry (\AA , $^\circ$).

$D-\text{H}\cdots A$	$D-\text{H}$	$\text{H}\cdots A$	$D\cdots A$	$D-\text{H}\cdots A$
N1—H1 \cdots O1 ⁱ	0.86	1.99	2.8363 (19)	167

Symmetry code: (i) $-x + 1, -y + 1, -z + 2$.

Data collection: *APEX2* (Bruker, 2007); cell refinement: *SAINT* (Bruker, 2007); data reduction: *SAINT*; program(s) used to solve structure: *SHELXS97* (Sheldrick, 2008); program(s) used to refine structure: *SHELXL97* (Sheldrick, 2008); molecular graphics: *SHELXTL* (Sheldrick, 2008); software used to prepare material for publication: *SHELXTL*.

The authors are grateful to the National Natural Science Foundation of China for financial support (grant No. 21001040).

Supplementary data and figures for this paper are available from the IUCr electronic archives (Reference: GK2381).

References

- Bruker (2007). *APEX2, SAINT* and *SADABS*. Bruker AXS Inc., Madison, Wisconsin, USA.
- Motekaitis, R. J., Heinert, H. D. & Martell, A. E. (1970). *J. Org. Chem.* **35**, 2504–2511.
- Sheldrick, G. M. (2008). *Acta Cryst. A* **64**, 112–122.
- Zhang, Y.-H., Yin, Z., Li, X.-F., He, J. & Cheng, J.-P. (2003). *Acta Cryst. E* **59**, o1881–o1882.

supplementary materials

Acta Cryst. (2011). E67, o1585 [doi:10.1107/S1600536811021180]

Ethyl 5-methyl-1*H*-pyrrole-2-carboxylate

Z.-P. Zhang, Y. Wang and X.-X. Li

Comment

As part of our studies on hydrazone ligands bearing pyrrole unit the title compound was synthesized and characterized by X-ray diffraction.

The non-hydrogen atoms of the title molecule (Fig. 1) are situated in a fair plane (r.m.s. deviation of the non-hydrogen atoms being 0.0306 Å). In the crystal, the molecules are linked into a centrosymmetric dimer by two intermolecular N—H···O hydrogen bonds, forming a $R_2^2(10)$ ring motif (Table 1, Fig. 2).

Experimental

The title compound was synthesized according to the literature procedure (Motekaitis *et al.*, 1970). The crude orange solid was washed well with ice water and dried in air. Colorless plates of the title compound were obtained by slow evaporation of petroleum ether/ethyl acetate (100:1) solution.

Refinement

All H atoms were placed in calculated positions, with C—H = 0.93–0.97 Å and N—H = 0.86 Å, and were thereafter treated as riding, with $U_{\text{iso}}(\text{H})$ values of 1.5Ueq(C) for methyl groups and 1.2Ueq(C,N) for others.

Figures




Fig. 1. The title compound with the displacement ellipsoids shown at the 50% probability level.




Fig. 2. The centrosymmetric dimer *via* N—H···O hydrogen bonds of the title compound (hydrogen bonds shown as dashed lines, symmetry code:: 1 - x , 1 - y , 2 - z).

Ethyl 5-methyl-1*H*-pyrrole-2-carboxylate

Crystal data

C₈H₁₁NO₂

$F(000) = 328$

$M_r = 153.18$

$D_x = 1.218 \text{ Mg m}^{-3}$

Monoclinic, $P2_1/c$

Mo $K\alpha$ radiation, $\lambda = 0.71073 \text{ \AA}$

supplementary materials

Hall symbol: -P 2ybc	Cell parameters from 2070 reflections
$a = 7.0759 (3) \text{ \AA}$	$\theta = 2.3\text{--}24.2^\circ$
$b = 18.0705 (9) \text{ \AA}$	$\mu = 0.09 \text{ mm}^{-1}$
$c = 6.6606 (3) \text{ \AA}$	$T = 296 \text{ K}$
$\beta = 101.349 (3)^\circ$	Plate, colorless
$V = 835.01 (7) \text{ \AA}^3$	$0.20 \times 0.15 \times 0.13 \text{ mm}$
$Z = 4$	

Data collection

Bruker APEXII CCD diffractometer	1991 independent reflections
Radiation source: fine-focus sealed tube graphite	1157 reflections with $I > 2\sigma(I)$
φ and ω scans	$R_{\text{int}} = 0.029$
Absorption correction: multi-scan (<i>SADABS</i> ; Bruker, 2007)	$\theta_{\text{max}} = 27.9^\circ, \theta_{\text{min}} = 2.3^\circ$
$T_{\text{min}} = 0.984, T_{\text{max}} = 0.989$	$h = -9 \rightarrow 9$
7496 measured reflections	$k = -23 \rightarrow 23$
	$l = -8 \rightarrow 8$

Refinement

Refinement on F^2	Primary atom site location: structure-invariant direct methods
Least-squares matrix: full	Secondary atom site location: difference Fourier map
$R[F^2 > 2\sigma(F^2)] = 0.045$	Hydrogen site location: inferred from neighbouring sites
$wR(F^2) = 0.143$	H-atom parameters constrained
$S = 1.02$	$w = 1/[\sigma^2(F_o^2) + (0.0683P)^2 + 0.0738P]$ where $P = (F_o^2 + 2F_c^2)/3$
1991 reflections	$(\Delta/\sigma)_{\text{max}} < 0.001$
102 parameters	$\Delta\rho_{\text{max}} = 0.16 \text{ e \AA}^{-3}$
0 restraints	$\Delta\rho_{\text{min}} = -0.16 \text{ e \AA}^{-3}$

Special details

Geometry. All e.s.d.'s (except the e.s.d. in the dihedral angle between two l.s. planes) are estimated using the full covariance matrix. The cell e.s.d.'s are taken into account individually in the estimation of e.s.d.'s in distances, angles and torsion angles; correlations between e.s.d.'s in cell parameters are only used when they are defined by crystal symmetry. An approximate (isotropic) treatment of cell e.s.d.'s is used for estimating e.s.d.'s involving l.s. planes.

Refinement. Refinement of F^2 against ALL reflections. The weighted R -factor wR and goodness of fit S are based on F^2 , conventional R -factors R are based on F , with F set to zero for negative F^2 . The threshold expression of $F^2 > \sigma(F^2)$ is used only for calculating R -factors(gt) etc. and is not relevant to the choice of reflections for refinement. R -factors based on F^2 are statistically about twice as large as those based on F , and R -factors based on ALL data will be even larger.

Fractional atomic coordinates and isotropic or equivalent isotropic displacement parameters (\AA^2)

	<i>x</i>	<i>y</i>	<i>z</i>	$U_{\text{iso}}^*/U_{\text{eq}}$
O2	0.12507 (16)	0.64959 (6)	0.92921 (18)	0.0629 (4)
N1	0.52408 (19)	0.57393 (7)	0.7402 (2)	0.0581 (4)
H1	0.5637	0.5360	0.8146	0.087*
C6	0.2631 (2)	0.60036 (9)	0.9187 (3)	0.0585 (4)
O1	0.2888 (2)	0.54602 (8)	1.0272 (2)	0.0861 (5)
C5	0.3729 (2)	0.61798 (9)	0.7644 (2)	0.0548 (4)
C2	0.6023 (2)	0.59835 (10)	0.5831 (3)	0.0600 (5)
C8	-0.1423 (3)	0.69326 (11)	1.0591 (3)	0.0768 (6)
H8A	-0.0794	0.7400	1.0916	0.115*
H8B	-0.2268	0.6837	1.1527	0.115*
H8C	-0.2159	0.6946	0.9217	0.115*
C4	0.3561 (2)	0.67196 (10)	0.6178 (3)	0.0651 (5)
H4	0.2658	0.7100	0.5976	0.078*
C7	0.0051 (3)	0.63353 (10)	1.0768 (3)	0.0691 (5)
H7A	0.0825	0.6324	1.2142	0.083*
H7B	-0.0569	0.5858	1.0480	0.083*
C1	0.7719 (3)	0.56133 (11)	0.5261 (3)	0.0780 (6)
H1A	0.8862	0.5893	0.5791	0.117*
H1B	0.7543	0.5585	0.3796	0.117*
H1C	0.7847	0.5123	0.5827	0.117*
C3	0.4987 (3)	0.65956 (10)	0.5044 (3)	0.0712 (5)
H3	0.5200	0.6879	0.3944	0.085*

Atomic displacement parameters (\AA^2)

	U^{11}	U^{22}	U^{33}	U^{12}	U^{13}	U^{23}
O2	0.0581 (7)	0.0624 (7)	0.0748 (8)	0.0073 (5)	0.0294 (6)	0.0035 (6)
N1	0.0533 (8)	0.0560 (8)	0.0691 (9)	0.0001 (6)	0.0221 (7)	0.0000 (6)
C6	0.0519 (9)	0.0563 (9)	0.0698 (11)	0.0019 (8)	0.0182 (8)	-0.0026 (9)
O1	0.0847 (10)	0.0820 (9)	0.1032 (11)	0.0273 (8)	0.0467 (8)	0.0332 (8)
C5	0.0486 (9)	0.0532 (9)	0.0649 (11)	-0.0015 (7)	0.0165 (8)	-0.0035 (8)
C2	0.0539 (9)	0.0652 (10)	0.0644 (11)	-0.0091 (8)	0.0202 (8)	-0.0063 (8)
C8	0.0689 (12)	0.0819 (13)	0.0884 (15)	0.0074 (10)	0.0373 (11)	-0.0054 (10)
C4	0.0587 (10)	0.0616 (10)	0.0766 (12)	0.0036 (8)	0.0172 (9)	0.0060 (9)
C7	0.0635 (11)	0.0788 (12)	0.0716 (12)	0.0050 (9)	0.0293 (9)	0.0048 (9)
C1	0.0664 (12)	0.0894 (14)	0.0867 (14)	-0.0022 (10)	0.0360 (10)	-0.0094 (11)
C3	0.0683 (11)	0.0781 (12)	0.0715 (12)	-0.0023 (10)	0.0246 (10)	0.0107 (10)

Geometric parameters (\AA , $^\circ$)

C2—C1	1.487 (2)	C1—H1A	0.9600
C2—C3	1.373 (3)	C1—H1B	0.9600
C4—C3	1.392 (2)	C1—H1C	0.9600
C5—C4	1.369 (2)	C3—H3	0.9300
N1—C2	1.351 (2)	C4—H4	0.9300

supplementary materials


N1—C5	1.368 (2)	C7—H7A	0.9700
C6—C5	1.440 (2)	C7—H7B	0.9700
C6—O1	1.212 (2)	C8—H8A	0.9600
O2—C6	1.3333 (19)	C8—H8B	0.9600
O2—C7	1.4490 (19)	C8—H8C	0.9600
C8—C7	1.490 (2)	N1—H1	0.8600
C6—O2—C7	115.81 (13)	C5—C4—C3	107.57 (16)
C2—N1—C5	110.53 (14)	C5—C4—H4	126.2
C2—N1—H1	124.7	C3—C4—H4	126.2
C5—N1—H1	124.7	O2—C7—C8	107.22 (15)
O1—C6—O2	122.31 (15)	O2—C7—H7A	110.3
O1—C6—C5	124.50 (15)	C8—C7—H7A	110.3
O2—C6—C5	113.19 (15)	O2—C7—H7B	110.3
C4—C5—N1	106.89 (14)	C8—C7—H7B	110.3
C4—C5—C6	133.09 (16)	H7A—C7—H7B	108.5
N1—C5—C6	119.97 (15)	C2—C1—H1A	109.5
N1—C2—C3	106.81 (15)	C2—C1—H1B	109.5
N1—C2—C1	121.69 (16)	H1A—C1—H1B	109.5
C3—C2—C1	131.48 (17)	C2—C1—H1C	109.5
C7—C8—H8A	109.5	H1A—C1—H1C	109.5
C7—C8—H8B	109.5	H1B—C1—H1C	109.5
H8A—C8—H8B	109.5	C2—C3—C4	108.20 (16)
C7—C8—H8C	109.5	C2—C3—H3	125.9
H8A—C8—H8C	109.5	C4—C3—H3	125.9
H8B—C8—H8C	109.5		
C7—O2—C6—O1	-0.9 (2)	C5—N1—C2—C3	-0.1 (2)
C7—O2—C6—C5	177.93 (14)	C5—N1—C2—C1	178.73 (15)
C2—N1—C5—C4	-0.12 (19)	N1—C5—C4—C3	0.26 (19)
C2—N1—C5—C6	177.58 (14)	C6—C5—C4—C3	-177.02 (18)
O1—C6—C5—C4	173.57 (18)	C6—O2—C7—C8	-178.10 (14)
O2—C6—C5—C4	-5.3 (3)	N1—C2—C3—C4	0.2 (2)
O1—C6—C5—N1	-3.4 (3)	C1—C2—C3—C4	-178.40 (18)
O2—C6—C5—N1	177.75 (13)	C5—C4—C3—C2	-0.3 (2)

Hydrogen-bond geometry (Å, °)

D—H···A	D—H	H···A	D···A	D—H···A
N1—H1···O1 ⁱ	0.86	1.99	2.8363 (19)	167.

Symmetry codes: (i) $-x+1, -y+1, -z+2$.

Fig. 1



supplementary materials

Fig. 2

