

## N'-(4-Diethylamino-2-hydroxybenzylidene)-4-(dimethylamino)benzo-hydrazide methanol monosolvate

Fu Su, Zheng-Gui Gu and Jun Lin\*

Jiangsu Centre of Extraction Separation Engineering Technology, College of Chemistry and Materials Science, Nanjing Normal University, Nanjing 210046, People's Republic of China  
Correspondence e-mail: douduo2011@163.com


Received 25 May 2011; accepted 25 May 2011

Key indicators: single-crystal X-ray study;  $T = 298\text{ K}$ ; mean  $\sigma(\text{C}-\text{C}) = 0.005\text{ \AA}$ ;  $R$  factor = 0.070;  $wR$  factor = 0.189; data-to-parameter ratio = 16.8.

The title compound,  $\text{C}_{20}\text{H}_{26}\text{N}_4\text{O}_2\cdot\text{CH}_3\text{OH}$ , was prepared by the reaction of 4-diethylamino-2-hydroxybenzaldehyde with 4-(dimethylamino)benzohydrazide. The dihedral angle between the two benzene rings is  $13.6(3)^\circ$  and an intramolecular  $\text{O}-\text{H}\cdots\text{N}$  hydrogen bond generates an  $S(6)$  ring. In the crystal, the hydrazone and methanol molecules are linked through intermolecular  $\text{O}-\text{H}\cdots\text{O}$  and  $\text{N}-\text{H}\cdots\text{O}$  hydrogen bonds, forming chains along  $a$ .

### Related literature

For the biological properties of hydrazones, see: Ajani *et al.* (2010); Zhang *et al.* (2010); Angelusiu *et al.* (2010). For similar structures, see: Huang & Wu (2010); Khaledi *et al.* (2010); Zhou & Yang (2010); Ji & Lu (2010); Singh & Singh (2010); Ahmad *et al.* (2010). For hydrogen-bond motifs, see Bernstein *et al.* (1995).



### Experimental

#### Crystal data

$\text{C}_{20}\text{H}_{26}\text{N}_4\text{O}_2\cdot\text{CH}_3\text{OH}$   
 $M_r = 386.49$   
Triclinic,  $P\bar{1}$

$a = 6.786(3)\text{ \AA}$   
 $b = 11.791(3)\text{ \AA}$   
 $c = 14.252(2)\text{ \AA}$

$\alpha = 111.511(3)^\circ$   
 $\beta = 92.811(2)^\circ$   
 $\gamma = 96.492(2)^\circ$   
 $V = 1049.1(6)\text{ \AA}^3$   
 $Z = 2$

Mo  $K\alpha$  radiation  
 $\mu = 0.08\text{ mm}^{-1}$   
 $T = 298\text{ K}$   
 $0.27 \times 0.23 \times 0.21\text{ mm}$

#### Data collection

Bruker SMART CCD area-detector diffractometer  
Absorption correction: multi-scan (*SADABS*; Bruker, 2001)  
 $T_{\min} = 0.978$ ,  $T_{\max} = 0.983$

7543 measured reflections  
4421 independent reflections  
1809 reflections with  $I > 2\sigma(I)$   
 $R_{\text{int}} = 0.045$

#### Refinement

$R[F^2 > 2\sigma(F^2)] = 0.070$   
 $wR(F^2) = 0.189$   
 $S = 1.00$   
4421 reflections  
263 parameters  
1 restraint

H atoms treated by a mixture of independent and constrained refinement  
 $\Delta\rho_{\max} = 0.30\text{ e \AA}^{-3}$   
 $\Delta\rho_{\min} = -0.21\text{ e \AA}^{-3}$

**Table 1**  
Hydrogen-bond geometry ( $\text{\AA}$ ,  $^\circ$ ).

$D-\text{H}\cdots A$	$D-\text{H}$	$\text{H}\cdots A$	$D\cdots A$	$D-\text{H}\cdots A$
$\text{N}2-\text{H2}\cdots\text{O}3$	0.90 (1)	2.07 (2)	2.936 (4)	160 (3)
$\text{O}3-\text{H3}\cdots\text{O}2^i$	0.82	1.84	2.661 (3)	177
$\text{O}1-\text{H1}\cdots\text{N}1$	0.82	2.02	2.727 (3)	145

Symmetry code: (i)  $x + 1, y, z$ .

Data collection: *SMART* (Bruker, 2007); cell refinement: *SAINT* (Bruker, 2007); data reduction: *SAINT*; program(s) used to solve structure: *SHELXTL* (Sheldrick, 2008); program(s) used to refine structure: *SHELXTL*; molecular graphics: *SHELXTL*; software used to prepare material for publication: *SHELXTL*.

We acknowledge the project funded by the Priority Academic Program Development of Jiangsu Higher Education Institutions.

Supplementary data and figures for this paper are available from the IUCr electronic archives (Reference: SJ5154).

### References

- Ahmad, T., Zia-ur-Rehman, M., Siddiqui, H. L., Mahmud, S. & Parvez, M. (2010). *Acta Cryst. E66*, o1022.  
Ajani, O. O., Obafemi, C. A., Nwinyi, O. C. & Akinpelu, D. A. (2010). *Bioorg. Med. Chem.* **18**, 214–221.  
Angelusiu, M. V., Barbuceanu, S. F., Draghici, C. & Almajan, G. L. (2010). *Eur. J. Med. Chem.* **45**, 2055–2062.  
Bernstein, J., Davis, R. E., Shimoni, L. & Chang, N.-L. (1995). *Angew. Chem. Int. Ed. Engl.* **34**, 1555–1573.  
Bruker (2001). *SADABS*. Bruker AXS Inc., Madison, Wisconsin, USA.  
Bruker (2007). *SMART* and *SAINT*. Bruker AXS Inc., Madison, Wisconsin, USA.  
Huang, H.-T. & Wu, H.-Y. (2010). *Acta Cryst. E66*, o2729–o2730.  
Ji, X.-H. & Lu, J.-F. (2010). *Acta Cryst. E66*, o1514.  
Khaledi, H., Alhadi, A. A., Mohd Ali, H., Robinson, W. T. & Abdulla, M. A. (2010). *Acta Cryst. E66*, o105–o106.  
Sheldrick, G. M. (2008). *Acta Cryst. A64*, 112–122.  
Singh, V. P. & Singh, S. (2010). *Acta Cryst. E66*, o1172.  
Zhang, Y.-H., Zhang, L., Liu, L., Guo, J.-X., Wu, D.-L., Xu, G.-C., Wang, X.-H. & Jia, D.-Z. (2010). *Inorg. Chim. Acta*, **363**, 289–293.  
Zhou, C.-S. & Yang, T. (2010). *Acta Cryst. E66*, o290.

## **supplementary materials**

*Acta Cryst.* (2011). E67, o1634 [doi:10.1107/S1600536811019957]

## ***N'*-(4-Diethylamino-2-hydroxybenzylidene)-4-(dimethylamino)benzohydrazide methanol monosolvate**

**F. Su, Z.-G. Gu and J. Lin**

### **Comment**

In the last year, much attention has been focused on the biological properties of hydrazone compounds (Ajani *et al.*, 2010; Zhang *et al.*, 2010; Angelusiu *et al.*, 2010). The crystal structures of a number of hydrazone compounds have also been determined (Huang & Wu, 2010; Khaledi *et al.*, 2010; Zhou & Yang, 2010; Ji & Lu, 2010; Singh & Singh, 2010; Ahmad *et al.*, 2010). In this paper, the title new hydrazone compound, Fig. 1, is reported.

The asymmetric unit of the compound contains a hydrazone molecule and a methanol molecule. The dihedral angle between the two benzene rings is 13.6 (3) $^{\circ}$  and an intramolecular O1—H1 $\cdots$ N1 hydrogen bond forms an S(6) ring (Bernstein *et al.*, 1995). In the crystal structure, the hydrazone and methanol molecules are linked through intermolecular O—H $\cdots$ O and N—H $\cdots$ O hydrogen bonds (Table 1), to form 1D chains along  $a$  (Fig. 2).

### **Experimental**

The reaction of 4-diethylamino-2-hydroxybenzaldehyde (0.193 g, 1 mmol) with 4-(dimethylamino)benzohydrazide (0.179 g, 1 mmol) in 30 ml methanol at room temperature afforded the title compound. Colorless single crystals were formed by gradual evaporation of the solution in air.

### **Refinement**

The amino H atom was located in a difference Fourier map and refined with the N—H distance restrained to be 0.90 (1) Å, and with  $U_{\text{iso}} = 0.08 \text{ \AA}^2$ . The remaining H atoms were positioned geometrically (C—H = 0.93–0.97 Å, O—H = 0.82 Å) and refined as riding with  $U_{\text{iso}}(\text{H}) = 1.2U_{\text{eq}}(\text{C})$  or 1.5  $U_{\text{eq}}(\text{O3 and C}_\text{methyl})$ .

### **Figures**




Fig. 1. The molecular structure of the title compound showing 30% probability displacement ellipsoids and the atomic numbering. Hydrogen bonds are drawn as dashed lines.

## supplementary materials

---




Fig. 2. Crystal packing of the title compound, viewed down the *b* axis. Intermolecular interactions are drawn as dashed lines.

### ***N<sup>1</sup>-(4-Diethylamino-2-hydroxybenzylidene)-4-(dimethylamino)benzohydrazide methanol monosolvate***

#### *Crystal data*

C <sub>20</sub> H <sub>26</sub> N <sub>4</sub> O <sub>2</sub> ·CH <sub>4</sub> O	Z = 2
$M_r = 386.49$	$F(000) = 416$
Triclinic, $P\bar{1}$	$D_x = 1.224 \text{ Mg m}^{-3}$
Hall symbol: -P 1	Mo $K\alpha$ radiation, $\lambda = 0.71073 \text{ \AA}$
$a = 6.786 (3) \text{ \AA}$	Cell parameters from 765 reflections
$b = 11.791 (3) \text{ \AA}$	$\theta = 2.6\text{--}24.5^\circ$
$c = 14.252 (2) \text{ \AA}$	$\mu = 0.08 \text{ mm}^{-1}$
$\alpha = 111.511 (3)^\circ$	$T = 298 \text{ K}$
$\beta = 92.811 (2)^\circ$	Block, colorless
$\gamma = 96.492 (2)^\circ$	$0.27 \times 0.23 \times 0.21 \text{ mm}$
$V = 1049.1 (6) \text{ \AA}^3$	

#### *Data collection*

Bruker SMART CCD area-detector diffractometer	4421 independent reflections
Radiation source: fine-focus sealed tube graphite	1809 reflections with $I > 2\sigma(I)$
$\omega$ scans	$R_{\text{int}} = 0.045$
Absorption correction: multi-scan ( <i>SADABS</i> ; Bruker, 2001)	$\theta_{\text{max}} = 27.0^\circ, \theta_{\text{min}} = 2.8^\circ$
$T_{\text{min}} = 0.978, T_{\text{max}} = 0.983$	$h = -8 \rightarrow 8$
7543 measured reflections	$k = -15 \rightarrow 15$
	$l = -15 \rightarrow 18$

#### *Refinement*

Refinement on $F^2$	Primary atom site location: structure-invariant direct methods
Least-squares matrix: full	Secondary atom site location: difference Fourier map
$R[F^2 > 2\sigma(F^2)] = 0.070$	Hydrogen site location: inferred from neighbouring sites
$wR(F^2) = 0.189$	H atoms treated by a mixture of independent and constrained refinement

$S = 1.00$	$w = 1/[\sigma^2(F_o^2) + (0.0676P)^2]$
	where $P = (F_o^2 + 2F_c^2)/3$
4421 reflections	$(\Delta/\sigma)_{\max} < 0.001$
263 parameters	$\Delta\rho_{\max} = 0.30 \text{ e \AA}^{-3}$
1 restraint	$\Delta\rho_{\min} = -0.21 \text{ e \AA}^{-3}$

### Special details

**Geometry.** All esds (except the esd in the dihedral angle between two l.s. planes) are estimated using the full covariance matrix. The cell esds are taken into account individually in the estimation of esds in distances, angles and torsion angles; correlations between esds in cell parameters are only used when they are defined by crystal symmetry. An approximate (isotropic) treatment of cell esds is used for estimating esds involving l.s. planes.

**Refinement.** Refinement of  $F^2$  against ALL reflections. The weighted R-factor wR and goodness of fit S are based on  $F^2$ , conventional R-factors R are based on F, with F set to zero for negative  $F^2$ . The threshold expression of  $F^2 > 2\text{sigma}(F^2)$  is used only for calculating R-factors(gt) etc. and is not relevant to the choice of reflections for refinement. R-factors based on  $F^2$  are statistically about twice as large as those based on F, and R- factors based on ALL data will be even larger.

### Fractional atomic coordinates and isotropic or equivalent isotropic displacement parameters ( $\text{\AA}^2$ )

	<i>x</i>	<i>y</i>	<i>z</i>	$U_{\text{iso}}^*/U_{\text{eq}}$
O1	-0.1048 (3)	0.5092 (2)	0.2045 (2)	0.0741 (8)
H1	-0.0686	0.4564	0.2235	0.111*
O2	-0.0566 (3)	0.2247 (2)	0.3184 (2)	0.0770 (8)
O3	0.6751 (4)	0.3787 (2)	0.3427 (2)	0.0907 (9)
H3	0.7586	0.3321	0.3371	0.136*
N1	0.1680 (4)	0.3808 (2)	0.2520 (2)	0.0549 (7)
N2	0.2452 (4)	0.2985 (2)	0.2885 (2)	0.0564 (8)
N3	0.4202 (4)	-0.1790 (2)	0.3829 (2)	0.0632 (8)
N4	0.1131 (4)	0.7762 (2)	0.0366 (2)	0.0705 (9)
C1	0.2398 (4)	0.5230 (2)	0.1692 (2)	0.0488 (8)
C2	0.0478 (5)	0.5553 (3)	0.1644 (2)	0.0487 (8)
C3	0.0066 (4)	0.6387 (3)	0.1211 (2)	0.0535 (9)
H3A	-0.1218	0.6588	0.1194	0.064*
C4	0.1541 (5)	0.6933 (3)	0.0801 (2)	0.0525 (8)
C5	0.3450 (5)	0.6589 (3)	0.0828 (3)	0.0649 (10)
H5	0.4457	0.6922	0.0546	0.078*
C6	0.3848 (5)	0.5769 (3)	0.1263 (3)	0.0594 (9)
H6	0.5130	0.5564	0.1273	0.071*
C7	0.2927 (5)	0.4383 (3)	0.2145 (2)	0.0523 (8)
H7	0.4248	0.4246	0.2168	0.063*
C8	0.1245 (5)	0.2195 (3)	0.3155 (2)	0.0522 (8)
C9	0.2154 (4)	0.1219 (3)	0.3371 (2)	0.0470 (8)
C10	0.0978 (5)	0.0474 (3)	0.3749 (3)	0.0573 (9)
H10	-0.0301	0.0644	0.3897	0.069*
C11	0.1637 (5)	-0.0504 (3)	0.3910 (2)	0.0596 (9)
H11	0.0806	-0.0973	0.4173	0.071*
C12	0.3538 (5)	-0.0806 (3)	0.3687 (2)	0.0505 (8)

## supplementary materials

---

C13	0.4723 (4)	-0.0056 (3)	0.3301 (2)	0.0575 (9)
H13	0.6001	-0.0224	0.3148	0.069*
C14	0.4036 (5)	0.0927 (3)	0.3145 (2)	0.0553 (9)
H14	0.4855	0.1402	0.2883	0.066*
C15	0.2976 (6)	-0.2536 (3)	0.4256 (3)	0.0780 (11)
H15A	0.2808	-0.2051	0.4946	0.117*
H15B	0.3614	-0.3231	0.4233	0.117*
H15C	0.1696	-0.2821	0.3869	0.117*
C16	0.6191 (5)	-0.2071 (3)	0.3638 (3)	0.0789 (12)
H16A	0.6412	-0.2160	0.2956	0.118*
H16B	0.6347	-0.2824	0.3728	0.118*
H16C	0.7139	-0.1414	0.4103	0.118*
C17	0.2791 (6)	0.8609 (3)	0.0203 (3)	0.0792 (12)
H17A	0.2346	0.9396	0.0307	0.095*
H17B	0.3917	0.8751	0.0698	0.095*
C18	0.3427 (6)	0.8094 (4)	-0.0821 (3)	0.1027 (14)
H18A	0.3853	0.7310	-0.0929	0.154*
H18B	0.4511	0.8645	-0.0893	0.154*
H18C	0.2332	0.7989	-0.1312	0.154*
C19	-0.0856 (5)	0.8062 (3)	0.0254 (3)	0.0603 (9)
H19A	-0.1818	0.7332	0.0115	0.072*
H19B	-0.0975	0.8310	-0.0322	0.072*
C20	-0.1341 (6)	0.9075 (3)	0.1181 (3)	0.0841 (12)
H20A	-0.1327	0.8810	0.1742	0.126*
H20B	-0.2638	0.9272	0.1057	0.126*
H20C	-0.0366	0.9791	0.1336	0.126*
C21	0.7364 (6)	0.4885 (4)	0.4255 (3)	0.0977 (14)
H21A	0.8506	0.5328	0.4108	0.146*
H21B	0.7706	0.4704	0.4841	0.146*
H21C	0.6301	0.5378	0.4384	0.146*
H2	0.3791 (16)	0.309 (3)	0.290 (3)	0.080*

Atomic displacement parameters ( $\text{\AA}^2$ )

	$U^{11}$	$U^{22}$	$U^{33}$	$U^{12}$	$U^{13}$	$U^{23}$
O1	0.0555 (14)	0.0899 (18)	0.111 (2)	0.0186 (12)	0.0232 (14)	0.0730 (17)
O2	0.0539 (15)	0.0793 (16)	0.118 (2)	0.0240 (13)	0.0139 (14)	0.0555 (15)
O3	0.0596 (17)	0.0795 (18)	0.137 (3)	0.0214 (13)	-0.0033 (17)	0.0440 (18)
N1	0.0575 (18)	0.0537 (15)	0.062 (2)	0.0165 (13)	-0.0006 (15)	0.0301 (15)
N2	0.0473 (16)	0.0592 (16)	0.078 (2)	0.0149 (14)	0.0037 (16)	0.0420 (16)
N3	0.071 (2)	0.0561 (16)	0.077 (2)	0.0165 (15)	0.0121 (16)	0.0391 (16)
N4	0.0546 (18)	0.0831 (19)	0.108 (3)	0.0214 (15)	0.0239 (17)	0.070 (2)
C1	0.0493 (19)	0.0476 (17)	0.055 (2)	0.0121 (15)	0.0030 (16)	0.0239 (16)
C2	0.048 (2)	0.0527 (18)	0.051 (2)	0.0079 (15)	0.0085 (16)	0.0261 (17)
C3	0.0428 (19)	0.0593 (19)	0.071 (3)	0.0145 (16)	0.0099 (17)	0.0364 (18)
C4	0.050 (2)	0.0564 (18)	0.062 (2)	0.0109 (16)	0.0111 (17)	0.0329 (18)
C5	0.049 (2)	0.078 (2)	0.091 (3)	0.0170 (18)	0.0227 (19)	0.054 (2)
C6	0.047 (2)	0.064 (2)	0.078 (3)	0.0168 (16)	0.0093 (18)	0.036 (2)

C7	0.051 (2)	0.0497 (18)	0.058 (2)	0.0168 (16)	0.0002 (17)	0.0206 (17)
C8	0.046 (2)	0.0579 (19)	0.058 (2)	0.0172 (16)	0.0043 (17)	0.0248 (17)
C9	0.0447 (19)	0.0527 (18)	0.049 (2)	0.0133 (15)	0.0031 (16)	0.0233 (16)
C10	0.0446 (19)	0.071 (2)	0.070 (3)	0.0175 (16)	0.0158 (17)	0.038 (2)
C11	0.057 (2)	0.066 (2)	0.069 (3)	0.0114 (18)	0.0119 (19)	0.0401 (19)
C12	0.058 (2)	0.0480 (17)	0.050 (2)	0.0094 (16)	-0.0002 (17)	0.0233 (16)
C13	0.0447 (19)	0.061 (2)	0.075 (3)	0.0165 (16)	0.0112 (18)	0.0314 (19)
C14	0.050 (2)	0.0555 (18)	0.072 (3)	0.0133 (16)	0.0117 (18)	0.0353 (18)
C15	0.108 (3)	0.061 (2)	0.079 (3)	0.020 (2)	0.015 (2)	0.040 (2)
C16	0.071 (3)	0.069 (2)	0.109 (3)	0.0221 (19)	0.004 (2)	0.043 (2)
C17	0.079 (3)	0.083 (2)	0.094 (4)	0.025 (2)	0.017 (2)	0.050 (2)
C18	0.113 (4)	0.129 (4)	0.092 (4)	0.038 (3)	0.036 (3)	0.062 (3)
C19	0.058 (2)	0.066 (2)	0.069 (3)	0.0134 (17)	0.0033 (19)	0.038 (2)
C20	0.085 (3)	0.080 (3)	0.089 (3)	0.023 (2)	0.011 (2)	0.030 (2)
C21	0.101 (3)	0.090 (3)	0.114 (4)	0.029 (3)	0.024 (3)	0.046 (3)

*Geometric parameters (Å, °)*

O1—C2	1.362 (3)	C10—C11	1.373 (4)
O1—H1	0.8200	C10—H10	0.9300
O2—C8	1.239 (3)	C11—C12	1.397 (4)
O3—C21	1.397 (4)	C11—H11	0.9300
O3—H3	0.8200	C12—C13	1.404 (4)
N1—C7	1.281 (4)	C13—C14	1.383 (4)
N1—N2	1.396 (3)	C13—H13	0.9300
N2—C8	1.344 (4)	C14—H14	0.9300
N2—H2	0.900 (10)	C15—H15A	0.9600
N3—C12	1.368 (3)	C15—H15B	0.9600
N3—C16	1.441 (4)	C15—H15C	0.9600
N3—C15	1.454 (4)	C16—H16A	0.9600
N4—C4	1.381 (4)	C16—H16B	0.9600
N4—C19	1.448 (4)	C16—H16C	0.9600
N4—C17	1.504 (4)	C17—C18	1.469 (5)
C1—C6	1.396 (4)	C17—H17A	0.9700
C1—C2	1.403 (4)	C17—H17B	0.9700
C1—C7	1.441 (4)	C18—H18A	0.9600
C2—C3	1.384 (4)	C18—H18B	0.9600
C3—C4	1.395 (4)	C18—H18C	0.9600
C3—H3A	0.9300	C19—C20	1.501 (5)
C4—C5	1.404 (4)	C19—H19A	0.9700
C5—C6	1.369 (4)	C19—H19B	0.9700
C5—H5	0.9300	C20—H20A	0.9600
C6—H6	0.9300	C20—H20B	0.9600
C7—H7	0.9300	C20—H20C	0.9600
C8—C9	1.484 (4)	C21—H21A	0.9600
C9—C14	1.382 (4)	C21—H21B	0.9600
C9—C10	1.389 (4)	C21—H21C	0.9600
C2—O1—H1	109.5	C14—C13—C12	121.6 (3)
C21—O3—H3	109.5	C14—C13—H13	119.2

## supplementary materials

---

C7—N1—N2	115.4 (3)	C12—C13—H13	119.2
C8—N2—N1	121.0 (3)	C9—C14—C13	121.3 (3)
C8—N2—H2	129 (2)	C9—C14—H14	119.4
N1—N2—H2	110 (2)	C13—C14—H14	119.4
C12—N3—C16	121.6 (3)	N3—C15—H15A	109.5
C12—N3—C15	121.0 (3)	N3—C15—H15B	109.5
C16—N3—C15	117.3 (3)	H15A—C15—H15B	109.5
C4—N4—C19	122.5 (3)	N3—C15—H15C	109.5
C4—N4—C17	120.5 (3)	H15A—C15—H15C	109.5
C19—N4—C17	115.6 (2)	H15B—C15—H15C	109.5
C6—C1—C2	116.7 (3)	N3—C16—H16A	109.5
C6—C1—C7	119.4 (3)	N3—C16—H16B	109.5
C2—C1—C7	123.9 (3)	H16A—C16—H16B	109.5
O1—C2—C3	116.9 (3)	N3—C16—H16C	109.5
O1—C2—C1	121.8 (3)	H16A—C16—H16C	109.5
C3—C2—C1	121.3 (3)	H16B—C16—H16C	109.5
C2—C3—C4	121.2 (3)	C18—C17—N4	111.6 (3)
C2—C3—H3A	119.4	C18—C17—H17A	109.3
C4—C3—H3A	119.4	N4—C17—H17A	109.3
N4—C4—C3	121.3 (3)	C18—C17—H17B	109.3
N4—C4—C5	121.2 (3)	N4—C17—H17B	109.3
C3—C4—C5	117.5 (3)	H17A—C17—H17B	108.0
C6—C5—C4	120.9 (3)	C17—C18—H18A	109.5
C6—C5—H5	119.6	C17—C18—H18B	109.5
C4—C5—H5	119.6	H18A—C18—H18B	109.5
C5—C6—C1	122.4 (3)	C17—C18—H18C	109.5
C5—C6—H6	118.8	H18A—C18—H18C	109.5
C1—C6—H6	118.8	H18B—C18—H18C	109.5
N1—C7—C1	123.7 (3)	N4—C19—C20	112.6 (3)
N1—C7—H7	118.2	N4—C19—H19A	109.1
C1—C7—H7	118.2	C20—C19—H19A	109.1
O2—C8—N2	121.6 (3)	N4—C19—H19B	109.1
O2—C8—C9	121.1 (3)	C20—C19—H19B	109.1
N2—C8—C9	117.2 (3)	H19A—C19—H19B	107.8
C14—C9—C10	117.2 (3)	C19—C20—H20A	109.5
C14—C9—C8	124.7 (3)	C19—C20—H20B	109.5
C10—C9—C8	117.9 (3)	H20A—C20—H20B	109.5
C11—C10—C9	122.3 (3)	C19—C20—H20C	109.5
C11—C10—H10	118.8	H20A—C20—H20C	109.5
C9—C10—H10	118.8	H20B—C20—H20C	109.5
C10—C11—C12	121.0 (3)	O3—C21—H21A	109.5
C10—C11—H11	119.5	O3—C21—H21B	109.5
C12—C11—H11	119.5	H21A—C21—H21B	109.5
N3—C12—C11	121.6 (3)	O3—C21—H21C	109.5
N3—C12—C13	121.7 (3)	H21A—C21—H21C	109.5
C11—C12—C13	116.7 (3)	H21B—C21—H21C	109.5

*Hydrogen-bond geometry (Å, °)*

$D-H\cdots A$	$D-H$	$H\cdots A$	$D\cdots A$	$D-H\cdots A$
N2—H2···O3	0.90 (1)	2.07 (2)	2.936 (4)	160 (3)
O3—H3···O2 <sup>i</sup>	0.82	1.84	2.661 (3)	177
O1—H1···N1	0.82	2.02	2.727 (3)	145

Symmetry codes: (i)  $x+1, y, z$ .

## supplementary materials

---

Fig. 1




Fig. 2

