

2-(4-Chlorophenyl)-2-oxoethyl benzoate

Hoong-Kun Fun,^{a,*†} Tara Shahani,^a B. Garudachari,^b
Arun M. Isloor^b and M. N. Satyanarayan^c

^aX-ray Crystallography Unit, School of Physics, Universiti Sains Malaysia, 11800 USM, Penang, Malaysia, ^bOrganic Electronics Division, Department of Chemistry, National Institute of Technology-Karnataka, Surathkal, Mangalore 575 025, India, and ^cDepartment of Physics, National Institute of Technology-Karnataka, Surathkal, Mangalore 575 025, India
Correspondence e-mail: hkfun@usm.my


Received 17 June 2011; accepted 17 June 2011

Key indicators: single-crystal X-ray study; $T = 296$ K; mean $\sigma(\text{C}-\text{C}) = 0.002$ Å; R factor = 0.044; wR factor = 0.126; data-to-parameter ratio = 23.6.

In the title compound, $\text{C}_{15}\text{H}_{11}\text{ClO}_3$, the dihedral angle between the aromatic rings is $84.29(8)^\circ$. In the crystal, molecules are linked by weak $\text{C}-\text{H}\cdots\pi$ interactions.

Related literature

For applications of phenacyl benzoate derivatives, see: Rather & Reid (1919); Litera *et al.* (2006); Huang *et al.* (1996); Gandhi *et al.* (1995). For related structures, see: Ogata *et al.* (1987); Wan *et al.* (2006); Zhang *et al.* (2006). For reported melting-point details, see: Le *et al.* (2009). For bond-length data, see: Allen *et al.* (1987).



Experimental

Crystal data

$\text{C}_{15}\text{H}_{11}\text{ClO}_3$	$V = 1303.6(2) \text{ \AA}^3$
$M_r = 274.69$	$Z = 4$
Monoclinic, $P2_1/c$	Mo $K\alpha$ radiation
$a = 8.1955(9) \text{ \AA}$	$\mu = 0.29 \text{ mm}^{-1}$
$b = 10.8717(12) \text{ \AA}$	$T = 296 \text{ K}$
$c = 16.5420(15) \text{ \AA}$	$0.34 \times 0.19 \times 0.19 \text{ mm}$
$\beta = 117.816(4)^\circ$	

Data collection

Bruker SMART APEXII CCD diffractometer	11201 measured reflections
Absorption correction: multi-scan (SADABS; Bruker, 2009)	4052 independent reflections
$T_{\min} = 0.908$, $T_{\max} = 0.948$	2720 reflections with $I > 2\sigma(I)$
	$R_{\text{int}} = 0.021$

Refinement

$R[F^2 > 2\sigma(F^2)] = 0.044$	172 parameters
$wR(F^2) = 0.126$	H-atom parameters constrained
$S = 1.03$	$\Delta\rho_{\text{max}} = 0.31 \text{ e \AA}^{-3}$
4052 reflections	$\Delta\rho_{\text{min}} = -0.49 \text{ e \AA}^{-3}$

Table 1

Hydrogen-bond geometry (Å, °).

$Cg2$ is the centroid of the C10–C15 ring.

$D-H\cdots A$	$D-H$	$H\cdots A$	$D\cdots A$	$D-H\cdots A$
$\text{C8}-\text{H8A}\cdots Cg2^i$	0.97	2.96	3.4952 (17)	116

Symmetry code: (i) $-x + 1, -y + 1, -z + 2$.

Data collection: APEX2 (Bruker, 2009); cell refinement: SAINT (Bruker, 2009); data reduction: SAINT; program(s) used to solve structure: SHELXTL (Sheldrick, 2008); program(s) used to refine structure: SHELXTL; molecular graphics: SHELXTL; software used to prepare material for publication: SHELXTL and PLATON (Spek, 2009).

HKF and TSH thank Universiti Sains Malaysia (USM) for the Research University Grant (1001/PFIZIK/811160). TSH also thanks USM for the award of a research fellowship. AMI is thankful to the Department of Atomic Energy, Board for Research in Nuclear Sciences, Government of India for the Young Scientist award. GB thanks the Department of Information Technology, New Delhi, India, for financial support.

Supplementary data and figures for this paper are available from the IUCr electronic archives (Reference: HB5918).

References

- Allen, F. H., Kennard, O., Watson, D. G., Brammer, L., Orpen, A. G. & Taylor, R. (1987). *J. Chem. Soc. Perkin Trans. 2*, pp. S1–19.
- Bruker (2009). APEX2, SAINT and SADABS. Bruker AXS Inc., Madison, Wisconsin, USA.
- Gandhi, S. S., Bell, K. L. & Gibson, M. S. (1995). *Tetrahedron*, **51**, 13301–13308.
- Huang, W., Pian, J., Chen, B., Pei, W. & Ye, X. (1996). *Tetrahedron*, **52**, 10131–10136.
- Le, Z.-G., Xie, Z.-B. & Xu, J.-P. (2009). *Synth. Commun.* **39**, 743–747.
- Litera, J. K., Loya, A. D. & Klan, P. (2006). *J. Org. Chem.* **71**, 713–723.
- Ogata, M., Matsumoto, H., Takahashi, K., Shimizu, S., Kida, S. & Murabayashi, A. (1987). *J. Med. Chem.* **30**, 1054–1068.
- Rather, J. B. & Reid, E. (1919). *J. Am. Chem. Soc.* **41**, 75–83.
- Sheldrick, G. M. (2008). *Acta Cryst.* **A64**, 112–122.
- Spek, A. L. (2009). *Acta Cryst.* **D65**, 148–155.
- Wan, J., Li, X.-M., Peng, Z.-Z., Ouyang, P.-K. & Zhang, S.-S. (2006). *Acta Cryst.* **E62**, o2793–o2794.
- Zhang, S.-S., Wan, J., Peng, Z.-Z. & Bi, S. (2006). *Acta Cryst.* **E62**, o4348–o4349.

† Thomson Reuters ResearcherID: A-3561-2009.

supplementary materials

Acta Cryst. (2011). E67, o1802 [doi:10.1107/S160053681102383X]

2-(4-Chlorophenyl)-2-oxoethyl benzoate

H.-K. Fun, T. Shahani, B. Garudachari, A. M. Isloor and M. N. Satyganarayan

Comment

Phenacyl benzoate derivatives are very important in identification of organic acids (Rather & Reid, 1919) as they undergo photolysis in neutral and mild conditions (Litera *et al.*, 2006). They find applications in the field of synthetic chemistry for the synthesis of oxazoles, imidazoles (Huang *et al.*, 1996) and benzoxazepine (Gandhi *et al.*, 1995). We hereby report the crystal structure of the title compound, (I).

The asymmetric unit of title compound is shown in Fig. 1. The dihedral angle between the phenyl (C10–C15) ring and the chloro-substituted phenyl (C1–C6) ring is 84.29 (8)°. The bond lengths (Allen *et al.*, 1987) and angles are within normal ranges and are comparable to those closely related structures (Ogata *et al.*, 1987; Wan *et al.*, 2006; Zhang *et al.*, 2006).

In the crystal (Fig. 2), there are no classical hydrogen bonds but stabilization is provided by weak C—H \cdots π (Table 1) interactions, involving the Cg2 (C10–C15) ring.

Experimental

A mixture of benzoic acid (1.0 g, 0.0081 mol), potassium carbonate (1.23 g, 0.0089 mol) and 2-bromo-1-(4-chlorophenyl) ethanone (1.81 g, 0.0081 mol) in dimethylformamide (10 ml) was stirred at room temperature for 2 h. On cooling, colorless needle-shaped crystals of 2-(4-chlorophenyl)-2-oxoethyl benzoate begin to separate out. These were collected by filtration and recrystallized from ethanol to yield colourless blocks of (I). Yield: 2.10 g, 93.7%, *Mp*: 119–120 °C (Le *et al.*, 2009).

Refinement

All the H atoms were positioned geometrically [C–H = 0.93–0.97 Å] and were refined using a riding model, with $U_{\text{iso}}(\text{H}) = 1.2$. $U_{\text{eq}}(\text{C})$.

Figures




Fig. 1. The molecular structure of the title compound, showing 30% probability displacement ellipsoids.




Fig. 2. The crystal packing of the title compound (I).

2-(4-Chlorophenyl)-2-oxoethyl benzoate

Crystal data

$C_{15}H_{11}ClO_3$	$F(000) = 568$
$M_r = 274.69$	$D_x = 1.400 \text{ Mg m}^{-3}$
Monoclinic, $P2_1/c$	Mo $K\alpha$ radiation, $\lambda = 0.71073 \text{ \AA}$
Hall symbol: -P 2ybc	Cell parameters from 2797 reflections
$a = 8.1955 (9) \text{ \AA}$	$\theta = 2.3\text{--}27.1^\circ$
$b = 10.8717 (12) \text{ \AA}$	$\mu = 0.29 \text{ mm}^{-1}$
$c = 16.5420 (15) \text{ \AA}$	$T = 296 \text{ K}$
$\beta = 117.816 (4)^\circ$	Block, colourless
$V = 1303.6 (2) \text{ \AA}^3$	$0.34 \times 0.19 \times 0.19 \text{ mm}$
$Z = 4$	

Data collection

Bruker SMART APEXII CCD diffractometer	4052 independent reflections
Radiation source: fine-focus sealed tube graphite	2720 reflections with $I > 2\sigma(I)$
φ and ω scans	$R_{\text{int}} = 0.021$
Absorption correction: multi-scan (SADABS; Bruker, 2009)	$\theta_{\text{max}} = 30.8^\circ$, $\theta_{\text{min}} = 2.3^\circ$
$T_{\text{min}} = 0.908$, $T_{\text{max}} = 0.948$	$h = -11 \rightarrow 11$
11201 measured reflections	$k = -15 \rightarrow 15$
	$l = -20 \rightarrow 23$

Refinement

Refinement on F^2	Primary atom site location: structure-invariant direct methods
Least-squares matrix: full	Secondary atom site location: difference Fourier map
$R[F^2 > 2\sigma(F^2)] = 0.044$	Hydrogen site location: inferred from neighbouring sites
$wR(F^2) = 0.126$	H-atom parameters constrained
$S = 1.03$	$w = 1/[\sigma^2(F_o^2) + (0.0504P)^2 + 0.2649P]$
4052 reflections	where $P = (F_o^2 + 2F_c^2)/3$
172 parameters	$(\Delta/\sigma)_{\text{max}} < 0.001$
0 restraints	$\Delta\rho_{\text{max}} = 0.31 \text{ e \AA}^{-3}$
	$\Delta\rho_{\text{min}} = -0.49 \text{ e \AA}^{-3}$

Special details

Geometry. All e.s.d.'s (except the e.s.d. in the dihedral angle between two l.s. planes) are estimated using the full covariance matrix. The cell e.s.d.'s are taken into account individually in the estimation of e.s.d.'s in distances, angles and torsion angles; correlations

between e.s.d.'s in cell parameters are only used when they are defined by crystal symmetry. An approximate (isotropic) treatment of cell e.s.d.'s is used for estimating e.s.d.'s involving l.s. planes.

Refinement. Refinement of F^2 against ALL reflections. The weighted R -factor wR and goodness of fit S are based on F^2 , conventional R -factors R are based on F , with F set to zero for negative F^2 . The threshold expression of $F^2 > \sigma(F^2)$ is used only for calculating R -factors(gt) *etc.* and is not relevant to the choice of reflections for refinement. R -factors based on F^2 are statistically about twice as large as those based on F , and R -factors based on ALL data will be even larger.

Fractional atomic coordinates and isotropic or equivalent isotropic displacement parameters (\AA^2)

	x	y	z	$U_{\text{iso}}^*/U_{\text{eq}}$
C11	0.98407 (8)	1.29129 (4)	1.01748 (4)	0.07410 (18)
O1	0.41826 (16)	0.81706 (12)	0.81726 (7)	0.0648 (3)
O2	0.41542 (15)	0.64299 (10)	0.93059 (7)	0.0544 (3)
O3	0.59892 (16)	0.55553 (13)	0.87990 (9)	0.0722 (4)
C1	0.6218 (2)	1.03708 (15)	0.86536 (10)	0.0509 (3)
H1A	0.5394	1.0250	0.8040	0.061*
C2	0.7272 (2)	1.14249 (14)	0.89170 (12)	0.0545 (4)
H2A	0.7170	1.2011	0.8486	0.065*
C3	0.8484 (2)	1.15975 (14)	0.98327 (11)	0.0507 (3)
C4	0.8651 (2)	1.07434 (15)	1.04787 (11)	0.0555 (4)
H4A	0.9469	1.0874	1.1092	0.067*
C5	0.7593 (2)	0.96902 (14)	1.02070 (10)	0.0505 (3)
H5A	0.7701	0.9109	1.0642	0.061*
C6	0.63658 (18)	0.94852 (13)	0.92902 (9)	0.0435 (3)
C7	0.52322 (18)	0.83503 (14)	0.89703 (10)	0.0458 (3)
C8	0.5435 (2)	0.74141 (15)	0.96859 (10)	0.0524 (4)
H8A	0.6680	0.7087	0.9967	0.063*
H8B	0.5243	0.7814	1.0159	0.063*
C9	0.4586 (2)	0.55549 (15)	0.88590 (10)	0.0510 (3)
C10	0.3128 (2)	0.46016 (14)	0.84710 (9)	0.0473 (3)
C11	0.1476 (2)	0.47129 (15)	0.85064 (10)	0.0520 (3)
H11A	0.1261	0.5397	0.8781	0.062*
C12	0.0150 (2)	0.38051 (17)	0.81327 (11)	0.0622 (4)
H12A	-0.0966	0.3888	0.8147	0.075*
C13	0.0468 (3)	0.27865 (18)	0.77419 (12)	0.0688 (5)
H13A	-0.0424	0.2174	0.7498	0.083*
C14	0.2104 (3)	0.26665 (17)	0.77089 (12)	0.0706 (5)
H14A	0.2317	0.1970	0.7445	0.085*
C15	0.3436 (3)	0.35746 (15)	0.80651 (11)	0.0587 (4)
H15A	0.4534	0.3496	0.8032	0.070*

Atomic displacement parameters (\AA^2)

	U^{11}	U^{22}	U^{33}	U^{12}	U^{13}	U^{23}
C11	0.0961 (4)	0.0549 (3)	0.0902 (4)	-0.0203 (2)	0.0593 (3)	-0.0167 (2)
O1	0.0575 (7)	0.0764 (8)	0.0440 (6)	-0.0114 (6)	0.0100 (5)	0.0018 (5)
O2	0.0573 (6)	0.0533 (6)	0.0590 (6)	-0.0116 (5)	0.0325 (5)	-0.0065 (5)

supplementary materials

O3	0.0551 (7)	0.0808 (9)	0.0885 (9)	-0.0061 (6)	0.0399 (7)	-0.0124 (7)
C1	0.0525 (8)	0.0567 (9)	0.0423 (7)	0.0060 (7)	0.0212 (6)	0.0066 (6)
C2	0.0646 (9)	0.0483 (8)	0.0593 (9)	0.0064 (7)	0.0361 (8)	0.0109 (7)
C3	0.0559 (8)	0.0453 (7)	0.0615 (9)	-0.0017 (6)	0.0362 (7)	-0.0066 (7)
C4	0.0624 (9)	0.0564 (9)	0.0471 (8)	-0.0057 (7)	0.0251 (7)	-0.0062 (7)
C5	0.0583 (8)	0.0518 (8)	0.0404 (7)	-0.0027 (7)	0.0222 (6)	0.0035 (6)
C6	0.0434 (7)	0.0475 (7)	0.0411 (7)	0.0036 (6)	0.0210 (6)	0.0020 (6)
C7	0.0404 (7)	0.0535 (8)	0.0419 (7)	0.0010 (6)	0.0179 (6)	0.0008 (6)
C8	0.0554 (8)	0.0541 (8)	0.0462 (8)	-0.0102 (7)	0.0224 (7)	-0.0010 (7)
C9	0.0506 (8)	0.0537 (8)	0.0485 (8)	0.0010 (7)	0.0229 (7)	0.0053 (7)
C10	0.0526 (8)	0.0463 (7)	0.0405 (7)	0.0016 (6)	0.0196 (6)	0.0066 (6)
C11	0.0545 (8)	0.0520 (8)	0.0486 (8)	-0.0029 (7)	0.0233 (7)	0.0030 (7)
C12	0.0579 (9)	0.0679 (10)	0.0539 (9)	-0.0110 (8)	0.0203 (8)	0.0042 (8)
C13	0.0739 (12)	0.0609 (10)	0.0541 (10)	-0.0154 (9)	0.0151 (9)	0.0009 (8)
C14	0.0960 (14)	0.0496 (9)	0.0526 (10)	0.0016 (9)	0.0233 (10)	-0.0037 (8)
C15	0.0692 (10)	0.0547 (9)	0.0520 (9)	0.0092 (8)	0.0282 (8)	0.0051 (7)

Geometric parameters (\AA , $^\circ$)

C11—C3	1.7362 (16)	C7—C8	1.510 (2)
O1—C7	1.2083 (17)	C8—H8A	0.9700
O2—C9	1.3492 (19)	C8—H8B	0.9700
O2—C8	1.4235 (18)	C9—C10	1.482 (2)
O3—C9	1.1996 (18)	C10—C15	1.385 (2)
C1—C2	1.378 (2)	C10—C11	1.388 (2)
C1—C6	1.390 (2)	C11—C12	1.382 (2)
C1—H1A	0.9300	C11—H11A	0.9300
C2—C3	1.383 (2)	C12—C13	1.367 (3)
C2—H2A	0.9300	C12—H12A	0.9300
C3—C4	1.373 (2)	C13—C14	1.373 (3)
C4—C5	1.379 (2)	C13—H13A	0.9300
C4—H4A	0.9300	C14—C15	1.384 (3)
C5—C6	1.3920 (19)	C14—H14A	0.9300
C5—H5A	0.9300	C15—H15A	0.9300
C6—C7	1.486 (2)		
C9—O2—C8	116.36 (12)	O2—C8—H8B	109.3
C2—C1—C6	121.10 (14)	C7—C8—H8B	109.3
C2—C1—H1A	119.5	H8A—C8—H8B	107.9
C6—C1—H1A	119.5	O3—C9—O2	123.48 (15)
C1—C2—C3	118.90 (14)	O3—C9—C10	125.06 (15)
C1—C2—H2A	120.6	O2—C9—C10	111.46 (13)
C3—C2—H2A	120.6	C15—C10—C11	119.52 (15)
C4—C3—C2	121.36 (15)	C15—C10—C9	118.75 (14)
C4—C3—C11	119.15 (13)	C11—C10—C9	121.72 (14)
C2—C3—C11	119.49 (13)	C12—C11—C10	119.86 (16)
C3—C4—C5	119.25 (15)	C12—C11—H11A	120.1
C3—C4—H4A	120.4	C10—C11—H11A	120.1
C5—C4—H4A	120.4	C13—C12—C11	120.41 (18)
C4—C5—C6	120.85 (14)	C13—C12—H12A	119.8

C4—C5—H5A	119.6	C11—C12—H12A	119.8
C6—C5—H5A	119.6	C12—C13—C14	120.06 (17)
C1—C6—C5	118.54 (14)	C12—C13—H13A	120.0
C1—C6—C7	119.12 (13)	C14—C13—H13A	120.0
C5—C6—C7	122.34 (13)	C13—C14—C15	120.43 (17)
O1—C7—C6	122.11 (14)	C13—C14—H14A	119.8
O1—C7—C8	120.59 (14)	C15—C14—H14A	119.8
C6—C7—C8	117.30 (12)	C14—C15—C10	119.69 (17)
O2—C8—C7	111.81 (12)	C14—C15—H15A	120.2
O2—C8—H8A	109.3	C10—C15—H15A	120.2
C7—C8—H8A	109.3		
C6—C1—C2—C3	-0.4 (2)	C6—C7—C8—O2	174.36 (12)
C1—C2—C3—C4	-0.1 (2)	C8—O2—C9—O3	2.5 (2)
C1—C2—C3—C11	179.31 (11)	C8—O2—C9—C10	-178.30 (12)
C2—C3—C4—C5	0.3 (2)	O3—C9—C10—C15	5.2 (2)
C11—C3—C4—C5	-179.14 (12)	O2—C9—C10—C15	-173.94 (13)
C3—C4—C5—C6	0.0 (2)	O3—C9—C10—C11	-174.87 (16)
C2—C1—C6—C5	0.7 (2)	O2—C9—C10—C11	5.9 (2)
C2—C1—C6—C7	-178.58 (13)	C15—C10—C11—C12	-0.4 (2)
C4—C5—C6—C1	-0.5 (2)	C9—C10—C11—C12	179.72 (14)
C4—C5—C6—C7	178.73 (14)	C10—C11—C12—C13	1.1 (2)
C1—C6—C7—O1	0.0 (2)	C11—C12—C13—C14	-0.8 (3)
C5—C6—C7—O1	-179.21 (15)	C12—C13—C14—C15	-0.3 (3)
C1—C6—C7—C8	-179.92 (13)	C13—C14—C15—C10	1.1 (3)
C5—C6—C7—C8	0.9 (2)	C11—C10—C15—C14	-0.7 (2)
C9—O2—C8—C7	79.04 (17)	C9—C10—C15—C14	179.19 (15)
O1—C7—C8—O2	-5.6 (2)		

Hydrogen-bond geometry (Å, °)

Cg2 is the centroid of the C10—C15 ring.

<i>D</i> —H... <i>A</i>	<i>D</i> —H	H... <i>A</i>	<i>D</i> ... <i>A</i>	<i>D</i> —H... <i>A</i>
C8—H8A...Cg2 ⁱ	0.97	2.96	3.4952 (17)	116

Symmetry codes: (i) $-x+1, -y+1, -z+2$.

Fig. 1




Fig. 2

