

Poly[aqua μ_5 -5-(isonicotinamido)-isophthalato] μ_4 -5-(isonicotinamido)-isophthalato]cerium(III)silver(I)]

Xue Nie* and Jing-Nian Qu

Department of Chemistry and Materials Science, Hengyang, Hunan 421008, People's Republic of China
Correspondence e-mail: xuehuanie@126.com


Received 22 June 2011; accepted 11 July 2011

Key indicators: single-crystal X-ray study; $T = 293\text{ K}$; mean $\sigma(\text{C}-\text{C}) = 0.008\text{ \AA}$; R factor = 0.035; wR factor = 0.081; data-to-parameter ratio = 11.8.

The $4d$ - $4f$ heteronuclear title complex, $[\text{AgCe}(\text{C}_{14}\text{H}_8\text{N}_2\text{O}_5)_2(\text{H}_2\text{O})]_n$, has a three-dimensional framework structure, generated by the carboxylate and pyridyl groups of the 5-(isonicotinamido)isophthalate (INAIP) ligands bridging the metal ions. The Ce^{III} atom is coordinated by eight O atoms from six INAIP ligands and a water molecule in a distorted tricapped trigonal-prismatic geometry, while the Ag^I atom has a distorted trigonal-planar AgN_2O geometry. $\text{O}-\text{H}\cdots\text{O}$ and $\text{N}-\text{H}\cdots\text{O}$ hydrogen bonds and $\pi-\pi$ interactions between the pyridine and benzene rings [centroid–centroid distances = 3.642 (4) and 3.624 (3) \AA] stabilize the crystal structure.

Related literature

For background to coordination polymers, see: Abourahma *et al.* (2002); Costes *et al.* (2004); Kapoor *et al.* (2002). For background to lanthanide and transition metal heterometallic compounds, see: Chen *et al.* (2010); Cheng *et al.* (2006); Lin *et al.* (2009); Zhang *et al.* (2005).



Experimental

Crystal data

$[\text{AgCe}(\text{C}_{14}\text{H}_8\text{N}_2\text{O}_5)_2(\text{H}_2\text{O})]$

$M_r = 834.45$	$\gamma = 102.885(2)^\circ$
Triclinic, $\bar{P}\bar{1}$	$V = 1375.2(3)\text{ \AA}^3$
$a = 10.4869(14)\text{ \AA}$	$Z = 2$
$b = 11.1540(15)\text{ \AA}$	Mo $K\alpha$ radiation
$c = 13.7276(18)\text{ \AA}$	$\mu = 2.42\text{ mm}^{-1}$
$\alpha = 107.853(1)^\circ$	$T = 293\text{ K}$
$\beta = 106.778(2)^\circ$	$0.20 \times 0.14 \times 0.10\text{ mm}$

Data collection

Bruker APEX CCD diffractometer	6903 measured reflections
Absorption correction: multi-scan (<i>SADABS</i> ; Bruker, 2001)	4777 independent reflections
$(SADABS$; Bruker, 2001)	4178 reflections with $I > 2\sigma(I)$
$T_{\min} = 0.643$, $T_{\max} = 0.794$	$R_{\text{int}} = 0.053$

Refinement

$R[F^2 > 2\sigma(F^2)] = 0.035$	406 parameters
$wR(F^2) = 0.081$	H-atom parameters constrained
$S = 0.97$	$\Delta\rho_{\max} = 1.40\text{ e \AA}^{-3}$
4777 reflections	$\Delta\rho_{\min} = -1.44\text{ e \AA}^{-3}$

Table 1
Hydrogen-bond geometry (\AA , $^\circ$).

$D-\text{H}\cdots A$	$D-\text{H}$	$\text{H}\cdots A$	$D\cdots A$	$D-\text{H}\cdots A$
O1W—H1WA \cdots O5 ⁱ	0.85	2.08	2.836 (5)	147
O1W—H1WB \cdots O4 ⁱⁱ	0.85	2.01	2.691 (6)	137
N2—H2 \cdots O10 ⁱⁱⁱ	0.86	2.01	2.788 (7)	149
N4—H4 \cdots O4 ⁱ	0.86	2.08	2.926 (6)	167

Symmetry codes: (i) $-x + 1, -y + 1, -z + 2$; (ii) $x + 1, y, z$; (iii) $x, y + 1, z$.

Data collection: *SMART* (Bruker, 2007); cell refinement: *SAINT* (Bruker, 2007); data reduction: *SAINT*; program(s) used to solve structure: *SHELXTL* (Sheldrick, 2008); program(s) used to refine structure: *SHELXTL*; molecular graphics: *SHELXTL*; software used to prepare material for publication: *SHELXTL*.

This work was supported by the Hengyang Bureau of Science & Technology (grant No. 2009 K J28).

Supplementary data and figures for this paper are available from the IUCr electronic archives (Reference: HY2446).

References

- Abourahma, H., Moulton, B., Kravtsov, V. & Zaworotko, M. J. (2002). *J. Am. Chem. Soc.* **124**, 9990–9991.
- Bruker (2001). *SADABS*. Bruker AXS Inc., Madison, Wisconsin, USA.
- Bruker (2007). *SMART* and *SAINT*. Bruker AXS Inc., Madison, Wisconsin, USA.
- Chen, M.-S., Su, Z., Chen, M., Chen, S.-S., Li, Y.-Z. & Sun, W.-Y. (2010). *CrystEngComm*, **12**, 3267–3276.
- Cheng, J.-W., Zhang, J., Zheng, S.-T., Zhang, M.-B. & Yang, G.-Y. (2006). *Angew. Chem. Int. Ed.* **45**, 73–77.
- Costes, J. P., Dahan, F., Donnadieu, B., Douton, M. J. R., Garcia, M. I. F., Bousseksou, A. & Tuchagues, J. P. (2004). *Inorg. Chem.* **43**, 2736–2744.
- Kapoor, P., Pathak, A., Kapoor, R., Venugopalan, P., Corbella, M., Rodrmiguez, M., Robles, J. & Llobet, A. (2002). *Inorg. Chem.* **41**, 6153–6160.
- Lin, X.-M., Ying, Y., Chen, L., Fang, H.-C., Zhou, Z.-Y., Zhan, Q.-G. & Cai, Y.-P. (2009). *Inorg. Chem. Commun.* **12**, 316–320.
- Sheldrick, G. M. (2008). *Acta Cryst. A* **64**, 112–122.
- Zhang, M.-B., Zhang, J., Zheng, S.-T. & Yang, G.-Y. (2005). *Angew. Chem. Int. Ed.* **44**, 1385–1388.

supplementary materials

Acta Cryst. (2011). E67, m1107-m1108 [doi:10.1107/S1600536811027668]

Poly[aqua μ_5 -5-(isonicotinamido)isophthalato] $[\mu_4$ -5-(isonicotinamido)isophthalato]cerium(III)silver(I)]

X. Nie and J.-N. Qu

Comment

In recent years, coordination polymeric frameworks have attracted great attention because of their potential applications and intriguing structure topologies (Abourahma *et al.*, 2002; Costes *et al.*, 2004; Kapoor *et al.*, 2002). To obtain d-f coordination polymers is more important. In general, multidentate ligands containing both N- and O-donor atoms are usually employed in the construction of lanthanide (Ln) and transition metal (M) heterometallic structures, in keeping with the typical coordination behaviors of Ln and M ions under different reaction conditions (Cheng *et al.*, 2006; Lin *et al.*, 2009; Zhang *et al.*, 2005). Compared with other N-heterocyclic acids, 5-(isonicotinamido)isophthalic acid (H₂INAIP) shows richer coordination modes due to its two carboxylate groups and one pyridyl group, and accordingly it is an excellent candidate for the construction of metal-organic frameworks (Chen *et al.*, 2010). In this paper, we report the synthesis and structure of the title complex (Fig. 1), a 4d-4f heterometallic coordination polymer.

It is interesting that two INAIP ligands exhibit different coordination modes: one coordinated to three Ce^{III} atoms and two Ag^I atoms while the other coordinated to three Ce^{III} atoms and one Ag^I atom, originated from the different coordination modes of the carboxylate groups. If the Ag—N and Ag—O connections are neglected, a two-dimensional (4, 4) bilayer network is formed by the Ce^{III}–carboxylate coordination. The (4, 4) nets are linked together by Ag—N and Ag—O coordination interactions, forming a complicated three-dimensional coordination net (Fig. 2).

Experimental

A mixture of Ce(NO₃)₃·6H₂O (22.0 mg, 0.05 mmol), H₂INAIP (28.6 mg, 0.1 mmol), AgNO₃ (8.5 mg, 0.05 mmol), NaOH (6.0 mg, 0.15 mmol) and H₂O (10 ml) was heated in a 16 ml Teflon-lined reaction vessel at 453 K for 4 d. The reaction mixture was cooled to room temperature over a period of 40 h. The product was collected by filtration, washed with H₂O and air dried.

Refinement

H atoms bonded to C and N atoms were positioned geometrically and refined as riding atoms, with C—H = 0.93 and N—H = 0.86 Å and $U_{\text{iso}}(\text{H}) = 1.2U_{\text{eq}}(\text{C}, \text{N})$. The water H atoms were found from a difference Fourier maps and refined as riding, with O—H = 0.85 Å and with $U_{\text{iso}}(\text{H}) = 1.2U_{\text{eq}}(\text{O})$. The highest residual electron density was found at 0.93 Å from Ag1 atom and the deepest hole at 0.86 Å from Ag1 atom.

supplementary materials

Figures




Fig. 1. The asymmetric unit of the title compound. Displacement ellipsoids are drawn at the 30% probability level. [Symmetry codes: (A) 1-x, 1-y, 1-z; (B) 1+x, y, z; (C) 1-x, -y, 1-z; (D) x, 1+y, z; (E) x, y, 1+z; (F) 2-x, 1-y, 3-z.]




Fig. 2. A view of the three-dimensional framework of the title compound.

Poly[aqua[μ_5 -5-(isonicotinamido)isophthalato][μ_4 -5- (isonicotinamido)isophthalato]cerium(III)silver(I)]

Crystal data

[AgCe(C ₁₄ H ₈ N ₂ O ₅) ₂ (H ₂ O)]	Z = 2
M _r = 834.45	F(000) = 814
Triclinic, PT	D _x = 2.015 Mg m ⁻³
Hall symbol: -P 1	Mo K α radiation, λ = 0.71073 Å
a = 10.4869 (14) Å	Cell parameters from 4777 reflections
b = 11.1540 (15) Å	θ = 2.0–25.0°
c = 13.7276 (18) Å	μ = 2.42 mm ⁻¹
α = 107.853 (1)°	T = 293 K
β = 106.778 (2)°	Block, colorless
γ = 102.885 (2)°	0.20 × 0.14 × 0.10 mm
V = 1375.2 (3) Å ³	

Data collection

Bruker APEX CCD diffractometer	4777 independent reflections
Radiation source: fine-focus sealed tube graphite	4178 reflections with $I > 2\sigma(I)$
φ and ω scans	$R_{\text{int}} = 0.053$
Absorption correction: multi-scan (SADABS; Bruker, 2001)	$\theta_{\text{max}} = 25.0^\circ$, $\theta_{\text{min}} = 2.0^\circ$
$T_{\text{min}} = 0.643$, $T_{\text{max}} = 0.794$	$h = -12 \rightarrow 12$
6903 measured reflections	$k = -11 \rightarrow 13$
	$l = -16 \rightarrow 12$

Refinement

Refinement on F^2	Primary atom site location: structure-invariant direct methods
Least-squares matrix: full	Secondary atom site location: difference Fourier map
$R[F^2 > 2\sigma(F^2)] = 0.035$	Hydrogen site location: inferred from neighbouring sites
$wR(F^2) = 0.081$	H-atom parameters constrained
$S = 0.97$	$w = 1/[\sigma^2(F_o^2) + (0.0347P)^2]$ where $P = (F_o^2 + 2F_c^2)/3$
4777 reflections	$(\Delta/\sigma)_{\max} = 0.001$
406 parameters	$\Delta\rho_{\max} = 1.40 \text{ e \AA}^{-3}$
0 restraints	$\Delta\rho_{\min} = -1.44 \text{ e \AA}^{-3}$

Fractional atomic coordinates and isotropic or equivalent isotropic displacement parameters (\AA^2)

	<i>x</i>	<i>y</i>	<i>z</i>	$U_{\text{iso}}^*/U_{\text{eq}}$
Ag1	0.88225 (5)	0.23184 (7)	1.55049 (4)	0.06266 (18)
Ce1	0.63603 (3)	0.45288 (2)	0.61285 (2)	0.01607 (9)
C1	0.3260 (5)	0.5683 (4)	0.5961 (4)	0.0200 (10)
C2	0.2981 (5)	0.5843 (4)	0.7000 (4)	0.0202 (10)
C3	0.1600 (5)	0.5553 (4)	0.6961 (4)	0.0222 (10)
H3	0.0837	0.5240	0.6288	0.027*
C4	0.1376 (5)	0.5735 (5)	0.7933 (4)	0.0234 (10)
C5	0.2530 (5)	0.6195 (5)	0.8932 (4)	0.0247 (11)
H5	0.2378	0.6312	0.9584	0.030*
C6	0.3879 (5)	0.6476 (4)	0.8964 (4)	0.0209 (10)
C7	0.4116 (5)	0.6312 (4)	0.8000 (4)	0.0223 (10)
H7	0.5037	0.6517	0.8023	0.027*
C8	-0.0077 (5)	0.5580 (5)	0.7942 (4)	0.0240 (11)
C9	0.5223 (5)	0.6542 (5)	1.0769 (4)	0.0236 (11)
C10	0.6688 (5)	0.7096 (5)	1.1656 (4)	0.0260 (11)
C11	0.6840 (6)	0.7319 (6)	1.2735 (4)	0.0398 (14)
H11	0.6052	0.7199	1.2925	0.048*
C12	0.8176 (6)	0.7723 (7)	1.3521 (4)	0.0469 (16)
H12	0.8272	0.7904	1.4252	0.056*
C13	0.9185 (6)	0.7691 (7)	1.2276 (5)	0.0481 (16)
H13	0.9993	0.7828	1.2112	0.058*
C14	0.7895 (6)	0.7313 (6)	1.1433 (4)	0.0418 (14)
H14	0.7840	0.7205	1.0721	0.050*
C15	0.6561 (5)	0.2158 (5)	0.6570 (4)	0.0240 (11)
C16	0.6919 (5)	0.1164 (4)	0.7036 (4)	0.0217 (10)
C17	0.7361 (5)	0.1512 (5)	0.8175 (4)	0.0258 (11)
H17	0.7431	0.2353	0.8638	0.031*
C18	0.7693 (5)	0.0620 (5)	0.8619 (4)	0.0252 (11)
C19	0.7550 (5)	-0.0648 (5)	0.7921 (4)	0.0241 (11)
H19	0.7773	-0.1255	0.8219	0.029*

supplementary materials

C20	0.7076 (5)	-0.1014 (4)	0.6784 (4)	0.0215 (10)
C21	0.6768 (5)	-0.0101 (4)	0.6344 (4)	0.0223 (10)
H21	0.6460	-0.0340	0.5582	0.027*
C22	0.6810 (5)	-0.2419 (5)	0.6031 (4)	0.0212 (10)
C23	0.7450 (6)	0.0246 (5)	1.0210 (4)	0.0356 (13)
C24	0.7834 (6)	0.0781 (5)	1.1428 (4)	0.0318 (12)
C25	0.6756 (6)	0.0579 (6)	1.1812 (5)	0.0428 (14)
H25	0.5819	0.0151	1.1312	0.051*
C26	0.7064 (7)	0.1008 (6)	1.2925 (5)	0.0437 (14)
H26	0.6321	0.0865	1.3165	0.052*
C27	0.9447 (6)	0.1839 (6)	1.3330 (5)	0.0452 (15)
H27	1.0375	0.2270	1.3846	0.054*
C28	0.9196 (6)	0.1435 (6)	1.2209 (4)	0.0384 (14)
H28	0.9950	0.1608	1.1985	0.046*
N1	0.9338 (5)	0.7871 (5)	1.3303 (4)	0.0409 (12)
N2	0.5085 (4)	0.6990 (4)	0.9965 (3)	0.0249 (9)
H2	0.5787	0.7647	1.0061	0.030*
N3	0.8395 (5)	0.1627 (5)	1.3687 (4)	0.0401 (12)
N4	0.8112 (5)	0.0994 (4)	0.9783 (3)	0.0285 (10)
H4	0.8810	0.1721	1.0226	0.034*
O1	0.4458 (3)	0.5621 (3)	0.5960 (2)	0.0233 (7)
O2	0.2324 (4)	0.5616 (4)	0.5135 (3)	0.0329 (9)
O3	-0.1121 (3)	0.4777 (3)	0.7074 (3)	0.0280 (8)
O4	-0.0163 (4)	0.6309 (4)	0.8809 (3)	0.0341 (9)
O5	0.4260 (4)	0.5710 (3)	1.0791 (3)	0.0316 (8)
O6	0.6166 (4)	0.3009 (3)	0.7118 (3)	0.0280 (8)
O7	0.6661 (4)	0.2109 (3)	0.5664 (2)	0.0241 (7)
O8	0.6024 (4)	-0.2802 (3)	0.5023 (3)	0.0270 (8)
O9	0.7369 (3)	-0.3125 (3)	0.6448 (3)	0.0259 (8)
O10	0.6548 (5)	-0.0862 (4)	0.9625 (3)	0.0639 (14)
O1W	0.7032 (3)	0.5815 (3)	0.8217 (3)	0.0266 (8)
H1WB	0.7778	0.5704	0.8568	0.032*
H1WA	0.6372	0.5446	0.8383	0.032*

Atomic displacement parameters (\AA^2)

	U^{11}	U^{22}	U^{33}	U^{12}	U^{13}	U^{23}
Ag1	0.0349 (3)	0.1311 (5)	0.0321 (3)	0.0401 (3)	0.0142 (2)	0.0360 (3)
Ce1	0.01710 (15)	0.01732 (14)	0.01439 (14)	0.00767 (11)	0.00504 (11)	0.00675 (10)
C1	0.022 (3)	0.018 (2)	0.019 (2)	0.006 (2)	0.008 (2)	0.005 (2)
C2	0.020 (3)	0.025 (2)	0.020 (2)	0.012 (2)	0.009 (2)	0.011 (2)
C3	0.020 (3)	0.025 (2)	0.017 (2)	0.008 (2)	0.003 (2)	0.007 (2)
C4	0.022 (3)	0.027 (3)	0.021 (3)	0.009 (2)	0.008 (2)	0.009 (2)
C5	0.026 (3)	0.033 (3)	0.015 (2)	0.015 (2)	0.006 (2)	0.009 (2)
C6	0.021 (3)	0.021 (2)	0.018 (2)	0.007 (2)	0.004 (2)	0.008 (2)
C7	0.019 (3)	0.027 (3)	0.024 (3)	0.010 (2)	0.009 (2)	0.012 (2)
C8	0.022 (3)	0.030 (3)	0.020 (3)	0.009 (2)	0.006 (2)	0.013 (2)
C9	0.024 (3)	0.026 (3)	0.015 (2)	0.006 (2)	0.005 (2)	0.005 (2)

C10	0.028 (3)	0.027 (3)	0.025 (3)	0.011 (2)	0.008 (2)	0.013 (2)
C11	0.026 (3)	0.072 (4)	0.025 (3)	0.019 (3)	0.011 (2)	0.021 (3)
C12	0.035 (4)	0.081 (5)	0.023 (3)	0.024 (3)	0.006 (3)	0.020 (3)
C13	0.025 (3)	0.082 (5)	0.040 (4)	0.017 (3)	0.011 (3)	0.030 (3)
C14	0.032 (3)	0.070 (4)	0.022 (3)	0.011 (3)	0.011 (3)	0.021 (3)
C15	0.021 (3)	0.024 (3)	0.029 (3)	0.010 (2)	0.008 (2)	0.015 (2)
C16	0.022 (3)	0.019 (2)	0.026 (3)	0.008 (2)	0.008 (2)	0.012 (2)
C17	0.034 (3)	0.020 (2)	0.024 (3)	0.010 (2)	0.012 (2)	0.008 (2)
C18	0.029 (3)	0.024 (3)	0.019 (2)	0.006 (2)	0.009 (2)	0.008 (2)
C19	0.027 (3)	0.023 (2)	0.024 (3)	0.011 (2)	0.005 (2)	0.013 (2)
C20	0.023 (3)	0.018 (2)	0.024 (3)	0.007 (2)	0.008 (2)	0.010 (2)
C21	0.023 (3)	0.023 (2)	0.020 (2)	0.008 (2)	0.006 (2)	0.010 (2)
C22	0.017 (3)	0.026 (2)	0.022 (3)	0.006 (2)	0.009 (2)	0.011 (2)
C23	0.041 (3)	0.029 (3)	0.030 (3)	0.007 (3)	0.009 (3)	0.011 (3)
C24	0.040 (3)	0.030 (3)	0.028 (3)	0.011 (3)	0.013 (3)	0.015 (2)
C25	0.034 (3)	0.048 (3)	0.032 (3)	0.006 (3)	0.005 (3)	0.011 (3)
C26	0.041 (4)	0.063 (4)	0.035 (3)	0.020 (3)	0.020 (3)	0.024 (3)
C27	0.035 (4)	0.076 (4)	0.033 (3)	0.026 (3)	0.013 (3)	0.028 (3)
C28	0.041 (4)	0.058 (4)	0.028 (3)	0.023 (3)	0.018 (3)	0.024 (3)
N1	0.026 (3)	0.064 (3)	0.028 (3)	0.016 (2)	0.004 (2)	0.018 (2)
N2	0.019 (2)	0.030 (2)	0.020 (2)	0.0024 (18)	0.0017 (18)	0.0124 (18)
N3	0.043 (3)	0.061 (3)	0.030 (3)	0.029 (3)	0.018 (2)	0.023 (2)
N4	0.036 (3)	0.023 (2)	0.018 (2)	0.0039 (19)	0.0031 (19)	0.0088 (18)
O1	0.0193 (18)	0.0286 (18)	0.0219 (17)	0.0091 (15)	0.0082 (15)	0.0089 (15)
O2	0.028 (2)	0.060 (2)	0.0197 (18)	0.0253 (19)	0.0103 (16)	0.0198 (18)
O3	0.0173 (18)	0.038 (2)	0.0197 (18)	0.0067 (16)	0.0035 (15)	0.0058 (16)
O4	0.0186 (19)	0.049 (2)	0.0256 (19)	0.0095 (17)	0.0089 (16)	0.0045 (17)
O5	0.025 (2)	0.042 (2)	0.0259 (19)	0.0067 (17)	0.0073 (16)	0.0177 (17)
O6	0.043 (2)	0.0247 (18)	0.0283 (19)	0.0201 (17)	0.0185 (17)	0.0148 (16)
O7	0.032 (2)	0.0250 (17)	0.0207 (18)	0.0128 (15)	0.0115 (15)	0.0124 (15)
O8	0.031 (2)	0.0209 (17)	0.0209 (18)	0.0057 (15)	0.0036 (16)	0.0054 (15)
O9	0.0257 (19)	0.0216 (17)	0.0313 (19)	0.0120 (15)	0.0069 (16)	0.0128 (15)
O10	0.083 (4)	0.044 (2)	0.033 (2)	-0.021 (2)	0.020 (2)	0.007 (2)
O1W	0.0233 (19)	0.0369 (19)	0.0245 (18)	0.0140 (16)	0.0117 (15)	0.0134 (16)

Geometric parameters (Å, °)

Ag1—N1 ⁱ	2.239 (4)	C13—N1	1.318 (7)
Ag1—N3	2.246 (4)	C13—C14	1.374 (8)
Ag1—O7 ⁱⁱ	2.306 (3)	C13—H13	0.9300
Ce1—O9 ⁱⁱⁱ	2.439 (3)	C14—H14	0.9300
Ce1—O8 ^{iv}	2.483 (3)	C15—O6	1.247 (5)
Ce1—O6	2.485 (3)	C15—O7	1.264 (5)
Ce1—O3 ^v	2.493 (3)	C15—C16	1.503 (6)
Ce1—O2 ^{vi}	2.499 (3)	C16—C21	1.383 (6)
Ce1—O1	2.551 (3)	C16—C17	1.393 (6)
Ce1—O1W	2.576 (3)	C17—C18	1.374 (6)
Ce1—O1 ^{vi}	2.685 (3)	C17—H17	0.9300

supplementary materials

Ce1—O7	2.690 (3)	C18—C19	1.390 (6)
C1—O2	1.237 (5)	C18—N4	1.423 (6)
C1—O1	1.274 (5)	C19—C20	1.386 (6)
C1—C2	1.503 (6)	C19—H19	0.9300
C2—C7	1.383 (6)	C20—C21	1.385 (6)
C2—C3	1.393 (6)	C20—C22	1.499 (6)
C3—C4	1.384 (6)	C21—H21	0.9300
C3—H3	0.9300	C22—O9	1.256 (5)
C4—C5	1.394 (6)	C22—O8	1.265 (5)
C4—C8	1.498 (6)	C23—O10	1.224 (6)
C5—C6	1.364 (6)	C23—N4	1.341 (6)
C5—H5	0.9300	C23—C24	1.485 (7)
C6—C7	1.382 (6)	C24—C28	1.376 (8)
C6—N2	1.416 (6)	C24—C25	1.382 (8)
C7—H7	0.9300	C25—C26	1.365 (7)
C8—O3	1.255 (6)	C25—H25	0.9300
C8—O4	1.260 (5)	C26—N3	1.343 (8)
C9—O5	1.225 (6)	C26—H26	0.9300
C9—N2	1.329 (6)	C27—N3	1.335 (7)
C9—C10	1.503 (7)	C27—C28	1.388 (7)
C10—C14	1.374 (7)	C27—H27	0.9300
C10—C11	1.378 (7)	C28—H28	0.9300
C11—C12	1.371 (7)	N2—H2	0.8600
C11—H11	0.9300	N4—H4	0.8600
C12—N1	1.324 (7)	O1W—H1WB	0.8500
C12—H12	0.9300	O1W—H1WA	0.8500
N1 ⁱ —Ag1—N3	121.79 (16)	O5—C9—C10	121.0 (4)
N1 ⁱ —Ag1—O7 ⁱⁱ	124.05 (14)	N2—C9—C10	115.1 (4)
N3—Ag1—O7 ⁱⁱ	106.74 (14)	C14—C10—C11	117.8 (5)
O9 ⁱⁱⁱ —Ce1—O8 ^{iv}	133.09 (10)	C14—C10—C9	122.9 (4)
O9 ⁱⁱⁱ —Ce1—O6	142.14 (11)	C11—C10—C9	119.1 (5)
O8 ^{iv} —Ce1—O6	76.29 (11)	C12—C11—C10	118.6 (5)
O9 ⁱⁱⁱ —Ce1—O3 ^v	83.67 (11)	C12—C11—H11	120.7
O8 ^{iv} —Ce1—O3 ^v	141.72 (10)	C10—C11—H11	120.7
O6—Ce1—O3 ^v	77.23 (11)	N1—C12—C11	123.9 (5)
O9 ⁱⁱⁱ —Ce1—O2 ^{vi}	77.92 (11)	N1—C12—H12	118.0
O8 ^{iv} —Ce1—O2 ^{vi}	105.14 (11)	C11—C12—H12	118.0
O6—Ce1—O2 ^{vi}	122.35 (10)	N1—C13—C14	123.5 (5)
O3 ^v —Ce1—O2 ^{vi}	67.59 (10)	N1—C13—H13	118.2
O9 ⁱⁱⁱ —Ce1—O1	69.33 (10)	C14—C13—H13	118.2
O8 ^{iv} —Ce1—O1	69.60 (10)	C10—C14—C13	119.1 (5)
O6—Ce1—O1	113.51 (10)	C10—C14—H14	120.4
O3 ^v —Ce1—O1	147.60 (10)	C13—C14—H14	120.4
O2 ^{vi} —Ce1—O1	120.84 (10)	O6—C15—O7	122.8 (4)
O9 ⁱⁱⁱ —Ce1—O1W	75.72 (10)	O6—C15—C16	116.8 (4)

O8 ^{iv} —Ce1—O1W	119.75 (11)	O7—C15—C16	120.4 (4)
O6—Ce1—O1W	67.66 (10)	C21—C16—C17	119.9 (4)
O3 ^v —Ce1—O1W	73.47 (10)	C21—C16—C15	120.7 (4)
O2 ^{vi} —Ce1—O1W	134.75 (11)	C17—C16—C15	119.4 (4)
O1—Ce1—O1W	82.50 (10)	C18—C17—C16	120.4 (4)
O9 ⁱⁱⁱ —Ce1—O1 ^{vi}	79.04 (10)	C18—C17—H17	119.8
O8 ^{iv} —Ce1—O1 ^{vi}	69.80 (10)	C16—C17—H17	119.8
O6—Ce1—O1 ^{vi}	138.81 (10)	C17—C18—C19	119.6 (4)
O3 ^v —Ce1—O1 ^{vi}	117.28 (10)	C17—C18—N4	119.0 (4)
O2 ^{vi} —Ce1—O1 ^{vi}	49.96 (10)	C19—C18—N4	121.3 (4)
O1—Ce1—O1 ^{vi}	75.80 (10)	C20—C19—C18	120.4 (4)
O1W—Ce1—O1 ^{vi}	151.22 (9)	C20—C19—H19	119.8
O9 ⁱⁱⁱ —Ce1—O7	147.39 (10)	C18—C19—H19	119.8
O8 ^{iv} —Ce1—O7	72.15 (10)	C21—C20—C19	119.7 (4)
O6—Ce1—O7	50.26 (9)	C21—C20—C22	120.2 (4)
O3 ^v —Ce1—O7	69.70 (10)	C19—C20—C22	120.0 (4)
O2 ^{vi} —Ce1—O7	74.76 (10)	C16—C21—C20	120.0 (4)
O1—Ce1—O7	141.32 (10)	C16—C21—H21	120.0
O1W—Ce1—O7	112.36 (9)	C20—C21—H21	120.0
O1 ^{vi} —Ce1—O7	96.35 (9)	O9—C22—O8	125.2 (4)
O9 ⁱⁱⁱ —Ce1—C1 ^{vi}	77.77 (11)	O9—C22—C20	117.9 (4)
O8 ^{iv} —Ce1—C1 ^{vi}	87.39 (12)	O8—C22—C20	116.9 (4)
O6—Ce1—C1 ^{vi}	134.58 (11)	O10—C23—N4	121.8 (5)
O3 ^v —Ce1—C1 ^{vi}	91.94 (12)	O10—C23—C24	119.4 (5)
O2 ^{vi} —Ce1—C1 ^{vi}	24.43 (11)	N4—C23—C24	118.8 (5)
O1—Ce1—C1 ^{vi}	99.20 (11)	C28—C24—C25	117.2 (5)
O1W—Ce1—C1 ^{vi}	150.89 (12)	C28—C24—C23	124.7 (5)
O1 ^{vi} —Ce1—C1 ^{vi}	25.54 (11)	C25—C24—C23	118.1 (5)
O7—Ce1—C1 ^{vi}	84.49 (10)	C26—C25—C24	120.1 (6)
O9 ⁱⁱⁱ —Ce1—Ce1 ^{vi}	70.04 (7)	C26—C25—H25	119.9
O8 ^{iv} —Ce1—Ce1 ^{vi}	63.92 (7)	C24—C25—H25	119.9
O6—Ce1—Ce1 ^{vi}	137.33 (8)	N3—C26—C25	122.7 (6)
O3 ^v —Ce1—Ce1 ^{vi}	144.99 (8)	N3—C26—H26	118.6
O2 ^{vi} —Ce1—Ce1 ^{vi}	84.17 (7)	C25—C26—H26	118.6
O1—Ce1—Ce1 ^{vi}	39.05 (7)	N3—C27—C28	121.9 (6)
O1W—Ce1—Ce1 ^{vi}	119.27 (7)	N3—C27—H27	119.1
O1 ^{vi} —Ce1—Ce1 ^{vi}	36.76 (6)	C28—C27—H27	119.1
O7—Ce1—Ce1 ^{vi}	123.63 (7)	C24—C28—C27	120.2 (5)
C1 ^{vi} —Ce1—Ce1 ^{vi}	60.81 (9)	C24—C28—H28	119.9
O2—C1—O1	122.0 (4)	C27—C28—H28	119.9
O2—C1—C2	119.2 (4)	C13—N1—C12	116.9 (5)
O1—C1—C2	118.8 (4)	C13—N1—Ag1 ⁱ	123.5 (4)

supplementary materials

O2—C1—Ce1 ^{vi}	56.7 (2)	C12—N1—Ag1 ⁱ	116.1 (3)
O1—C1—Ce1 ^{vi}	65.3 (2)	C9—N2—C6	125.4 (4)
C2—C1—Ce1 ^{vi}	175.4 (3)	C9—N2—H2	117.3
C7—C2—C3	120.4 (4)	C6—N2—H2	117.3
C7—C2—C1	119.1 (4)	C27—N3—C26	117.9 (5)
C3—C2—C1	120.4 (4)	C27—N3—Ag1	121.3 (4)
C4—C3—C2	119.2 (4)	C26—N3—Ag1	120.8 (4)
C4—C3—H3	120.4	C23—N4—C18	122.0 (4)
C2—C3—H3	120.4	C23—N4—H4	119.0
C3—C4—C5	119.6 (4)	C18—N4—H4	119.0
C3—C4—C8	120.7 (4)	C1—O1—Ce1	152.6 (3)
C5—C4—C8	119.5 (4)	C1—O1—Ce1 ^{vi}	89.1 (3)
C6—C5—C4	120.9 (4)	Ce1—O1—Ce1 ^{vi}	104.20 (10)
C6—C5—H5	119.6	C1—O2—Ce1 ^{vi}	98.9 (3)
C4—C5—H5	119.6	C8—O3—Ce1 ^{vii}	137.7 (3)
C5—C6—C7	120.0 (4)	C15—O6—Ce1	97.7 (3)
C5—C6—N2	122.7 (4)	C15—O7—Ag1 ^{viii}	119.3 (3)
C7—C6—N2	117.2 (4)	C15—O7—Ce1	87.7 (2)
C6—C7—C2	119.9 (4)	Ag1 ^{viii} —O7—Ce1	107.58 (12)
C6—C7—H7	120.1	C22—O8—Ce1 ^{iv}	131.5 (3)
C2—C7—H7	120.1	C22—O9—Ce1 ^{ix}	129.5 (3)
O3—C8—O4	124.5 (4)	Ce1—O1W—H1WB	108.0
O3—C8—C4	118.0 (4)	Ce1—O1W—H1WA	107.2
O4—C8—C4	117.4 (4)	H1WB—O1W—H1WA	107.1
O5—C9—N2	123.9 (4)		
O2—C1—C2—C7	−160.8 (4)	N1 ⁱ —Ag1—N3—C26	140.5 (4)
O1—C1—C2—C7	19.1 (6)	O7 ⁱⁱ —Ag1—N3—C26	−10.4 (5)
O2—C1—C2—C3	17.6 (7)	O10—C23—N4—C18	7.5 (8)
O1—C1—C2—C3	−162.4 (4)	C24—C23—N4—C18	−172.8 (4)
C7—C2—C3—C4	−0.3 (7)	C17—C18—N4—C23	124.9 (5)
C1—C2—C3—C4	−178.7 (4)	C19—C18—N4—C23	−52.0 (7)
C2—C3—C4—C5	−0.5 (7)	O2—C1—O1—Ce1	−118.1 (6)
C2—C3—C4—C8	173.2 (4)	C2—C1—O1—Ce1	62.0 (8)
C3—C4—C5—C6	0.5 (7)	Ce1 ^{vi} —C1—O1—Ce1	−120.2 (6)
C8—C4—C5—C6	−173.3 (4)	O2—C1—O1—Ce1 ^{vi}	2.1 (4)
C4—C5—C6—C7	0.3 (7)	C2—C1—O1—Ce1 ^{vi}	−177.8 (4)
C4—C5—C6—N2	177.7 (4)	O9 ⁱⁱⁱ —Ce1—O1—C1	−159.6 (6)
C5—C6—C7—C2	−1.0 (7)	O8 ^{iv} —Ce1—O1—C1	43.6 (6)
N2—C6—C7—C2	−178.6 (4)	O6—Ce1—O1—C1	−20.4 (6)
C3—C2—C7—C6	1.0 (7)	O3 ^v —Ce1—O1—C1	−124.1 (6)
C1—C2—C7—C6	179.4 (4)	O2 ^{vi} —Ce1—O1—C1	139.5 (6)
C3—C4—C8—O3	29.7 (7)	O1W—Ce1—O1—C1	−82.1 (6)
C5—C4—C8—O3	−156.5 (4)	O1 ^{vi} —Ce1—O1—C1	117.0 (7)
C3—C4—C8—O4	−147.4 (4)	O7—Ce1—O1—C1	34.7 (7)

C5—C4—C8—O4	26.3 (6)	C1 ^{vi} —Ce1—O1—C1	127.3 (6)
O5—C9—C10—C14	139.0 (5)	Ce1 ^{vi} —Ce1—O1—C1	117.0 (7)
N2—C9—C10—C14	−38.8 (7)	O9 ⁱⁱⁱ —Ce1—O1—Ce1 ^{vi}	83.44 (12)
O5—C9—C10—C11	−36.7 (7)	O8 ^{iv} —Ce1—O1—Ce1 ^{vi}	−73.38 (11)
N2—C9—C10—C11	145.5 (5)	O6—Ce1—O1—Ce1 ^{vi}	−137.43 (11)
C14—C10—C11—C12	−1.0 (8)	O3 ^v —Ce1—O1—Ce1 ^{vi}	118.93 (17)
C9—C10—C11—C12	174.9 (5)	O2 ^{vi} —Ce1—O1—Ce1 ^{vi}	22.50 (16)
C10—C11—C12—N1	−2.4 (10)	O1W—Ce1—O1—Ce1 ^{vi}	160.95 (12)
C11—C10—C14—C13	2.4 (8)	O1 ^{vi} —Ce1—O1—Ce1 ^{vi}	0.0
C9—C10—C14—C13	−173.4 (5)	O7—Ce1—O1—Ce1 ^{vi}	−82.34 (16)
N1—C13—C14—C10	−0.5 (10)	C1 ^{vi} —Ce1—O1—Ce1 ^{vi}	10.33 (13)
O6—C15—C16—C21	151.8 (5)	O1—C1—O2—Ce1 ^{vi}	−2.3 (5)
O7—C15—C16—C21	−28.0 (7)	C2—C1—O2—Ce1 ^{vi}	177.6 (3)
O6—C15—C16—C17	−26.0 (7)	O4—C8—O3—Ce1 ^{vii}	26.3 (8)
O7—C15—C16—C17	154.2 (5)	C4—C8—O3—Ce1 ^{vii}	−150.6 (3)
C21—C16—C17—C18	2.2 (7)	O7—C15—O6—Ce1	−13.1 (5)
C15—C16—C17—C18	180.0 (4)	C16—C15—O6—Ce1	167.0 (4)
C16—C17—C18—C19	−1.7 (8)	O9 ⁱⁱⁱ —Ce1—O6—C15	−129.0 (3)
C16—C17—C18—N4	−178.7 (4)	O8 ^{iv} —Ce1—O6—C15	84.7 (3)
C17—C18—C19—C20	0.0 (7)	O3 ^v —Ce1—O6—C15	−67.4 (3)
N4—C18—C19—C20	176.9 (5)	O2 ^{vi} —Ce1—O6—C15	−14.7 (3)
C18—C19—C20—C21	1.2 (7)	O1—Ce1—O6—C15	144.9 (3)
C18—C19—C20—C22	−175.0 (4)	O1W—Ce1—O6—C15	−144.5 (3)
C17—C16—C21—C20	−1.0 (7)	O1 ^{vi} —Ce1—O6—C15	49.7 (4)
C15—C16—C21—C20	−178.7 (4)	O7—Ce1—O6—C15	6.7 (3)
C19—C20—C21—C16	−0.7 (7)	C1 ^{vi} —Ce1—O6—C15	12.6 (4)
C22—C20—C21—C16	175.4 (4)	Ce1 ^{vi} —Ce1—O6—C15	105.9 (3)
C21—C20—C22—O9	165.3 (4)	O6—C15—O7—Ag1 ^{viii}	121.0 (4)
C19—C20—C22—O9	−18.5 (6)	C16—C15—O7—Ag1 ^{viii}	−59.2 (5)
C21—C20—C22—O8	−15.4 (7)	O6—C15—O7—Ce1	12.0 (5)
C19—C20—C22—O8	160.8 (4)	C16—C15—O7—Ce1	−168.1 (4)
O10—C23—C24—C28	137.3 (6)	O9 ⁱⁱⁱ —Ce1—O7—C15	120.7 (3)
N4—C23—C24—C28	−42.4 (8)	O8 ^{iv} —Ce1—O7—C15	−93.3 (3)
O10—C23—C24—C25	−40.9 (8)	O6—Ce1—O7—C15	−6.6 (3)
N4—C23—C24—C25	139.4 (5)	O3 ^v —Ce1—O7—C15	83.5 (3)
C28—C24—C25—C26	−0.8 (8)	O2 ^{vi} —Ce1—O7—C15	154.8 (3)
C23—C24—C25—C26	177.5 (5)	O1—Ce1—O7—C15	−84.5 (3)
C24—C25—C26—N3	−0.3 (9)	O1W—Ce1—O7—C15	22.2 (3)
C25—C24—C28—C27	1.4 (8)	O1 ^{vi} —Ce1—O7—C15	−159.7 (3)
C23—C24—C28—C27	−176.8 (5)	C1 ^{vi} —Ce1—O7—C15	177.6 (3)
N3—C27—C28—C24	−0.9 (9)	Ce1 ^{vi} —Ce1—O7—C15	−133.1 (3)
C14—C13—N1—C12	−2.7 (10)	O9 ⁱⁱⁱ —Ce1—O7—Ag1 ^{viii}	0.7 (2)

supplementary materials

C14—C13—N1—Ag1 ⁱ	154.9 (5)	O8 ^{iv} —Ce1—O7—Ag1 ^{viii}	146.62 (14)
C11—C12—N1—C13	4.2 (10)	O6—Ce1—O7—Ag1 ^{viii}	-126.60 (18)
C11—C12—N1—Ag1 ⁱ	-155.1 (5)	O3 ^v —Ce1—O7—Ag1 ^{viii}	-36.53 (12)
O5—C9—N2—C6	-7.6 (7)	O2 ^{vi} —Ce1—O7—Ag1 ^{viii}	34.77 (12)
C10—C9—N2—C6	170.2 (4)	O1—Ce1—O7—Ag1 ^{viii}	155.43 (12)
C5—C6—N2—C9	45.2 (7)	O1W—Ce1—O7—Ag1 ^{viii}	-97.83 (13)
C7—C6—N2—C9	-137.3 (5)	O1 ^{vi} —Ce1—O7—Ag1 ^{viii}	80.25 (12)
C28—C27—N3—C26	-0.2 (8)	C1 ^{vi} —Ce1—O7—Ag1 ^{viii}	57.59 (13)
C28—C27—N3—Ag1	-179.0 (4)	Ce1 ^{vi} —Ce1—O7—Ag1 ^{viii}	106.86 (10)
C25—C26—N3—C27	0.8 (9)	O9—C22—O8—Ce1 ^{iv}	65.6 (6)
C25—C26—N3—Ag1	179.6 (4)	C20—C22—O8—Ce1 ^{iv}	-113.6 (4)
N1 ⁱ —Ag1—N3—C27	-40.7 (5)	O8—C22—O9—Ce1 ^{ix}	-41.8 (6)
O7 ⁱⁱ —Ag1—N3—C27	168.3 (4)	C20—C22—O9—Ce1 ^{ix}	137.4 (3)


Symmetry codes: (i) $-x+2, -y+1, -z+3$; (ii) $x, y, z+1$; (iii) $x, y+1, z$; (iv) $-x+1, -y, -z+1$; (v) $x+1, y, z$; (vi) $-x+1, -y+1, -z+1$; (vii) $x-1, y, z$; (viii) $x, y, z-1$; (ix) $x, y-1, z$.

Hydrogen-bond geometry (\AA , $^\circ$)

$D\text{—H}\cdots A$	$D\text{—H}$	$H\cdots A$	$D\cdots A$	$D\text{—H}\cdots A$
O1W—H1WA \cdots O5 ^x	0.85	2.08	2.836 (5)	147
O1W—H1WB \cdots O4 ^v	0.85	2.01	2.691 (6)	137
N2—H2 \cdots O10 ⁱⁱⁱ	0.86	2.01	2.788 (7)	149
N4—H4 \cdots O4 ^x	0.86	2.08	2.926 (6)	167

Symmetry codes: (x) $-x+1, -y+1, -z+2$; (v) $x+1, y, z$; (iii) $x, y+1, z$.

Fig. 1



supplementary materials

Fig. 2

