

Tetraethylammonium tri- μ -phenolato-bis[tricarbonylmanganate(I)]

 Timothy J. McNeese^{a*} and Robert D. Pike^b

^aDepartment of Chemistry, Loyola University Maryland, 4501 N. Charles Street, Baltimore, MD 21210-2699, USA, and ^bDepartment of Chemistry, College of William and Mary, PO Box 8795, Williamsburg, VA 23187-8795, USA

Correspondence e-mail: tmcneese@loyola.edu


Received 15 July 2011; accepted 19 July 2011

Key indicators: single-crystal X-ray study; $T = 100$ K; mean $\sigma(\text{C}-\text{C}) = 0.004$ Å; R factor = 0.033; wR factor = 0.084; data-to-parameter ratio = 12.0.

The title compound, $(\text{C}_8\text{H}_{20}\text{N})[\text{Mn}_2(\text{C}_6\text{H}_5\text{O})_3(\text{CO})_6]$, was synthesized from $[\text{Mn}(\text{CO})_3(\text{CH}_3\text{CN})_3]\text{BF}_4$ and $(\text{C}_8\text{H}_{20}\text{N})-(\text{OC}_6\text{H}_5)$. The binuclear anion exhibits a pseudo-threefold symmetry and contains two six-coordinate Mn atoms. Each metal atom is coordinated by three facially oriented CO ligands and three doubly-bridging phenolate ligands. The average O–Mn–O bond angle is $74.9(7)^\circ$ in the Mn_2O_3 metal–phenolate dimeric core, yielding a distorted octahedron for each metal.

Related literature

For the synthesis of the starting materials, see: Riemann & Singleton (1973); McNeese *et al.* (1985). For related metal phenolate complexes, see: Darensbourg *et al.* (1988, 1989); McNeese *et al.* (1985); Lee *et al.* (1995). For analogous tungsten and rhenium dimers, see: Darensbourg *et al.* (1988); Beringhelli *et al.* (1985).



Experimental

Crystal data

 $(\text{C}_8\text{H}_{20}\text{N})[\text{Mn}_2(\text{C}_6\text{H}_5\text{O})_3(\text{CO})_6]$
 $M_r = 687.49$

Orthorhombic, $Pna2_1$
 $a = 18.6831(4)$ Å
 $b = 9.2037(2)$ Å
 $c = 18.5999(4)$ Å
 $V = 3198.32(12)$ Å³

$Z = 4$
 Cu $K\alpha$ radiation
 $\mu = 6.88$ mm⁻¹
 $T = 100$ K
 $0.45 \times 0.43 \times 0.30$ mm

Data collection

Bruker SMART APEXII CCD diffractometer
 Absorption correction: numerical (SADABS; Sheldrick, 2004)
 $T_{\min} = 0.148$, $T_{\max} = 0.229$

30613 measured reflections
 4817 independent reflections
 4660 reflections with $I > 2\sigma(I)$
 $R_{\text{int}} = 0.059$

Refinement

$R[F^2 > 2\sigma(F^2)] = 0.033$
 $wR(F^2) = 0.084$
 $S = 1.07$
 4817 reflections
 402 parameters
 19 restraints

H-atom parameters constrained
 $\Delta\rho_{\max} = 0.79$ e Å⁻³
 $\Delta\rho_{\min} = -0.57$ e Å⁻³
 Absolute structure: Flack (1983), 1874 Friedel pairs
 Flack parameter: 0.051 (5)

Data collection: APEX2 (Bruker, 2004); cell refinement: SAINT-Plus (Bruker, 2004); data reduction: SAINT-Plus; program(s) used to solve structure: SHELXS97 (Sheldrick, 2008); program(s) used to refine structure: SHELXL97 (Sheldrick, 2008); molecular graphics: ORTEP-3 (Farrugia, 1997) and Mercury (Macrae *et al.*, 2006); software used to prepare material for publication: SHELXL97.

TJM acknowledges Daniel M. Perrine for technical assistance and Loyola University Maryland for financial support. RDP thanks the NSF (CHE-0443345) and the College of William and Mary for the purchase of the X-ray equipment.

Supplementary data and figures for this paper are available from the IUCr electronic archives (Reference: BG2415).

References

- Beringhelli, T., Gianfranco, C., Giuseppe, A., Sironi, A. & Freni, M. (1985). *J. Chem. Soc. Dalton Trans.* pp. 1507–1512.
- Bruker (2004). APEX2 and SAINT-Plus. Bruker AXS Inc., Madison, Wisconsin, USA.
- Darensbourg, D. J., Sanchez, K. M. & Reibenspies, J. H. (1988). *Inorg. Chem.* **27**, 3269–3270.
- Darensbourg, D. J., Sanchez, K. M., Reibenspies, J. H. & Rheingold, A. L. (1989). *J. Am. Chem. Soc.* **111**, 7094–7103.
- Farrugia, L. J. (1997). *J. Appl. Cryst.* **30**, 565.
- Flack, H. D. (1983). *Acta Cryst.* **A39**, 876–881.
- Lee, S.-G., Kim, J.-A., Chung, Y. K., Yoon, T.-S., Kim, N. & Whanchul, S. (1995). *Organometallics*, **14**, 1023–1029.
- Macrae, C. F., Edgington, P. R., McCabe, P., Pidcock, E., Shields, G. P., Taylor, R., Towler, M. & van de Streek, J. (2006). *J. Appl. Cryst.* **39**, 453–457.
- McNeese, T. J., Mueller, T. E., Wierda, D. A., Darensbourg, D. J. & DeLord, T. J. (1985). *Inorg. Chem.* **24**, 3465–3468.
- Riemann, R. H. & Singleton, E. (1973). *J. Organomet. Chem.* **59**, C24–C26.
- Sheldrick, G. M. (2004). SADABS. University of Göttingen, Germany.
- Sheldrick, G. M. (2008). *Acta Cryst.* **A64**, 112–122.

supplementary materials

Acta Cryst. (2011). E67, m1150 [doi:10.1107/S1600536811029266]

Tetraethylammonium tri- μ -phenolato-bis[tricarbonylmanganate(I)]

T. J. McNeese and R. D. Pike

Comment

The phenolate ligand, OC_6H_5^- , exhibits a variety of bonding modes in organometallic carbonyl compounds. Examples include mono- and polynuclear complexes where the ligand is bonded by its oxygen atom to one or more metals in a terminal (Darensbourg *et al.*, 1989), doubly- (Darensbourg *et al.*, 1988) and triply- (McNeese, *et al.*, 1985) bridging fashion, and through its aromatic ring as an oxocyclohexadienyl ligand (Lee *et al.*, 1995).

The title compound, $(\text{Et}_4\text{N})[(\text{CO})_3\text{Mn}(\mu\text{-OC}_6\text{H}_5)_3\text{Mn}(\text{CO})_3]$ (**1**), was synthesized by displacement of labile acetonitrile ligands of the precursor, $[\text{Mn}(\text{CO})_3(\text{CH}_3\text{CN})_3]\text{BF}_4$ (Riemann & Singleton, 1973), by phenoxide ions of $(\text{Et}_4\text{N})(\text{OC}_6\text{H}_5)$ (McNeese *et al.*, 1985). The doubly-bridging phenolate ligands of the dimer are basic, reaction of (**1**) in CH_3CN with HBF_4 re-forms the Mn-containing starting material and $\text{C}_6\text{H}_5\text{OH}$.

The bimetallic compound crystallizes in the orthorhombic space group $\text{Pna}2(1)$, and exhibits a striking pseudo threefold symmetry. The structure presents bond lengths and angles that are comparable to analogous Re (Beringhelli *et al.*, 1985) and W (Darensbourg *et al.*, 1988) dimers. The geometry for each Mn atom in the organometallic anion of (**1**) is a distorted octahedron (Figure 1), with average O—Mn—O angles in the Mn_2O_3 metal-phenolate group of $75.12(7)^\circ$. Each Mn atom has a noble-gas configuration and a nonbonding metal distance of $2.8759(5)$ Å. Figure 2 illustrates the packing diagram for the molecular structure of the title compound.

Experimental

Solid $(\text{Et}_4\text{N})(\text{OC}_6\text{H}_5)$ (500 mg, 2.24 mmol) (McNeese *et al.*, 1985) was added with a funnel to a 100-ml Schlenk tube containing a 50-ml CH_3CN solution of $[\text{Mn}(\text{CO})_3(\text{CH}_3\text{CN})_3]\text{BF}_4$ (521 mg, 1.49 mmol) (Riemann & Singleton, 1973). The orange-colored solution was stirred under argon at room temperature for 18 h and evaporated to dryness. THF (25 ml) was added to the solid to dissolve the desired compound. The reaction mixture was filtered under vacuum to separate insoluble Et_4NBF_4 and the resulting solution was evaporated. The orange-colored product, $(\text{Et}_4\text{N})[(\text{CO})_3\text{Mn}(\mu\text{-OC}_6\text{H}_5)_3\text{Mn}(\text{CO})_3]$, was crystallized from THF/cyclohexane. Yield: 359 mg (69%). IR ($\nu(\text{CO})$, CH_3CN) 2013 (s), 1912 (s) cm^{-1} ; ^1H NMR (CD_3CN): cation, δ 1.25 (12H, t, $-\text{CH}_3$), 3.25 (8H, q, $-\text{CH}_2-$); anion, δ 6.75–7.45 (15H, m, OC_6H_5). Elemental analysis, calcd for $\text{C}_{32}\text{H}_{35}\text{Mn}_2\text{NO}_9$: C, 55.90; H, 5.14. Found: C, 55.67; H, 5.28.

Refinement

All hydrogen atoms were placed in theoretical positions (C—H: 0.95–0.99 Å) riding on the atoms to which they are attached.

Displacement factors of the atoms attached to Mn were restrained *via* DELU instructions in *SHELXL*.

Figures




Fig. 1. Ellipsoid plot of the organometallic anion in (1). Displacement ellipsoids are drawn at the 50% probability level.




Fig. 2. Packing diagram of (1) viewed down the *b* axis.

Tetraethylammonium tri- μ -phenolato-bis[tricarboxylmanganate(I)]

Crystal data

(C₈H₂₀N)[Mn₂(C₆H₅O)₃(CO)₆]

$M_r = 687.49$

Orthorhombic, *Pna*2₁

Hall symbol: P 2c -2n

$a = 18.6831$ (4) Å

$b = 9.2037$ (2) Å

$c = 18.5999$ (4) Å

$V = 3198.32$ (12) Å³

$Z = 4$

$F(000) = 1424$

$D_x = 1.428$ Mg m⁻³

Cu $K\alpha$ radiation, $\lambda = 1.54178$ Å

Cell parameters from 9989 reflections

$\theta = 3.4$ – 68.1°

$\mu = 6.88$ mm⁻¹

$T = 100$ K

Block, brown

$0.45 \times 0.43 \times 0.30$ mm

Data collection

Bruker SMART APEXII CCD
diffractometer

Radiation source: fine-focus sealed tube
graphite

ω and ψ scans

Absorption correction: numerical
(*SADABS*; Sheldrick, 2004)

$T_{\min} = 0.148$, $T_{\max} = 0.229$

30613 measured reflections

4817 independent reflections

4660 reflections with $I > 2\sigma(I)$

$R_{\text{int}} = 0.059$

$\theta_{\max} = 67.0^\circ$, $\theta_{\min} = 4.7^\circ$

$h = -22 \rightarrow 22$

$k = -10 \rightarrow 10$

$l = -19 \rightarrow 22$

Refinement

Refinement on F^2

Secondary atom site location: difference Fourier map

Least-squares matrix: full	Hydrogen site location: inferred from neighbouring sites
$R[F^2 > 2\sigma(F^2)] = 0.033$	H-atom parameters constrained
$wR(F^2) = 0.084$	$w = 1/[\sigma^2(F_o^2) + (0.0591P)^2]$
$S = 1.07$	where $P = (F_o^2 + 2F_c^2)/3$
4817 reflections	$(\Delta/\sigma)_{\max} = 0.001$
402 parameters	$\Delta\rho_{\max} = 0.79 \text{ e } \text{\AA}^{-3}$
19 restraints	$\Delta\rho_{\min} = -0.57 \text{ e } \text{\AA}^{-3}$
Primary atom site location: structure-invariant direct methods	Absolute structure: Flack (1983), 1874 Friedel pairs
	Flack parameter: 0.051 (5)

Special details

Geometry. All e.s.d.'s (except the e.s.d. in the dihedral angle between two l.s. planes) are estimated using the full covariance matrix. The cell e.s.d.'s are taken into account individually in the estimation of e.s.d.'s in distances, angles and torsion angles; correlations between e.s.d.'s in cell parameters are only used when they are defined by crystal symmetry. An approximate (isotropic) treatment of cell e.s.d.'s is used for estimating e.s.d.'s involving l.s. planes.

Refinement. Refinement of F^2 against ALL reflections. The weighted R -factor wR and goodness of fit S are based on F^2 , conventional R -factors R are based on F , with F set to zero for negative F^2 . The threshold expression of $F^2 > \sigma(F^2)$ is used only for calculating R -factors(gt) *etc.* and is not relevant to the choice of reflections for refinement. R -factors based on F^2 are statistically about twice as large as those based on F , and R -factors based on ALL data will be even larger.

Fractional atomic coordinates and isotropic or equivalent isotropic displacement parameters (\AA^2)

	x	y	z	$U_{\text{iso}}^*/U_{\text{eq}}$
Mn1	0.942524 (18)	0.23445 (4)	0.69042 (3)	0.00987 (11)
Mn2	0.864233 (19)	0.49426 (4)	0.72511 (3)	0.00854 (11)
O1	0.88111 (12)	-0.0575 (2)	0.71027 (13)	0.0277 (5)
O2	1.00084 (11)	0.1572 (2)	0.54831 (12)	0.0255 (5)
O3	1.07657 (10)	0.1197 (2)	0.75028 (13)	0.0253 (5)
O4	0.71971 (11)	0.5205 (2)	0.78704 (13)	0.0246 (5)
O5	0.82718 (11)	0.7131 (2)	0.61509 (12)	0.0197 (4)
O6	0.90503 (11)	0.7115 (2)	0.83376 (12)	0.0194 (4)
O7	0.96310 (8)	0.45028 (19)	0.68333 (11)	0.0120 (4)
O8	0.89854 (9)	0.3187 (2)	0.78004 (11)	0.0139 (4)
O9	0.85015 (9)	0.32527 (19)	0.65656 (11)	0.0118 (4)
C1	0.90466 (14)	0.0552 (3)	0.70182 (17)	0.0151 (5)
C2	0.97751 (12)	0.1871 (3)	0.60351 (16)	0.0134 (5)
C3	1.02446 (13)	0.1687 (3)	0.72812 (18)	0.0160 (5)
C4	0.77549 (14)	0.5058 (3)	0.76187 (18)	0.0133 (5)
C5	0.84219 (13)	0.6313 (3)	0.65873 (15)	0.0114 (5)
C6	0.89061 (13)	0.6281 (3)	0.79007 (15)	0.0110 (5)
C7	1.02389 (13)	0.5276 (3)	0.67916 (14)	0.0101 (5)
C8	1.08961 (14)	0.4598 (3)	0.66590 (16)	0.0156 (6)
H8	1.0915	0.3575	0.6593	0.019*
C9	1.15232 (15)	0.5415 (4)	0.66233 (17)	0.0191 (6)

supplementary materials

H9	1.1965	0.4939	0.6531	0.023*
C10	1.15143 (15)	0.6886 (3)	0.67177 (17)	0.0212 (6)
H10	1.1947	0.7428	0.6708	0.025*
C11	1.08608 (16)	0.7580 (3)	0.68280 (19)	0.0196 (6)
H11	1.0847	0.8606	0.6881	0.024*
C12	1.02274 (13)	0.6785 (3)	0.68610 (16)	0.0141 (5)
H12	0.9785	0.7273	0.6931	0.017*
C13	0.89238 (14)	0.2703 (3)	0.84769 (16)	0.0134 (6)
C14	0.92658 (15)	0.1421 (3)	0.86978 (17)	0.0186 (6)
H14	0.9543	0.0884	0.8361	0.022*
C15	0.92062 (16)	0.0925 (3)	0.93968 (18)	0.0232 (6)
H15	0.9441	0.0052	0.9532	0.028*
C16	0.88101 (18)	0.1679 (3)	0.99032 (18)	0.0244 (7)
H16	0.8772	0.1331	1.0382	0.029*
C17	0.84700 (17)	0.2953 (3)	0.96968 (17)	0.0209 (6)
H17	0.8199	0.3487	1.0039	0.025*
C18	0.85229 (14)	0.3455 (3)	0.89947 (16)	0.0156 (6)
H18	0.8283	0.4325	0.8862	0.019*
C19	0.79925 (13)	0.2836 (3)	0.60953 (16)	0.0104 (5)
C20	0.81015 (12)	0.1642 (3)	0.56366 (16)	0.0137 (6)
H20	0.8535	0.1106	0.5666	0.016*
C21	0.75808 (14)	0.1237 (3)	0.51403 (16)	0.0176 (6)
H21	0.7666	0.0431	0.4833	0.021*
C22	0.69416 (14)	0.1987 (4)	0.50854 (18)	0.0193 (6)
H22	0.6588	0.1708	0.4745	0.023*
C23	0.68308 (13)	0.3167 (3)	0.55445 (17)	0.0175 (6)
H23	0.6393	0.3689	0.5518	0.021*
C24	0.73455 (14)	0.3590 (3)	0.60368 (16)	0.0143 (5)
H24	0.7258	0.4404	0.6339	0.017*
N1	0.62259 (13)	0.0037 (2)	0.90791 (16)	0.0168 (5)
C25	0.70244 (15)	-0.0258 (4)	0.91741 (19)	0.0243 (7)
H25A	0.7224	0.0490	0.9500	0.029*
H25B	0.7082	-0.1212	0.9413	0.029*
C26	0.74587 (17)	-0.0265 (4)	0.8490 (3)	0.0383 (9)
H26A	0.7438	0.0698	0.8265	0.057*
H26B	0.7263	-0.0990	0.8158	0.057*
H26C	0.7957	-0.0506	0.8601	0.057*
C27	0.58925 (14)	-0.1050 (3)	0.85663 (16)	0.0168 (6)
H27A	0.5380	-0.0803	0.8506	0.020*
H27B	0.6125	-0.0945	0.8091	0.020*
C28	0.59471 (18)	-0.2619 (3)	0.87963 (19)	0.0251 (7)
H28A	0.5685	-0.2757	0.9247	0.038*
H28B	0.6451	-0.2875	0.8868	0.038*
H28C	0.5741	-0.3243	0.8423	0.038*
C29	0.58983 (18)	-0.0064 (3)	0.98216 (18)	0.0214 (7)
H29A	0.6006	-0.1036	1.0023	0.026*
H29B	0.6131	0.0666	1.0135	0.026*
C30	0.5099 (2)	0.0171 (4)	0.9847 (2)	0.0307 (8)
H30A	0.4862	-0.0537	0.9534	0.046*

H30B	0.4988	0.1157	0.9681	0.046*
H30C	0.4929	0.0048	1.0341	0.046*
C31	0.61045 (16)	0.1516 (3)	0.87507 (17)	0.0205 (6)
H31A	0.5585	0.1640	0.8669	0.025*
H31B	0.6342	0.1539	0.8275	0.025*
C32	0.6371 (2)	0.2799 (4)	0.9187 (2)	0.0340 (9)
H32A	0.6888	0.2707	0.9262	0.051*
H32B	0.6127	0.2817	0.9653	0.051*
H32C	0.6271	0.3702	0.8927	0.051*

Atomic displacement parameters (\AA^2)

	U^{11}	U^{22}	U^{33}	U^{12}	U^{13}	U^{23}
Mn1	0.00445 (18)	0.0091 (2)	0.0160 (2)	0.00351 (13)	-0.00178 (18)	-0.00336 (17)
Mn2	0.00319 (18)	0.0078 (2)	0.0146 (2)	0.00217 (13)	0.00009 (17)	-0.00081 (15)
O1	0.0346 (11)	0.0128 (8)	0.0358 (15)	-0.0038 (8)	-0.0064 (10)	0.0000 (9)
O2	0.0132 (9)	0.0400 (13)	0.0232 (9)	0.0042 (9)	0.0009 (8)	-0.0140 (9)
O3	0.0156 (8)	0.0239 (11)	0.0364 (13)	0.0116 (7)	-0.0130 (9)	-0.0079 (9)
O4	0.0079 (8)	0.0358 (13)	0.0300 (14)	0.0031 (8)	0.0050 (8)	-0.0003 (9)
O5	0.0188 (10)	0.0174 (10)	0.0230 (11)	0.0053 (8)	0.0002 (9)	0.0044 (7)
O6	0.0192 (9)	0.0179 (10)	0.0211 (11)	-0.0012 (8)	0.0008 (9)	-0.0062 (7)
O7	0.0040 (7)	0.0110 (9)	0.0209 (11)	0.0000 (6)	0.0017 (8)	-0.0027 (8)
O8	0.0139 (8)	0.0117 (9)	0.0160 (10)	0.0058 (7)	0.0018 (8)	-0.0004 (7)
O9	0.0052 (8)	0.0104 (9)	0.0198 (11)	0.0020 (7)	-0.0046 (7)	-0.0042 (7)
C1	0.0124 (12)	0.0101 (9)	0.0227 (15)	0.0045 (8)	-0.0028 (11)	-0.0053 (11)
C2	0.0025 (11)	0.0164 (13)	0.0213 (11)	0.0031 (10)	-0.0014 (10)	-0.0057 (11)
C3	0.0102 (10)	0.0131 (13)	0.0249 (15)	0.0047 (9)	-0.0060 (10)	-0.0064 (11)
C4	0.0063 (9)	0.0131 (14)	0.0203 (16)	0.0024 (9)	-0.0003 (10)	0.0008 (10)
C5	0.0054 (11)	0.0101 (12)	0.0188 (13)	0.0024 (9)	0.0024 (10)	-0.0003 (8)
C6	0.0055 (10)	0.0106 (13)	0.0170 (13)	0.0016 (9)	-0.0002 (10)	0.0003 (8)
C7	0.0066 (11)	0.0168 (12)	0.0068 (13)	-0.0014 (10)	-0.0004 (10)	-0.0002 (10)
C8	0.0110 (13)	0.0190 (14)	0.0167 (15)	0.0005 (11)	-0.0013 (11)	-0.0035 (11)
C9	0.0068 (12)	0.0344 (17)	0.0160 (15)	0.0012 (12)	0.0035 (11)	-0.0031 (13)
C10	0.0102 (11)	0.0330 (17)	0.0204 (16)	-0.0083 (12)	0.0036 (12)	-0.0015 (12)
C11	0.0194 (14)	0.0174 (14)	0.0221 (18)	-0.0047 (10)	0.0044 (14)	-0.0019 (11)
C12	0.0075 (11)	0.0183 (13)	0.0165 (14)	0.0010 (9)	0.0039 (11)	-0.0007 (11)
C13	0.0109 (13)	0.0122 (14)	0.0169 (15)	-0.0022 (10)	-0.0018 (12)	-0.0015 (10)
C14	0.0197 (13)	0.0147 (14)	0.0214 (16)	0.0042 (11)	-0.0005 (12)	-0.0009 (11)
C15	0.0307 (15)	0.0172 (15)	0.0216 (16)	0.0063 (12)	-0.0041 (14)	0.0053 (12)
C16	0.0328 (16)	0.0238 (17)	0.0168 (15)	0.0005 (13)	-0.0025 (14)	0.0069 (12)
C17	0.0222 (14)	0.0193 (15)	0.0212 (16)	-0.0007 (12)	0.0021 (13)	-0.0026 (12)
C18	0.0130 (12)	0.0158 (14)	0.0179 (15)	0.0036 (10)	-0.0014 (11)	0.0000 (11)
C19	0.0029 (11)	0.0133 (13)	0.0150 (14)	-0.0030 (9)	-0.0007 (10)	0.0032 (10)
C20	0.0062 (11)	0.0155 (14)	0.0195 (15)	-0.0023 (9)	0.0007 (11)	0.0004 (11)
C21	0.0153 (13)	0.0180 (14)	0.0195 (15)	-0.0073 (11)	-0.0003 (12)	-0.0015 (11)
C22	0.0105 (13)	0.0271 (16)	0.0203 (16)	-0.0091 (11)	-0.0057 (11)	0.0040 (12)
C23	0.0042 (11)	0.0205 (15)	0.0277 (18)	-0.0022 (10)	-0.0043 (11)	0.0078 (12)
C24	0.0075 (11)	0.0140 (13)	0.0213 (15)	-0.0006 (10)	0.0011 (11)	0.0016 (11)

supplementary materials

N1	0.0099 (11)	0.0202 (14)	0.0203 (14)	-0.0005 (9)	-0.0044 (11)	0.0018 (9)
C25	0.0109 (14)	0.0270 (16)	0.035 (2)	0.0028 (12)	-0.0101 (14)	0.0021 (13)
C26	0.0109 (15)	0.055 (2)	0.049 (3)	0.0007 (14)	-0.0023 (16)	0.0094 (18)
C27	0.0106 (12)	0.0254 (16)	0.0145 (15)	-0.0006 (11)	-0.0020 (11)	-0.0018 (11)
C28	0.0275 (16)	0.0240 (17)	0.0239 (18)	-0.0002 (12)	0.0005 (15)	-0.0023 (12)
C29	0.0290 (17)	0.0224 (17)	0.0129 (16)	0.0028 (12)	-0.0032 (14)	0.0020 (10)
C30	0.034 (2)	0.035 (2)	0.0235 (19)	0.0127 (15)	0.0101 (16)	0.0052 (13)
C31	0.0185 (13)	0.0220 (16)	0.0210 (16)	0.0006 (11)	-0.0055 (13)	0.0064 (12)
C32	0.040 (2)	0.0233 (18)	0.039 (2)	-0.0032 (14)	-0.0147 (17)	0.0053 (15)

Geometric parameters (Å, °)

Mn1—C3	1.789 (3)	C17—H17	0.9500
Mn1—C2	1.797 (3)	C18—H18	0.9500
Mn1—C1	1.807 (3)	C19—C24	1.398 (4)
Mn1—O8	2.014 (2)	C19—C20	1.406 (4)
Mn1—O9	2.0184 (18)	C20—C21	1.392 (4)
Mn1—O7	2.0276 (18)	C20—H20	0.9500
Mn1—Mn2	2.8765 (5)	C21—C22	1.383 (4)
Mn2—C6	1.795 (3)	C21—H21	0.9500
Mn2—C4	1.796 (3)	C22—C23	1.397 (5)
Mn2—C5	1.812 (3)	C22—H22	0.9500
Mn2—O8	2.017 (2)	C23—C24	1.384 (4)
Mn2—O9	2.0284 (19)	C23—H23	0.9500
Mn2—O7	2.0444 (17)	C24—H24	0.9500
O1—C1	1.138 (4)	N1—C31	1.509 (4)
O2—C2	1.149 (4)	N1—C29	1.514 (4)
O3—C3	1.149 (3)	N1—C27	1.516 (4)
O4—C4	1.151 (4)	N1—C25	1.526 (4)
O5—C5	1.142 (3)	C25—C26	1.510 (6)
O6—C6	1.150 (3)	C25—H25A	0.9900
O7—C7	1.342 (3)	C25—H25B	0.9900
O8—C13	1.340 (4)	C26—H26A	0.9800
O9—C19	1.348 (3)	C26—H26B	0.9800
C7—C12	1.395 (4)	C26—H26C	0.9800
C7—C8	1.399 (4)	C27—C28	1.509 (4)
C8—C9	1.394 (4)	C27—H27A	0.9900
C8—H8	0.9500	C27—H27B	0.9900
C9—C10	1.365 (5)	C28—H28A	0.9800
C9—H9	0.9500	C28—H28B	0.9800
C10—C11	1.393 (4)	C28—H28C	0.9800
C10—H10	0.9500	C29—C30	1.509 (5)
C11—C12	1.393 (4)	C29—H29A	0.9900
C11—H11	0.9500	C29—H29B	0.9900
C12—H12	0.9500	C30—H30A	0.9800
C13—C14	1.404 (4)	C30—H30B	0.9800
C13—C18	1.403 (4)	C30—H30C	0.9800
C14—C15	1.382 (4)	C31—C32	1.517 (5)
C14—H14	0.9500	C31—H31A	0.9900

C15—C16	1.384 (5)	C31—H31B	0.9900
C15—H15	0.9500	C32—H32A	0.9800
C16—C17	1.388 (5)	C32—H32B	0.9800
C16—H16	0.9500	C32—H32C	0.9800
C17—C18	1.389 (4)		
C3—Mn1—C2	87.67 (13)	C16—C15—H15	119.4
C3—Mn1—C1	88.86 (12)	C15—C16—C17	118.7 (3)
C2—Mn1—C1	91.51 (13)	C15—C16—H16	120.7
C3—Mn1—O8	98.90 (12)	C17—C16—H16	120.7
C2—Mn1—O8	170.52 (11)	C18—C17—C16	120.6 (3)
C1—Mn1—O8	95.42 (11)	C18—C17—H17	119.7
C3—Mn1—O9	173.68 (11)	C16—C17—H17	119.7
C2—Mn1—O9	97.52 (10)	C17—C18—C13	121.3 (3)
C1—Mn1—O9	94.59 (10)	C17—C18—H18	119.3
O8—Mn1—O9	75.52 (8)	C13—C18—H18	119.3
C3—Mn1—O7	101.23 (10)	O9—C19—C24	121.3 (2)
C2—Mn1—O7	96.34 (11)	O9—C19—C20	120.9 (2)
C1—Mn1—O7	167.43 (9)	C24—C19—C20	117.8 (2)
O8—Mn1—O7	75.76 (8)	C21—C20—C19	120.7 (2)
O9—Mn1—O7	74.70 (7)	C21—C20—H20	119.7
C3—Mn1—Mn2	128.92 (9)	C19—C20—H20	119.7
C2—Mn1—Mn2	126.05 (9)	C22—C21—C20	121.3 (3)
C1—Mn1—Mn2	122.23 (8)	C22—C21—H21	119.4
O8—Mn1—Mn2	44.50 (6)	C20—C21—H21	119.4
O9—Mn1—Mn2	44.84 (5)	C21—C22—C23	118.1 (3)
O7—Mn1—Mn2	45.30 (5)	C21—C22—H22	121.0
C6—Mn2—C4	87.53 (12)	C23—C22—H22	121.0
C6—Mn2—C5	92.48 (12)	C24—C23—C22	121.4 (2)
C4—Mn2—C5	90.49 (12)	C24—C23—H23	119.3
C6—Mn2—O8	96.99 (10)	C22—C23—H23	119.3
C4—Mn2—O8	98.50 (10)	C23—C24—C19	120.8 (3)
C5—Mn2—O8	167.19 (10)	C23—C24—H24	119.6
C6—Mn2—O9	170.16 (9)	C19—C24—H24	119.6
C4—Mn2—O9	99.48 (10)	C31—N1—C29	111.4 (2)
C5—Mn2—O9	94.35 (11)	C31—N1—C27	106.2 (2)
O8—Mn2—O9	75.24 (8)	C29—N1—C27	111.5 (2)
C6—Mn2—O7	98.26 (10)	C31—N1—C25	110.7 (2)
C4—Mn2—O7	171.95 (10)	C29—N1—C25	106.2 (2)
C5—Mn2—O7	94.83 (10)	C27—N1—C25	110.9 (2)
O8—Mn2—O7	75.33 (7)	C26—C25—N1	115.4 (3)
O9—Mn2—O7	74.12 (7)	C26—C25—H25A	108.4
C6—Mn2—Mn1	125.60 (8)	N1—C25—H25A	108.4
C4—Mn2—Mn1	127.13 (8)	C26—C25—H25B	108.4
C5—Mn2—Mn1	122.77 (9)	N1—C25—H25B	108.4
O8—Mn2—Mn1	44.42 (6)	H25A—C25—H25B	107.5
O9—Mn2—Mn1	44.56 (5)	C25—C26—H26A	109.5
O7—Mn2—Mn1	44.82 (5)	C25—C26—H26B	109.5
C7—O7—Mn1	133.11 (15)	H26A—C26—H26B	109.5
C7—O7—Mn2	132.96 (16)	C25—C26—H26C	109.5

supplementary materials

Mn1—O7—Mn2	89.89 (7)	H26A—C26—H26C	109.5
C13—O8—Mn1	133.21 (17)	H26B—C26—H26C	109.5
C13—O8—Mn2	135.63 (17)	C28—C27—N1	115.2 (2)
Mn1—O8—Mn2	91.07 (9)	C28—C27—H27A	108.5
C19—O9—Mn1	133.46 (16)	N1—C27—H27A	108.5
C19—O9—Mn2	135.87 (16)	C28—C27—H27B	108.5
Mn1—O9—Mn2	90.60 (8)	N1—C27—H27B	108.5
O1—C1—Mn1	178.8 (3)	H27A—C27—H27B	107.5
O2—C2—Mn1	179.0 (2)	C27—C28—H28A	109.5
O3—C3—Mn1	176.3 (2)	C27—C28—H28B	109.5
O4—C4—Mn2	176.2 (3)	H28A—C28—H28B	109.5
O5—C5—Mn2	177.1 (2)	C27—C28—H28C	109.5
O6—C6—Mn2	176.8 (2)	H28A—C28—H28C	109.5
O7—C7—C12	120.6 (2)	H28B—C28—H28C	109.5
O7—C7—C8	121.1 (2)	C30—C29—N1	114.8 (3)
C12—C7—C8	118.3 (2)	C30—C29—H29A	108.6
C9—C8—C7	120.4 (3)	N1—C29—H29A	108.6
C9—C8—H8	119.8	C30—C29—H29B	108.6
C7—C8—H8	119.8	N1—C29—H29B	108.6
C10—C9—C8	121.2 (3)	H29A—C29—H29B	107.5
C10—C9—H9	119.4	C29—C30—H30A	109.5
C8—C9—H9	119.4	C29—C30—H30B	109.5
C9—C10—C11	119.0 (3)	H30A—C30—H30B	109.5
C9—C10—H10	120.5	C29—C30—H30C	109.5
C11—C10—H10	120.5	H30A—C30—H30C	109.5
C12—C11—C10	120.7 (3)	H30B—C30—H30C	109.5
C12—C11—H11	119.7	N1—C31—C32	115.9 (3)
C10—C11—H11	119.7	N1—C31—H31A	108.3
C11—C12—C7	120.4 (2)	C32—C31—H31A	108.3
C11—C12—H12	119.8	N1—C31—H31B	108.3
C7—C12—H12	119.8	C32—C31—H31B	108.3
O8—C13—C14	121.0 (3)	H31A—C31—H31B	107.4
O8—C13—C18	121.8 (2)	C31—C32—H32A	109.5
C14—C13—C18	117.2 (3)	C31—C32—H32B	109.5
C15—C14—C13	121.1 (3)	H32A—C32—H32B	109.5
C15—C14—H14	119.5	C31—C32—H32C	109.5
C13—C14—H14	119.5	H32A—C32—H32C	109.5
C14—C15—C16	121.2 (3)	H32B—C32—H32C	109.5
C14—C15—H15	119.4		
C3—Mn1—Mn2—C6	1.44 (17)	C5—Mn2—O8—Mn1	1.9 (5)
C2—Mn1—Mn2—C6	-120.10 (14)	O9—Mn2—O8—Mn1	38.38 (7)
C1—Mn1—Mn2—C6	120.06 (16)	O7—Mn2—O8—Mn1	-38.70 (7)
O8—Mn1—Mn2—C6	58.90 (13)	C2—Mn1—O9—C19	47.5 (3)
O9—Mn1—Mn2—C6	-179.94 (14)	C1—Mn1—O9—C19	-44.6 (3)
O7—Mn1—Mn2—C6	-61.99 (13)	O8—Mn1—O9—C19	-139.1 (2)
C3—Mn1—Mn2—C4	-116.94 (19)	O7—Mn1—O9—C19	142.1 (2)
C2—Mn1—Mn2—C4	121.51 (16)	Mn2—Mn1—O9—C19	-177.3 (3)
C1—Mn1—Mn2—C4	1.68 (18)	C2—Mn1—O9—Mn2	-135.16 (10)
O8—Mn1—Mn2—C4	-59.49 (15)	C1—Mn1—O9—Mn2	132.70 (11)

O9—Mn1—Mn2—C4	61.67 (15)	O8—Mn1—O9—Mn2	38.28 (7)
O7—Mn1—Mn2—C4	179.62 (16)	O7—Mn1—O9—Mn2	-40.61 (8)
C3—Mn1—Mn2—C5	123.05 (16)	C4—Mn2—O9—C19	42.6 (3)
C2—Mn1—Mn2—C5	1.50 (14)	C5—Mn2—O9—C19	-48.6 (3)
C1—Mn1—Mn2—C5	-118.33 (15)	O8—Mn2—O9—C19	138.9 (3)
O8—Mn1—Mn2—C5	-179.49 (12)	O7—Mn2—O9—C19	-142.4 (3)
O9—Mn1—Mn2—C5	-58.33 (13)	Mn1—Mn2—O9—C19	177.2 (3)
O7—Mn1—Mn2—C5	59.62 (13)	C4—Mn2—O9—Mn1	-134.64 (11)
C3—Mn1—Mn2—O8	-57.46 (15)	C5—Mn2—O9—Mn1	134.13 (9)
C2—Mn1—Mn2—O8	-179.00 (12)	O8—Mn2—O9—Mn1	-38.27 (7)
C1—Mn1—Mn2—O8	61.16 (14)	O7—Mn2—O9—Mn1	40.34 (7)
O9—Mn1—Mn2—O8	121.16 (11)	Mn1—O7—C7—C12	-169.8 (2)
O7—Mn1—Mn2—O8	-120.89 (11)	Mn2—O7—C7—C12	-19.6 (4)
C3—Mn1—Mn2—O9	-178.62 (16)	Mn1—O7—C7—C8	11.9 (4)
C2—Mn1—Mn2—O9	59.84 (12)	Mn2—O7—C7—C8	162.1 (2)
C1—Mn1—Mn2—O9	-60.00 (14)	O7—C7—C8—C9	-179.5 (3)
O8—Mn1—Mn2—O9	-121.16 (11)	C12—C7—C8—C9	2.2 (4)
O7—Mn1—Mn2—O9	117.95 (12)	C7—C8—C9—C10	0.2 (5)
C3—Mn1—Mn2—O7	63.43 (15)	C8—C9—C10—C11	-2.1 (5)
C2—Mn1—Mn2—O7	-58.11 (13)	C9—C10—C11—C12	1.7 (5)
C1—Mn1—Mn2—O7	-177.95 (15)	C10—C11—C12—C7	0.7 (5)
O8—Mn1—Mn2—O7	120.89 (11)	O7—C7—C12—C11	179.0 (3)
O9—Mn1—Mn2—O7	-117.95 (12)	C8—C7—C12—C11	-2.6 (4)
C3—Mn1—O7—C7	23.9 (3)	Mn1—O8—C13—C14	7.0 (4)
C2—Mn1—O7—C7	-65.0 (2)	Mn2—O8—C13—C14	-177.19 (19)
C1—Mn1—O7—C7	166.7 (5)	Mn1—O8—C13—C18	-173.50 (19)
O8—Mn1—O7—C7	120.3 (2)	Mn2—O8—C13—C18	2.3 (4)
O9—Mn1—O7—C7	-161.1 (2)	O8—C13—C14—C15	179.6 (3)
Mn2—Mn1—O7—C7	158.7 (3)	C18—C13—C14—C15	0.1 (4)
C3—Mn1—O7—Mn2	-134.81 (11)	C13—C14—C15—C16	-0.3 (5)
C2—Mn1—O7—Mn2	136.31 (10)	C14—C15—C16—C17	0.0 (5)
C1—Mn1—O7—Mn2	8.0 (6)	C15—C16—C17—C18	0.4 (5)
O8—Mn1—O7—Mn2	-38.36 (8)	C16—C17—C18—C13	-0.6 (5)
O9—Mn1—O7—Mn2	40.22 (8)	O8—C13—C18—C17	-179.2 (3)
C6—Mn2—O7—C7	-25.2 (3)	C14—C13—C18—C17	0.3 (4)
C5—Mn2—O7—C7	68.0 (2)	Mn1—O9—C19—C24	170.61 (19)
O8—Mn2—O7—C7	-120.3 (2)	Mn2—O9—C19—C24	-5.6 (4)
O9—Mn2—O7—C7	161.2 (2)	Mn1—O9—C19—C20	-10.7 (4)
Mn1—Mn2—O7—C7	-158.7 (3)	Mn2—O9—C19—C20	173.16 (19)
C6—Mn2—O7—Mn1	133.50 (10)	O9—C19—C20—C21	-178.5 (2)
C5—Mn2—O7—Mn1	-133.28 (10)	C24—C19—C20—C21	0.3 (4)
O8—Mn2—O7—Mn1	38.38 (8)	C19—C20—C21—C22	-0.4 (4)
O9—Mn2—O7—Mn1	-40.12 (8)	C20—C21—C22—C23	-0.1 (4)
C3—Mn1—O8—C13	-44.6 (2)	C21—C22—C23—C24	0.6 (4)
C1—Mn1—O8—C13	45.1 (2)	C22—C23—C24—C19	-0.7 (4)
O9—Mn1—O8—C13	138.5 (2)	O9—C19—C24—C23	179.0 (2)
O7—Mn1—O8—C13	-144.0 (2)	C20—C19—C24—C23	0.2 (4)
Mn2—Mn1—O8—C13	177.0 (3)	C31—N1—C25—C26	-61.1 (4)
C3—Mn1—O8—Mn2	138.41 (10)	C29—N1—C25—C26	177.9 (3)

supplementary materials

C1—Mn1—O8—Mn2	-131.90 (10)	C27—N1—C25—C26	56.5 (4)
O9—Mn1—O8—Mn2	-38.55 (7)	C31—N1—C27—C28	-179.5 (2)
O7—Mn1—O8—Mn2	39.00 (7)	C29—N1—C27—C28	-58.0 (3)
C6—Mn2—O8—C13	47.6 (3)	C25—N1—C27—C28	60.2 (3)
C4—Mn2—O8—C13	-40.9 (3)	C31—N1—C29—C30	60.3 (3)
C5—Mn2—O8—C13	-175.0 (4)	C27—N1—C29—C30	-58.1 (3)
O9—Mn2—O8—C13	-138.5 (2)	C25—N1—C29—C30	-179.0 (3)
O7—Mn2—O8—C13	144.4 (3)	C29—N1—C31—C32	56.2 (3)
Mn1—Mn2—O8—C13	-176.9 (3)	C27—N1—C31—C32	177.8 (3)
C6—Mn2—O8—Mn1	-135.45 (9)	C25—N1—C31—C32	-61.7 (4)
C4—Mn2—O8—Mn1	136.02 (11)		

Fig. 1




Fig. 2

