

## *N*-(2,6-Dichlorophenyl)succinamic acid

 B. S. Saraswathi,<sup>a</sup> Sabine Foro<sup>b</sup> and B. Thimme Gowda<sup>a\*</sup>

<sup>a</sup>Department of Chemistry, Mangalore University, Mangalagangothri 574 199, Mangalore, India, and <sup>b</sup>Institute of Materials Science, Darmstadt University of Technology, Petersenstrasse 23, D-64287 Darmstadt, Germany  
Correspondence e-mail: gowdabt@yahoo.com


Received 22 June 2011; accepted 25 June 2011

Key indicators: single-crystal X-ray study;  $T = 293$  K; mean  $\sigma(\text{C}-\text{C}) = 0.014$  Å;  $R$  factor = 0.119;  $wR$  factor = 0.222; data-to-parameter ratio = 13.6.

In the crystal of the title compound,  $\text{C}_{10}\text{H}_9\text{Cl}_2\text{NO}_3$ , the conformations of the amide O atom and the carbonyl O atom of the acid segment are *anti* to each other and to the H atoms on the adjacent  $-\text{CH}_2$  groups. The  $\text{C}=\text{O}$  and  $\text{O}-\text{H}$  bonds of the acid group are *syn* to one another. In the crystal, molecules are packed into infinite chains through intermolecular  $\text{O}-\text{H}\cdots\text{O}$  and  $\text{N}-\text{H}\cdots\text{O}$  hydrogen bonds.

### Related literature

For our studies of the effect of substituents on the structures and other aspects of *N*-(aryl)-amides, see: Bhat & Gowda (2000); Gowda *et al.* (2000, 2009*a,b*). For modes of interlinking carboxylic acids by hydrogen bonds, see: Leiserowitz (1976). For packing of molecules involving dimeric hydrogen-bonding associations of each carboxyl group with a centrosymmetrically related neighbor, see: Jagannathan *et al.* (1994).



### Experimental

#### Crystal data

$\text{C}_{10}\text{H}_9\text{Cl}_2\text{NO}_3$   
 $M_r = 262.08$   
Monoclinic,  $P2_1/n$

$a = 4.713$  (1) Å  
 $b = 11.963$  (3) Å  
 $c = 20.687$  (4) Å

$\beta = 94.64$  (2)°  
 $V = 1162.5$  (4) Å<sup>3</sup>  
 $Z = 4$   
Mo  $K\alpha$  radiation

$\mu = 0.55$  mm<sup>-1</sup>  
 $T = 293$  K  
 $0.48 \times 0.06 \times 0.04$  mm

#### Data collection

Oxford Diffraction Xcalibur diffractometer with a Sapphire CCD detector  
Absorption correction: multi-scan (*CrysAlis RED*; Oxford)

Diffraction, 2009)  
 $T_{\min} = 0.779$ ,  $T_{\max} = 0.978$   
3912 measured reflections  
1965 independent reflections  
1189 reflections with  $I > 2\sigma(I)$   
 $R_{\text{int}} = 0.052$

#### Refinement

$R[F^2 > 2\sigma(F^2)] = 0.119$   
 $wR(F^2) = 0.222$   
 $S = 1.34$   
1965 reflections  
145 parameters

18 restraints  
H-atom parameters constrained  
 $\Delta\rho_{\max} = 0.57$  e Å<sup>-3</sup>  
 $\Delta\rho_{\min} = -0.55$  e Å<sup>-3</sup>

**Table 1**

Hydrogen-bond geometry (Å, °).

$D-\text{H}\cdots A$	$D-\text{H}$	$\text{H}\cdots A$	$D\cdots A$	$D-\text{H}\cdots A$
$\text{N1}-\text{H1N}\cdots\text{O1}^{\text{i}}$	0.86	2.06	2.875 (9)	159
$\text{O2}-\text{H2O}\cdots\text{O3}^{\text{ii}}$	0.82	1.89	2.678 (11)	162

Symmetry codes: (i)  $x - 1, y, z$ ; (ii)  $-x + 1, -y - 1, -z + 1$ .

Data collection: *CrysAlis CCD* (Oxford Diffraction, 2009); cell refinement: *CrysAlis RED* (Oxford Diffraction, 2009); data reduction: *CrysAlis RED*; program(s) used to solve structure: *SHELXS97* (Sheldrick, 2008); program(s) used to refine structure: *SHELXL97* (Sheldrick, 2008); molecular graphics: *PLATON* (Spek, 2009); software used to prepare material for publication: *SHELXL97*.

BSS thanks the University Grants Commission, Government of India, New Delhi, for the award of a research fellowship under its faculty improvement program.

Supplementary data and figures for this paper are available from the IUCr electronic archives (Reference: SJ5173).

### References

- Bhat, D. K. & Gowda, B. T. (2000). *J. Indian Chem. Soc.* **77**, 279–284.  
Gowda, B. T., Foro, S., Saraswathi, B. S., Terao, H. & Fuess, H. (2009*a*). *Acta Cryst.* **E65**, o399.  
Gowda, B. T., Foro, S., Saraswathi, B. S., Terao, H. & Fuess, H. (2009*b*). *Acta Cryst.* **E65**, o466.  
Gowda, B. T., Kumar, B. H. A. & Fuess, H. (2000). *Z. Naturforsch. Teil A*, **55**, 721–728.  
Jagannathan, N. R., Rajan, S. S. & Subramanian, E. (1994). *J. Chem. Crystallogr.* **24**, 75–78.  
Leiserowitz, L. (1976). *Acta Cryst.* **B32**, 775–802.  
Oxford Diffraction (2009). *CrysAlis CCD* and *CrysAlis RED*. Oxford Diffraction Ltd, Yarnton, England.  
Sheldrick, G. M. (2008). *Acta Cryst.* **A64**, 112–122.  
Spek, A. L. (2009). *Acta Cryst.* **D65**, 148–155.

**supplementary materials**

*Acta Cryst.* (2011). E67, o1880 [ doi:10.1107/S1600536811024949 ]

## ***N*-(2,6-Dichlorophenyl)succinamic acid**

**B. S. Saraswathi, S. Foro and B. T. Gowda**

### **Comment**

The amide moiety is an important constituent of many biologically important compounds. As part of our studies into the substituent effects on the structures and other aspects of this class of compounds (Bhat & Gowda, 2000; Gowda *et al.*, 2000, 2009*a,b*), in the present work, the crystal structure of *N*-(2,6-dichlorophenyl)-succinamic acid (I) has been determined (Fig. 1). The conformation of the amide O atom and the carbonyl O atom of the acid segment are *anti* to each other and are also *anti* to the H atoms attached to the adjacent C atoms (Fig. 1). Further, C=O and O—H bonds of the acid group are *syn* to each other, similar to that observed in *N*-(2-chlorophenyl)-succinamic acid (Gowda *et al.*, 2009*a*) and *N*-(2,6-dimethylphenyl)-succinamic acid (Gowda *et al.*, 2009*b*).

In the structure, the intermolecular O—H...O and N—H...O hydrogen bonds pack the molecules into infinite chains (Table 1, Fig.2).

The modes of interlinking carboxylic acids by hydrogen bonds is described elsewhere (Leiserowitz, 1976). The packing of molecules involving dimeric hydrogen bonded association of each carboxyl group with a centrosymmetrically related neighbor has also been observed (Jagannathan *et al.*, 1994).

### **Experimental**

A solution of succinic anhydride (0.01 mole) in toluene (25 ml) was treated dropwise with the solution of 2,6-dichloroaniline (0.01 mole) also in toluene (20 ml) with constant stirring. The resulting mixture was stirred for about one hour and set aside for an additional hour at room temperature for completion of the reaction. The mixture was then treated with dilute hydrochloric acid to remove the unreacted 2,6-dichloroaniline. The resultant title compound was filtered under suction and washed thoroughly with water to remove the unreacted succinic anhydride and succinic acid. It was recrystallized to constant melting point from ethanol. The purity of the compound was checked and characterized by its infrared and NMR spectra.

Colorless needle like single crystals used in X-ray diffraction studies were grown in ethanolic solution by slow evaporation at room temperature.

### **Refinement**

The H atoms were positioned with idealized geometry using a riding model with the aromatic C—H = 0.93 Å, methylene C—H = 0.97 Å, N—H = 0.86 Å and O—H = 0.82 Å, and were refined with isotropic displacement parameters (set to 1.2 times of the  $U_{eq}$  of the parent atom).

The crystals available for X-ray analysis were of rather poor quality and weak scatterers at high theta value with very low intensity, resulting in relatively high *R* values. The crystal has 37.1% weak reflections.

## Figures




Fig. 1. Molecular structure of the title compound, showing the atom labelling scheme and displacement ellipsoids are drawn at the 50% probability level.




Fig. 2. Molecular packing of the title compound with hydrogen bonding shown as dashed lines.

## *N*-(2,6-Dichlorophenyl)succinamic acid

### Crystal data

$C_{10}H_9Cl_2NO_3$

$M_r = 262.08$

Monoclinic,  $P2_1/n$

Hall symbol: -P 2yn

$a = 4.713$  (1) Å

$b = 11.963$  (3) Å

$c = 20.687$  (4) Å

$\beta = 94.64$  (2)°

$V = 1162.5$  (4) Å<sup>3</sup>

$Z = 4$

$F(000) = 536$

$D_x = 1.497$  Mg m<sup>-3</sup>

Mo  $K\alpha$  radiation,  $\lambda = 0.71073$  Å

Cell parameters from 777 reflections

$\theta = 3.0$ – $27.7$ °

$\mu = 0.55$  mm<sup>-1</sup>

$T = 293$  K

Needle, colourless

$0.48 \times 0.06 \times 0.04$  mm

### Data collection

Oxford Diffraction Xcalibur diffractometer with a Sapphire CCD detector

Radiation source: fine-focus sealed tube

graphite

Rotation method data acquisition using  $\omega$  scans

Absorption correction: multi-scan (*CrysAlis RED*; Oxford Diffraction, 2009)

$T_{\min} = 0.779$ ,  $T_{\max} = 0.978$

3912 measured reflections

1965 independent reflections

1189 reflections with  $I > 2\sigma(I)$

$R_{\text{int}} = 0.052$

$\theta_{\max} = 25.2$ °,  $\theta_{\min} = 3.4$ °

$h = -5 \rightarrow 5$

$k = -12 \rightarrow 14$

$l = -16 \rightarrow 24$

### Refinement

Refinement on  $F^2$

Least-squares matrix: full

$R[F^2 > 2\sigma(F^2)] = 0.119$

$wR(F^2) = 0.222$

Primary atom site location: structure-invariant direct methods

Secondary atom site location: difference Fourier map

Hydrogen site location: inferred from neighbouring sites

H-atom parameters constrained

$S = 1.34$	$w = 1/[\sigma^2(F_o^2) + (0.P)^2 + 8.7677P]$
1965 reflections	where $P = (F_o^2 + 2F_c^2)/3$
145 parameters	$(\Delta/\sigma)_{\max} < 0.001$
18 restraints	$\Delta\rho_{\max} = 0.57 \text{ e } \text{\AA}^{-3}$
	$\Delta\rho_{\min} = -0.55 \text{ e } \text{\AA}^{-3}$

*Special details*

**Geometry.** All e.s.d.'s (except the e.s.d. in the dihedral angle between two l.s. planes) are estimated using the full covariance matrix. The cell e.s.d.'s are taken into account individually in the estimation of e.s.d.'s in distances, angles and torsion angles; correlations between e.s.d.'s in cell parameters are only used when they are defined by crystal symmetry. An approximate (isotropic) treatment of cell e.s.d.'s is used for estimating e.s.d.'s involving l.s. planes.

**Refinement.** Refinement of  $F^2$  against ALL reflections. The weighted  $R$ -factor  $wR$  and goodness of fit  $S$  are based on  $F^2$ , conventional  $R$ -factors  $R$  are based on  $F$ , with  $F$  set to zero for negative  $F^2$ . The threshold expression of  $F^2 > \sigma(F^2)$  is used only for calculating  $R$ -factors(gt) *etc.* and is not relevant to the choice of reflections for refinement.  $R$ -factors based on  $F^2$  are statistically about twice as large as those based on  $F$ , and  $R$ -factors based on ALL data will be even larger.

*Fractional atomic coordinates and isotropic or equivalent isotropic displacement parameters ( $\text{\AA}^2$ )*

	$x$	$y$	$z$	$U_{\text{iso}}^*/U_{\text{eq}}$
C1	0.5925 (18)	0.1082 (7)	0.3449 (4)	0.028 (2)
C2	0.4791 (19)	0.1566 (8)	0.2870 (5)	0.039 (2)
C3	0.554 (2)	0.2620 (8)	0.2688 (5)	0.049 (3)
H3	0.4712	0.2925	0.2304	0.059*
C4	0.749 (3)	0.3227 (9)	0.3068 (6)	0.063 (4)
H4	0.8010	0.3938	0.2938	0.076*
C5	0.871 (2)	0.2786 (9)	0.3643 (6)	0.058 (3)
H5	1.0064	0.3186	0.3899	0.069*
C6	0.785 (2)	0.1730 (8)	0.3832 (5)	0.041 (3)
C7	0.6868 (18)	-0.0859 (7)	0.3757 (4)	0.027 (2)
C8	0.5518 (18)	-0.1883 (8)	0.4030 (5)	0.039 (2)
H8A	0.3759	-0.2043	0.3771	0.047*
H8B	0.5035	-0.1716	0.4467	0.047*
C9	0.7363 (19)	-0.2911 (7)	0.4051 (4)	0.034 (2)
H9A	0.7487	-0.3182	0.3613	0.041*
H9B	0.9269	-0.2711	0.4225	0.041*
C10	0.627 (2)	-0.3827 (8)	0.4453 (5)	0.044 (3)
N1	0.5076 (14)	0.0022 (6)	0.3650 (3)	0.0312 (18)
H1N	0.3308	-0.0076	0.3709	0.037*
O1	0.9365 (12)	-0.0820 (5)	0.3658 (3)	0.0348 (16)
O2	0.738 (2)	-0.4776 (7)	0.4371 (4)	0.095 (3)
H2O	0.6475	-0.5255	0.4551	0.114*
O3	0.460 (2)	-0.3676 (7)	0.4844 (4)	0.089 (3)
Cl1	0.2395 (6)	0.0781 (3)	0.23715 (14)	0.0625 (9)
Cl2	0.9252 (7)	0.1234 (3)	0.45814 (13)	0.0621 (9)

## supplementary materials

---

### Atomic displacement parameters ( $\text{\AA}^2$ )

	$U^{11}$	$U^{22}$	$U^{33}$	$U^{12}$	$U^{13}$	$U^{23}$
C1	0.024 (5)	0.017 (5)	0.046 (6)	0.003 (4)	0.012 (4)	-0.001 (4)
C2	0.033 (6)	0.038 (6)	0.047 (6)	0.006 (5)	0.007 (5)	0.009 (5)
C3	0.061 (7)	0.032 (6)	0.059 (7)	0.012 (6)	0.026 (6)	0.020 (6)
C4	0.074 (9)	0.024 (6)	0.096 (10)	-0.002 (6)	0.030 (8)	0.006 (7)
C5	0.062 (8)	0.028 (6)	0.083 (9)	-0.003 (6)	0.009 (7)	-0.003 (6)
C6	0.045 (6)	0.029 (6)	0.051 (7)	0.012 (5)	0.017 (5)	0.002 (5)
C7	0.023 (5)	0.023 (5)	0.035 (5)	-0.003 (4)	0.000 (4)	0.004 (4)
C8	0.025 (5)	0.031 (6)	0.059 (7)	0.001 (4)	0.000 (5)	0.016 (5)
C9	0.032 (5)	0.030 (5)	0.042 (6)	-0.003 (4)	0.013 (4)	0.002 (4)
C10	0.048 (6)	0.022 (5)	0.065 (7)	0.021 (5)	0.025 (5)	0.015 (5)
N1	0.019 (4)	0.027 (4)	0.049 (5)	-0.003 (3)	0.010 (3)	0.014 (3)
O1	0.018 (3)	0.028 (4)	0.059 (4)	-0.001 (3)	0.011 (3)	0.005 (3)
O2	0.119 (7)	0.057 (5)	0.120 (6)	0.012 (5)	0.075 (5)	0.031 (5)
O3	0.112 (6)	0.048 (5)	0.116 (6)	0.021 (5)	0.068 (5)	0.028 (5)
Cl1	0.0596 (18)	0.063 (2)	0.0619 (19)	0.0017 (16)	-0.0145 (14)	0.0092 (15)
Cl2	0.080 (2)	0.0543 (18)	0.0492 (17)	0.0067 (16)	-0.0129 (15)	-0.0084 (15)

### Geometric parameters ( $\text{\AA}$ , $^\circ$ )

C1—C6	1.392 (13)	C7—N1	1.358 (10)
C1—C2	1.396 (12)	C7—C8	1.511 (12)
C1—N1	1.402 (10)	C8—C9	1.505 (12)
C2—C3	1.370 (13)	C8—H8A	0.9700
C2—Cl1	1.741 (10)	C8—H8B	0.9700
C3—C4	1.370 (15)	C9—C10	1.491 (12)
C3—H3	0.9300	C9—H9A	0.9700
C4—C5	1.382 (15)	C9—H9B	0.9700
C4—H4	0.9300	C10—O3	1.190 (11)
C5—C6	1.391 (14)	C10—O2	1.266 (11)
C5—H5	0.9300	N1—H1N	0.8600
C6—Cl2	1.739 (10)	O2—H2O	0.8200
C7—O1	1.212 (9)		
C6—C1—C2	116.4 (8)	N1—C7—C8	114.5 (7)
C6—C1—N1	121.6 (8)	C7—C8—C9	114.4 (7)
C2—C1—N1	122.0 (8)	C7—C8—H8A	108.7
C3—C2—C1	121.9 (10)	C9—C8—H8A	108.7
C3—C2—Cl1	120.1 (8)	C7—C8—H8B	108.7
C1—C2—Cl1	118.0 (7)	C9—C8—H8B	108.7
C2—C3—C4	120.4 (11)	H8A—C8—H8B	107.6
C2—C3—H3	119.8	C10—C9—C8	113.1 (7)
C4—C3—H3	119.8	C10—C9—H9A	109.0
C3—C4—C5	120.2 (11)	C8—C9—H9A	109.0
C3—C4—H4	119.9	C10—C9—H9B	109.0
C5—C4—H4	119.9	C8—C9—H9B	109.0

C4—C5—C6	118.7 (11)	H9A—C9—H9B	107.8
C4—C5—H5	120.7	O3—C10—O2	121.9 (9)
C6—C5—H5	120.7	O3—C10—C9	123.1 (9)
C5—C6—C1	122.4 (10)	O2—C10—C9	114.9 (8)
C5—C6—Cl2	117.6 (9)	C7—N1—C1	124.2 (7)
C1—C6—Cl2	120.0 (7)	C7—N1—H1N	117.9
O1—C7—N1	122.9 (8)	C1—N1—H1N	117.9
O1—C7—C8	122.6 (8)	C10—O2—H2O	109.5
C6—C1—C2—C3	0.2 (13)	C2—C1—C6—Cl2	176.8 (7)
N1—C1—C2—C3	177.3 (8)	N1—C1—C6—Cl2	-0.4 (12)
C6—C1—C2—Cl1	179.4 (7)	O1—C7—C8—C9	-11.0 (13)
N1—C1—C2—Cl1	-3.4 (11)	N1—C7—C8—C9	170.9 (8)
C1—C2—C3—C4	1.5 (15)	C7—C8—C9—C10	167.1 (9)
Cl1—C2—C3—C4	-177.8 (8)	C8—C9—C10—O3	-18.2 (16)
C2—C3—C4—C5	-0.9 (17)	C8—C9—C10—O2	165.2 (10)
C3—C4—C5—C6	-1.3 (17)	O1—C7—N1—C1	-3.5 (14)
C4—C5—C6—C1	3.1 (15)	C8—C7—N1—C1	174.5 (8)
C4—C5—C6—Cl2	-176.2 (8)	C6—C1—N1—C7	-62.7 (12)
C2—C1—C6—C5	-2.5 (13)	C2—C1—N1—C7	120.3 (10)
N1—C1—C6—C5	-179.6 (9)		

*Hydrogen-bond geometry (Å, °)*

<i>D</i> —H··· <i>A</i>	<i>D</i> —H	H··· <i>A</i>	<i>D</i> ··· <i>A</i>	<i>D</i> —H··· <i>A</i>
N1—H1N···O1 <sup>i</sup>	0.86	2.06	2.875 (9)	159
O2—H2O···O3 <sup>ii</sup>	0.82	1.89	2.678 (11)	162

Symmetry codes: (i)  $x-1, y, z$ ; (ii)  $-x+1, -y-1, -z+1$ .

Fig. 1






Fig. 2

