

N-(3-Chlorophenyl)succinimide**B. S. Saraswathi,^a Sabine Foro^b and B. Thimme Gowda^{a*}**

^aDepartment of Chemistry, Mangalore University, Mangalagangotri 574 199, Mangalore, India, and ^bInstitute of Materials Science, Darmstadt University of Technology, Petersenstrasse 23, D-64287 Darmstadt, Germany
Correspondence e-mail: gowdabt@yahoo.com


Received 4 July 2011; accepted 5 July 2011

Key indicators: single-crystal X-ray study; $T = 293\text{ K}$; mean $\sigma(\text{C}-\text{C}) = 0.006\text{ \AA}$; R factor = 0.082; wR factor = 0.137; data-to-parameter ratio = 13.8.

In the title compound, $\text{C}_{10}\text{H}_8\text{ClNO}_2$, the chlorobenzene and the essentially planar (r.m.s. deviation = 0.030 Å) pyrrolidine ring are tilted by 59.5 (1)° with respect to one another.

Related literature

For our studies on the effects of substituents on the structures of *N*-(aryl)-amides, see: Bhat & Gowda (2000); Gowda *et al.* (1999, 2007); Saraswathi *et al.* (2010*a,b*).

**Experimental***Crystal data*

$\text{C}_{10}\text{H}_8\text{ClNO}_2$
 $M_r = 209.62$
Orthorhombic, $Pbca$

$a = 12.884(2)\text{ \AA}$
 $b = 7.173(1)\text{ \AA}$
 $c = 20.805(3)\text{ \AA}$

$V = 1922.7(5)\text{ \AA}^3$
 $Z = 8$
Mo $K\alpha$ radiation

$\mu = 0.37\text{ mm}^{-1}$
 $T = 293\text{ K}$
 $0.46 \times 0.12 \times 0.09\text{ mm}$

Data collection

Oxford Diffraction Xcalibur diffractometer with a Sapphire CCD detector
Absorption correction: multi-scan (*CrysAlis RED*; Oxford)

Diffraction, 2009)
 $T_{\min} = 0.849$, $T_{\max} = 0.968$
6087 measured reflections
1755 independent reflections
1163 reflections with $I > 2\sigma(I)$
 $R_{\text{int}} = 0.044$

Refinement

$R[F^2 > 2\sigma(F^2)] = 0.082$
 $wR(F^2) = 0.137$
 $S = 1.33$
1755 reflections

127 parameters
H-atom parameters constrained
 $\Delta\rho_{\text{max}} = 0.32\text{ e \AA}^{-3}$
 $\Delta\rho_{\text{min}} = -0.46\text{ e \AA}^{-3}$

Data collection: *CrysAlis CCD* (Oxford Diffraction, 2009); cell refinement: *CrysAlis RED* (Oxford Diffraction, 2009); data reduction: *CrysAlis RED*; program(s) used to solve structure: *SHELXS97* (Sheldrick, 2008); program(s) used to refine structure: *SHELXL97* (Sheldrick, 2008); molecular graphics: *PLATON* (Spek, 2009); software used to prepare material for publication: *SHELXL97*.

BSS thanks the University Grants Commission, Government of India, New Delhi, for the award of a research fellowship under its faculty improvement program.

Supplementary data and figures for this paper are available from the IUCr electronic archives (Reference: BT5570).

References

- Bhat, D. K. & Gowda, B. T. (2000). *J. Indian Chem. Soc.* **77**, 279–284.
- Gowda, B. T., Bhat, D. K., Fuess, H. & Weiss, A. (1999). *Z. Naturforsch. Teil A*, **54**, 261–267.
- Gowda, B. T., Foro, S. & Fuess, H. (2007). *Acta Cryst. E* **63**, o1975–o1976.
- Oxford Diffraction (2009). *CrysAlis CCD* and *CrysAlis RED*. Oxford Diffraction Ltd, Yarnton, England.
- Saraswathi, B. S., Gowda, B. T., Foro, S. & Fuess, H. (2010*a*). *Acta Cryst. E* **66**, o921.
- Saraswathi, B. S., Gowda, B. T., Foro, S. & Fuess, H. (2010*b*). *Acta Cryst. E* **66**, o1269.
- Sheldrick, G. M. (2008). *Acta Cryst. A* **64**, 112–122.
- Spek, A. L. (2009). *Acta Cryst. D* **65**, 148–155.

supplementary materials

Acta Cryst. (2011). E67, o1977 [doi:10.1107/S1600536811026845]

N-(3-Chlorophenyl)succinimide

B. S. Saraswathi, S. Foro and B. T. Gowda

Comment

As a part of our studies on the effects of ring and side chain substitutions on the structures and other aspects of biologically significant compounds (Bhat & Gowda, 2000; Gowda *et al.*, 1999, 2007; Saraswathi *et al.*, 2010*a,b*), the crystal structure of *N*-(3-chlorophenyl)succinimide has been determined (Fig. 1). In the structure, the molecule is non-planar with the benzene and pyrrolidine rings tilted by 59.5 (1) $^{\circ}$ with respect to one another, compared to the values of 69.5 (1) $^{\circ}$ in *N*-(2-chlorophenyl)- succinimide (Saraswathi *et al.*, 2010*a*) and 52.5 (1) $^{\circ}$ in *N*-(3-methylphenyl)succinimide (Saraswathi *et al.*, 2010*b*).

The torsional angles of the groups, C2 - C1 - N1 - C7, C6 - C1 - N1 - C7, C2 - C1 - N1 - C10 and C6 - C1 - N1 - C10 in the molecule are -117.5 (5), 61.9 (5), 57.7 (5) $^{\circ}$ and -123.0 (4), respectively, while the torsional angles of the groups, O1 - C7 - N1 - C1, C8 - C7 - N1 - C1, O2 - C10 - N1 - C1 and C9 - C10 - N1 - C1 are 0.5 (6), -178.4 (4), 2.7 (7) and -177.6 (4) $^{\circ}$, respectively.

The packing of molecules into layered chains along *a*-axis is shown in Fig. 2.

Experimental

The solution of succinic anhydride (0.02 mole) in toluene (25 ml) was treated dropwise with the solution of 3-chloroaniline (0.02 mole) also in toluene (20 ml) with constant stirring. The resulting mixture was stirred for one hour and set aside for an additional hour at room temperature for the completion of reaction. The mixture was then treated with dilute hydrochloric acid to remove the unreacted 3-chloroaniline. The resultant solid *N*-(3-chlorophenyl)succinamic acid was filtered under suction and washed thoroughly with water to remove the unreacted succinic anhydride and succinic acid. It was recrystallized to constant melting point from ethanol.

N-(3-chlorophenyl)succinamic acid was heated for 2 h and then allowed to cool slowly to room temperature to get the compound, *N*-(3-chlorophenyl)succinimide. The purity of the compound was checked and characterized by its infrared spectra.

Needle like colourless single crystals of the compound used in X-ray diffraction studies were grown in ethanolic solution by a slow evaporation at room temperature.

Refinement

The H atoms were positioned with idealized geometry using a riding model with the aromatic C—H = 0.93 Å and methylene C—H = 0.97 Å and were refined with isotropic displacement parameters set to 1.2 times of the U_{eq} of the parent atom.

supplementary materials

Figures




Fig. 1. Molecular structure of (I), showing the atom labeling. Displacement ellipsoids drawn at the 50% probability level.




Fig. 2. Molecular packing of (I).

N-(3-Chlorophenyl)succinimide

Crystal data

| | |
|--|---|
| C ₁₀ H ₈ ClNO ₂ | <i>F</i> (000) = 864 |
| <i>M_r</i> = 209.62 | <i>D_x</i> = 1.448 Mg m ⁻³ |
| Orthorhombic, <i>Pbca</i> | Mo <i>Kα</i> radiation, λ = 0.71073 Å |
| Hall symbol: -P 2ac 2ab | Cell parameters from 1436 reflections |
| <i>a</i> = 12.884 (2) Å | θ = 2.8–27.8° |
| <i>b</i> = 7.173 (1) Å | μ = 0.37 mm ⁻¹ |
| <i>c</i> = 20.805 (3) Å | <i>T</i> = 293 K |
| <i>V</i> = 1922.7 (5) Å ³ | Needle, colourless |
| <i>Z</i> = 8 | 0.46 × 0.12 × 0.09 mm |

Data collection

| | |
|--|--|
| Oxford Diffraction Xcalibur | 1755 independent reflections |
| diffractometer with a Sapphire CCD detector | 1163 reflections with $I > 2\sigma(I)$ |
| Radiation source: fine-focus sealed tube | R_{int} = 0.044 |
| graphite | $\theta_{\text{max}} = 25.4^\circ$, $\theta_{\text{min}} = 3.2^\circ$ |
| Rotation method data acquisition using ω scans | $h = -9 \rightarrow 15$ |
| Absorption correction: multi-scan (<i>CrysAlis RED</i> ; Oxford Diffraction, 2009) | $k = -6 \rightarrow 8$ |
| $T_{\text{min}} = 0.849$, $T_{\text{max}} = 0.968$ | $l = -25 \rightarrow 22$ |
| 6087 measured reflections | |

Refinement

| | |
|---------------------------------|--|
| Refinement on F^2 | Primary atom site location: structure-invariant direct methods |
| Least-squares matrix: full | Secondary atom site location: difference Fourier map |
| $R[F^2 > 2\sigma(F^2)] = 0.082$ | Hydrogen site location: inferred from neighbouring sites |
| $wR(F^2) = 0.137$ | H-atom parameters constrained |

| | |
|------------------|--|
| $S = 1.33$ | $w = 1/[\sigma^2(F_o^2) + (0.0157P)^2 + 3.1516P]$ |
| | where $P = (F_o^2 + 2F_c^2)/3$ |
| 1755 reflections | $(\Delta/\sigma)_{\max} = 0.001$ |
| 127 parameters | $\Delta\rho_{\max} = 0.32 \text{ e } \text{\AA}^{-3}$ |
| 0 restraints | $\Delta\rho_{\min} = -0.46 \text{ e } \text{\AA}^{-3}$ |

Special details

Geometry. All e.s.d.'s (except the e.s.d. in the dihedral angle between two l.s. planes) are estimated using the full covariance matrix. The cell e.s.d.'s are taken into account individually in the estimation of e.s.d.'s in distances, angles and torsion angles; correlations between e.s.d.'s in cell parameters are only used when they are defined by crystal symmetry. An approximate (isotropic) treatment of cell e.s.d.'s is used for estimating e.s.d.'s involving l.s. planes.

Refinement. Refinement of F^2 against ALL reflections. The weighted R -factor wR and goodness of fit S are based on F^2 , conventional R -factors R are based on F , with F set to zero for negative F^2 . The threshold expression of $F^2 > \sigma(F^2)$ is used only for calculating R -factors(gt) etc. and is not relevant to the choice of reflections for refinement. R -factors based on F^2 are statistically about twice as large as those based on F , and R -factors based on ALL data will be even larger.

Fractional atomic coordinates and isotropic or equivalent isotropic displacement parameters (\AA^2)

| | x | y | z | $U_{\text{iso}}^*/U_{\text{eq}}$ |
|-----|--------------|------------|--------------|----------------------------------|
| C1 | 0.1240 (3) | 0.3985 (6) | 0.39944 (19) | 0.0406 (10) |
| C2 | 0.1122 (3) | 0.2512 (6) | 0.44197 (19) | 0.0427 (10) |
| H2 | 0.1035 | 0.1300 | 0.4269 | 0.051* |
| C3 | 0.1135 (3) | 0.2881 (7) | 0.5070 (2) | 0.0474 (12) |
| C4 | 0.1279 (3) | 0.4651 (8) | 0.5302 (2) | 0.0572 (13) |
| H4 | 0.1292 | 0.4874 | 0.5743 | 0.069* |
| C5 | 0.1403 (3) | 0.6090 (7) | 0.4874 (2) | 0.0590 (14) |
| H5 | 0.1504 | 0.7294 | 0.5028 | 0.071* |
| C6 | 0.1381 (3) | 0.5785 (6) | 0.4213 (2) | 0.0489 (12) |
| H6 | 0.1459 | 0.6769 | 0.3926 | 0.059* |
| C7 | 0.2069 (4) | 0.3968 (6) | 0.2912 (2) | 0.0437 (11) |
| C8 | 0.1764 (4) | 0.3477 (7) | 0.2235 (2) | 0.0557 (13) |
| H8A | 0.2260 | 0.2619 | 0.2047 | 0.067* |
| H8B | 0.1725 | 0.4586 | 0.1970 | 0.067* |
| C9 | 0.0704 (4) | 0.2568 (7) | 0.2297 (2) | 0.0600 (14) |
| H9A | 0.0203 | 0.3189 | 0.2023 | 0.072* |
| H9B | 0.0738 | 0.1263 | 0.2177 | 0.072* |
| C10 | 0.0405 (4) | 0.2769 (6) | 0.2994 (2) | 0.0498 (12) |
| N1 | 0.1227 (3) | 0.3623 (4) | 0.33150 (16) | 0.0408 (9) |
| O1 | 0.2895 (2) | 0.4592 (4) | 0.30949 (15) | 0.0591 (9) |
| O2 | -0.0397 (3) | 0.2302 (5) | 0.32476 (16) | 0.0706 (10) |
| Cl1 | 0.09625 (11) | 0.1056 (2) | 0.56153 (6) | 0.0722 (5) |

Atomic displacement parameters (\AA^2)

| U^{11} | U^{22} | U^{33} | U^{12} | U^{13} | U^{23} |
|----------|----------|----------|----------|----------|----------|
| | | | | | |

supplementary materials

| | | | | | | |
|-----|------------|-------------|-------------|--------------|--------------|-------------|
| C1 | 0.033 (2) | 0.048 (3) | 0.041 (2) | 0.002 (2) | -0.0020 (19) | -0.006 (2) |
| C2 | 0.042 (3) | 0.042 (2) | 0.043 (3) | 0.004 (2) | -0.001 (2) | -0.006 (2) |
| C3 | 0.036 (3) | 0.067 (3) | 0.039 (3) | 0.002 (2) | -0.001 (2) | -0.005 (2) |
| C4 | 0.042 (3) | 0.085 (4) | 0.044 (3) | -0.004 (3) | 0.003 (2) | -0.021 (3) |
| C5 | 0.041 (3) | 0.065 (3) | 0.071 (3) | -0.008 (3) | 0.006 (3) | -0.035 (3) |
| C6 | 0.038 (3) | 0.045 (3) | 0.064 (3) | -0.004 (2) | 0.005 (2) | -0.008 (2) |
| C7 | 0.055 (3) | 0.035 (2) | 0.042 (2) | 0.003 (2) | 0.000 (2) | 0.005 (2) |
| C8 | 0.076 (3) | 0.052 (3) | 0.040 (3) | 0.002 (3) | -0.003 (2) | 0.007 (2) |
| C9 | 0.081 (4) | 0.053 (3) | 0.046 (3) | -0.006 (3) | -0.018 (3) | 0.003 (2) |
| C10 | 0.057 (3) | 0.039 (3) | 0.053 (3) | -0.003 (2) | -0.013 (3) | 0.006 (2) |
| N1 | 0.044 (2) | 0.040 (2) | 0.0386 (19) | -0.0028 (17) | -0.0024 (17) | 0.0015 (17) |
| O1 | 0.053 (2) | 0.069 (2) | 0.055 (2) | -0.0133 (18) | 0.0055 (17) | 0.0002 (17) |
| O2 | 0.057 (2) | 0.087 (3) | 0.068 (2) | -0.024 (2) | -0.011 (2) | 0.003 (2) |
| Cl1 | 0.0824 (9) | 0.0924 (10) | 0.0418 (6) | 0.0078 (8) | 0.0035 (7) | 0.0082 (7) |

Geometric parameters (\AA , $^\circ$)

| | | | |
|-----------|-----------|------------|-----------|
| C1—C6 | 1.380 (6) | C7—O1 | 1.214 (5) |
| C1—C2 | 1.387 (6) | C7—N1 | 1.393 (5) |
| C1—N1 | 1.437 (5) | C7—C8 | 1.505 (6) |
| C2—C3 | 1.379 (6) | C8—C9 | 1.518 (7) |
| C2—H2 | 0.9300 | C8—H8A | 0.9700 |
| C3—C4 | 1.371 (6) | C8—H8B | 0.9700 |
| C3—Cl1 | 1.746 (5) | C9—C10 | 1.507 (6) |
| C4—C5 | 1.374 (7) | C9—H9A | 0.9700 |
| C4—H4 | 0.9300 | C9—H9B | 0.9700 |
| C5—C6 | 1.393 (6) | C10—O2 | 1.208 (5) |
| C5—H5 | 0.9300 | C10—N1 | 1.395 (5) |
| C6—H6 | 0.9300 | | |
| C6—C1—C2 | 121.2 (4) | N1—C7—C8 | 108.5 (4) |
| C6—C1—N1 | 119.6 (4) | C7—C8—C9 | 104.9 (4) |
| C2—C1—N1 | 119.2 (4) | C7—C8—H8A | 110.8 |
| C3—C2—C1 | 118.6 (4) | C9—C8—H8A | 110.8 |
| C3—C2—H2 | 120.7 | C7—C8—H8B | 110.8 |
| C1—C2—H2 | 120.7 | C9—C8—H8B | 110.8 |
| C4—C3—C2 | 121.7 (4) | H8A—C8—H8B | 108.8 |
| C4—C3—Cl1 | 118.9 (4) | C10—C9—C8 | 105.7 (4) |
| C2—C3—Cl1 | 119.4 (4) | C10—C9—H9A | 110.6 |
| C3—C4—C5 | 118.9 (4) | C8—C9—H9A | 110.6 |
| C3—C4—H4 | 120.6 | C10—C9—H9B | 110.6 |
| C5—C4—H4 | 120.6 | C8—C9—H9B | 110.6 |
| C4—C5—C6 | 121.4 (5) | H9A—C9—H9B | 108.7 |
| C4—C5—H5 | 119.3 | O2—C10—N1 | 124.2 (4) |
| C6—C5—H5 | 119.3 | O2—C10—C9 | 127.8 (4) |
| C1—C6—C5 | 118.3 (4) | N1—C10—C9 | 108.0 (4) |
| C1—C6—H6 | 120.8 | C7—N1—C10 | 112.4 (4) |
| C5—C6—H6 | 120.8 | C7—N1—C1 | 123.3 (3) |
| O1—C7—N1 | 124.1 (4) | C10—N1—C1 | 124.1 (4) |
| O1—C7—C8 | 127.5 (4) | | |

supplementary materials

| | | | |
|--------------|------------|--------------|------------|
| C6—C1—C2—C3 | 0.8 (6) | C8—C9—C10—N1 | -2.7 (5) |
| N1—C1—C2—C3 | -179.9 (4) | O1—C7—N1—C10 | -175.1 (4) |
| C1—C2—C3—C4 | -1.1 (7) | C8—C7—N1—C10 | 6.0 (5) |
| C1—C2—C3—Cl1 | 178.8 (3) | O1—C7—N1—C1 | 0.5 (6) |
| C2—C3—C4—C5 | 0.6 (7) | C8—C7—N1—C1 | -178.4 (4) |
| Cl1—C3—C4—C5 | -179.3 (3) | O2—C10—N1—C7 | 178.3 (4) |
| C3—C4—C5—C6 | 0.3 (7) | C9—C10—N1—C7 | -2.0 (5) |
| C2—C1—C6—C5 | 0.1 (6) | O2—C10—N1—C1 | 2.7 (7) |
| N1—C1—C6—C5 | -179.2 (4) | C9—C10—N1—C1 | -177.6 (4) |
| C4—C5—C6—C1 | -0.6 (7) | C6—C1—N1—C7 | 61.9 (5) |
| O1—C7—C8—C9 | 173.9 (4) | C2—C1—N1—C7 | -117.5 (4) |
| N1—C7—C8—C9 | -7.3 (5) | C6—C1—N1—C10 | -123.0 (4) |
| C7—C8—C9—C10 | 5.9 (5) | C2—C1—N1—C10 | 57.7 (5) |
| C8—C9—C10—O2 | 176.9 (5) | | |

supplementary materials

Fig. 1




Fig. 2

