

1,1-Dibenzyl-3-(4-fluorobenzoyl)thio-urea

Mohd Faizal Md Nasir,^a Ibrahim N. Hassan,^{a*} Wan Ramli Wan Daud,^{b,a} Bohari M. Yamin^c and Mohammad B. Kassim^{c,a}

^aFuel Cell Institute, Universiti Kebangsaan Malaysia, UKM 43600 Bangi Selangor, Malaysia, ^bDepartment of Chemical and Process Engineering, Faculty of Engineering, Universiti Kebangsaan Malaysia, UKM 43600 Bangi Selangor, Malaysia, and ^cSchool of Chemical Sciences and Food Technology, Faculty of Science and Technology, Universiti Kebangsaan Malaysia, UKM 43600 Bangi Selangor, Malaysia
Correspondence e-mail: ibnhum@gmail.com


Received 28 June 2011; accepted 4 July 2011

Key indicators: single-crystal X-ray study; $T = 298$ K; mean $\sigma(\text{C}-\text{C}) = 0.005$ Å; R factor = 0.059; wR factor = 0.138; data-to-parameter ratio = 14.0.

In the title compound, $\text{C}_{22}\text{H}_{19}\text{FN}_2\text{OS}$, the 2-fluorobenzoyl group adopts a *trans* conformation with respect to the thiono S atom across the N—C bond. In the crystal, intermolecular N—H \cdots S, C—H \cdots S and C—H \cdots O hydrogen bonds link the molecules, forming a two-dimensional network parallel to (101).

Related literature

For standard bond lengths, see: Allen *et al.* (1987). For related structures, see: Nasir *et al.* (2011); Yamin & Hassan (2004); Hassan *et al.* (2008*a,b,c*, 2009). For the synthesis, see: Hassan *et al.* (2008*a*).



Experimental

Crystal data

$\text{C}_{22}\text{H}_{19}\text{FN}_2\text{OS}$
 $M_r = 378.45$
Monoclinic, $P2_1/n$

$a = 10.683$ (3) Å
 $b = 7.026$ (2) Å
 $c = 26.435$ (7) Å

$\beta = 101.100$ (6) $^\circ$
 $V = 1946.9$ (10) Å³
 $Z = 4$
Mo $K\alpha$ radiation

$\mu = 0.19$ mm⁻¹
 $T = 298$ K
 $0.42 \times 0.21 \times 0.18$ mm

Data collection

Bruker SMART APEX CCD area-detector diffractometer
Absorption correction: multi-scan (SADABS; Sheldrick, 2000)
 $T_{\min} = 0.925$, $T_{\max} = 0.967$

8644 measured reflections
3425 independent reflections
2659 reflections with $I > 2\sigma(I)$
 $R_{\text{int}} = 0.027$

Refinement

$R[F^2 > 2\sigma(F^2)] = 0.059$
 $wR(F^2) = 0.138$
 $S = 1.15$
3425 reflections

244 parameters
H-atom parameters constrained
 $\Delta\rho_{\text{max}} = 0.25$ e Å⁻³
 $\Delta\rho_{\text{min}} = -0.23$ e Å⁻³

Table 1

Hydrogen-bond geometry (Å, $^\circ$).

$D-H\cdots A$	$D-H$	$H\cdots A$	$D\cdots A$	$D-H\cdots A$
N1—H1 \cdots S1 ⁱ	0.86	2.61	3.422 (2)	159
C1—H1A \cdots S1 ⁱ	0.93	2.87	3.727 (3)	154
C4—H4 \cdots O1 ⁱⁱ	0.93	2.50	3.322 (4)	147

Symmetry codes: (i) $-x + 1, -y, -z$; (ii) $-x + \frac{3}{2}, y - \frac{1}{2}, -z + \frac{1}{2}$.

Data collection: SMART (Bruker, 2000); cell refinement: SAINT (Bruker, 2000); data reduction: SAINT; program(s) used to solve structure: SHELXS97 (Sheldrick, 2008); program(s) used to refine structure: SHELXL97 (Sheldrick, 2008); molecular graphics: ORTEP3 (Burnett & Johnson, 1996), XP in SHELXTL (Sheldrick, 2008) and PLATON (Spek, 2009); software used to prepare material for publication: SHELXTL and PLATON.

The authors thank Universiti Kebangsaan Malaysia for providing facilities and grants (postdoctoral for INH, UKM-GUP-BTT-07–30-190 and UKM-OUP-TK-16– 73/2010&2011 for MBK sabbatical leave). They also express their appreciation to the Kementerian Pengajian Tinggi, Malaysia, for the research fund No. UKM-ST-06-FRGS0111–2009.

Supplementary data and figures for this paper are available from the IUCr electronic archives (Reference: DN2702).

References

- Allen, F. H., Kennard, O., Watson, D. G., Brammer, L., Orpen, A. G. & Taylor, R. (1987). *J. Chem. Soc. Perkin Trans. 2*, pp. S1–19.
- Bruker (2000). SADABS, SMART and SAINT. Bruker AXS Inc., Madison, Wisconsin, USA.
- Burnett, M. N. & Johnson, C. K. (1996). ORTEP3. Report ORNL-6895. Oak Ridge National Laboratory, Tennessee, USA.
- Hassan, I. N., Yamin, B. M. & Kassim, M. B. (2008*a*). *Acta Cryst. E64*, o1727.
- Hassan, I. N., Yamin, B. M. & Kassim, M. B. (2008*b*). *Acta Cryst. E64*, o2083.
- Hassan, I. N., Yamin, B. M. & Kassim, M. B. (2008*c*). *Acta Cryst. E64*, o2167.
- Hassan, I. N., Yamin, B. M. & Kassim, M. B. (2009). *Acta Cryst. E65*, o3078.
- Nasir, M. F. M., Hassan, I. N., Wan Daud, W. R., Yamin, B. M. & Kassim, M. B. (2011). *Acta Cryst. E67*, o1218.
- Sheldrick, G. M. (2000). SADABS. University of Göttingen, Germany.
- Sheldrick, G. M. (2008). *Acta Cryst. A64*, 112–122.
- Spek, A. L. (2009). *Acta Cryst. D65*, 148–155.
- Yamin, B. M. & Hassan, I. N. (2004). *Acta Cryst. E60*, o2513–o2514.

supplementary materials

Acta Cryst. (2011). E67, o1987 [doi:10.1107/S1600536811026687]

1,1-Dibenzyl-3-(4-fluorobenzoyl)thiourea

M. F. M. Nasir, I. N. Hassan, W. R. Wan Daud, B. M. Yamin and M. B. Kassim

Comment

The title compound, I, is a thiourea derivative of dibenzylamine analogous to our previous reported, 1,1-sibenzyl-3-(3-chlorobenzoyl)thiourea, II (Nasir *et al.* 2011). The molecule maintains the the same *trans* and *cis* conformation for both the 3-fluorobenzoyl and the dibenzylamine groups, respectively, relative to the S atom across the N2—C8 bond (Fig 1). The dihedral angle between the phenyl ring, (C1—C6), and the thiourea fragment, (S1/N1/N2/C8) is $70.95(13)^\circ$, whereas in II was $72.9(2)^\circ$. The bond lengths and angles in the molecules are in normal ranges (Allen *et al.*, 1987) and comparable with those of II. However, the C=S bond length [$1.678(2)\text{\AA}$] is slightly longer than that of (II) [$1.672(6)\text{\AA}$]

Both phenyl rings, [C10/C11/C12/C13/C14/C15] and [C17/C18/C19/C20/C21/C22] are essentially planar and are twisted to each other by a dihedral angle of $22.4(4)^\circ$. The intermolecular N1—H1 \cdots S1, C1—H1A \cdots S and C4—H4 \cdots O1 hydrogen bonds (Table 1,) links the molecules into two dimensional ribbon parallel to the (1 0 1) plane (Fig 2).

Experimental

The title compound was synthesized according to a previously reported compound (Hassan *et al.*, 2008a). A colourless crystal, suitable for X-ray crystallography, was obtained by a slow evaporation from ethanolic solution at room temperature (yield 87%).

Refinement

H atoms of C and N atoms were positioned geometrically and allowed to ride on their parent atoms, with $U_{\text{iso}} = 1.2U_{\text{eq}}(\text{C})$ for aromatic 0.93\AA , $U_{\text{iso}} = 1.2U_{\text{eq}}(\text{C})$ for CH_2 0.97\AA , $U_{\text{iso}} = 1.2U_{\text{eq}}(\text{N})$ for N—H 0.86\AA .

Figures




Fig. 1. The molecular structure of (I), with the atoms labeling scheme and displacement ellipsoids are drawn at the 50% probability level. H atoms are represented as small spheres of arbitrary radii.




Fig. 2. Partial packing of (I) view down the b axis. H bonds are shown as dashed lines. H atoms not involved in hydrogen bondings have been omitted for clarity.

1,1-Dibenzyl-3-(4-fluorobenzoyl)thiourea

Crystal data

$C_{22}H_{19}FN_2OS$	$F(000) = 792$
$M_r = 378.45$	$D_x = 1.291 \text{ Mg m}^{-3}$
Monoclinic, $P2_1/n$	Melting point: 410 K
Hall symbol: -P 2yn	Mo $K\alpha$ radiation, $\lambda = 0.71073 \text{ \AA}$
$a = 10.683 (3) \text{ \AA}$	Cell parameters from 2659 reflections
$b = 7.026 (2) \text{ \AA}$	$\theta = 2.0\text{--}25.0^\circ$
$c = 26.435 (7) \text{ \AA}$	$\mu = 0.19 \text{ mm}^{-1}$
$\beta = 101.100 (6)^\circ$	$T = 298 \text{ K}$
$V = 1946.9 (10) \text{ \AA}^3$	Block, colourless
$Z = 4$	$0.42 \times 0.21 \times 0.18 \text{ mm}$

Data collection

Bruker SMART APEX CCD area-detector diffractometer	3425 independent reflections
Radiation source: fine-focus sealed tube graphite	2659 reflections with $I > 2\sigma(I)$
ω scans	$R_{\text{int}} = 0.027$
Absorption correction: multi-scan (SADABS; Sheldrick, 2000)	$\theta_{\text{max}} = 25.0^\circ$, $\theta_{\text{min}} = 2.0^\circ$
$T_{\text{min}} = 0.925$, $T_{\text{max}} = 0.967$	$h = -12 \rightarrow 8$
8644 measured reflections	$k = -8 \rightarrow 8$
	$l = -31 \rightarrow 28$

Refinement

Refinement on F^2	Primary atom site location: structure-invariant direct methods
Least-squares matrix: full	Secondary atom site location: difference Fourier map
$R[F^2 > 2\sigma(F^2)] = 0.059$	Hydrogen site location: inferred from neighbouring sites
$wR(F^2) = 0.138$	H-atom parameters constrained
$S = 1.15$	$w = 1/[\sigma^2(F_o^2) + (0.0542P)^2 + 0.5146P]$
3425 reflections	where $P = (F_o^2 + 2F_c^2)/3$
244 parameters	$(\Delta/\sigma)_{\text{max}} < 0.001$
0 restraints	$\Delta\rho_{\text{max}} = 0.25 \text{ e \AA}^{-3}$
	$\Delta\rho_{\text{min}} = -0.23 \text{ e \AA}^{-3}$

Special details

Geometry. All esds (except the esd in the dihedral angle between two l.s. planes) are estimated using the full covariance matrix. The cell esds are taken into account individually in the estimation of esds in distances, angles and torsion angles; correlations between esds

in cell parameters are only used when they are defined by crystal symmetry. An approximate (isotropic) treatment of cell esds is used for estimating esds involving l.s. planes.

Refinement. Refinement of F^2 against ALL reflections. The weighted R-factor wR and goodness of fit S are based on F^2 , conventional R-factors R are based on F, with F set to zero for negative F^2 . The threshold expression of $F^2 > 2\sigma(F^2)$ is used only for calculating R-factors(gt) etc. and is not relevant to the choice of reflections for refinement. R-factors based on F^2 are statistically about twice as large as those based on F, and R- factors based on ALL data will be even larger.

Fractional atomic coordinates and isotropic or equivalent isotropic displacement parameters (\AA^2)

	x	y	z	$U_{\text{iso}}^*/U_{\text{eq}}$
F1	0.8992 (2)	-0.7024 (3)	0.17795 (9)	0.1137 (8)
S1	0.47635 (8)	0.26603 (10)	0.02677 (3)	0.0641 (3)
O1	0.5719 (2)	0.0400 (3)	0.17142 (7)	0.0736 (6)
N1	0.5164 (2)	-0.0377 (3)	0.08698 (7)	0.0502 (6)
H1	0.5210	-0.1222	0.0639	0.060*
N2	0.3452 (2)	0.1514 (3)	0.09743 (7)	0.0453 (5)
C1	0.6626 (3)	-0.3835 (4)	0.11002 (10)	0.0570 (7)
H1A	0.6047	-0.3773	0.0789	0.068*
C2	0.7407 (3)	-0.5402 (5)	0.12067 (12)	0.0697 (8)
H2	0.7369	-0.6391	0.0970	0.084*
C3	0.8236 (3)	-0.5463 (5)	0.16670 (13)	0.0742 (9)
C4	0.8339 (3)	-0.4041 (6)	0.20216 (12)	0.0807 (10)
H4	0.8924	-0.4120	0.2331	0.097*
C5	0.7554 (3)	-0.2481 (5)	0.19116 (10)	0.0694 (9)
H5	0.7605	-0.1501	0.2151	0.083*
C6	0.6688 (2)	-0.2352 (4)	0.14481 (9)	0.0495 (6)
C7	0.5840 (2)	-0.0665 (4)	0.13658 (9)	0.0507 (7)
C8	0.4407 (3)	0.1238 (3)	0.07298 (8)	0.0460 (6)
C9	0.2742 (3)	0.3312 (4)	0.09357 (10)	0.0532 (7)
H9A	0.1842	0.3052	0.0815	0.064*
H9B	0.3027	0.4123	0.0683	0.064*
C10	0.2914 (3)	0.4349 (3)	0.14432 (9)	0.0482 (6)
C11	0.4099 (3)	0.4550 (4)	0.17540 (11)	0.0627 (8)
H11	0.4801	0.3968	0.1659	0.075*
C12	0.4261 (4)	0.5594 (5)	0.22019 (12)	0.0768 (10)
H12	0.5069	0.5718	0.2406	0.092*
C13	0.3240 (4)	0.6450 (4)	0.23477 (12)	0.0796 (10)
H13	0.3351	0.7164	0.2649	0.096*
C14	0.2052 (4)	0.6251 (4)	0.20481 (13)	0.0776 (10)
H14	0.1352	0.6823	0.2148	0.093*
C15	0.1887 (3)	0.5196 (4)	0.15949 (11)	0.0625 (8)
H15	0.1076	0.5062	0.1393	0.075*
C16	0.2919 (3)	0.0015 (4)	0.12613 (9)	0.0506 (7)
H16A	0.2790	0.0517	0.1589	0.061*
H16B	0.3517	-0.1036	0.1330	0.061*
C17	0.1667 (3)	-0.0689 (4)	0.09543 (9)	0.0521 (7)
C18	0.0539 (3)	-0.0329 (5)	0.11120 (12)	0.0716 (9)
H18	0.0539	0.0402	0.1405	0.086*

supplementary materials

C19	-0.0598 (3)	-0.1044 (7)	0.08383 (17)	0.0976 (12)
H19	-0.1357	-0.0802	0.0950	0.117*
C20	-0.0607 (5)	-0.2105 (7)	0.04053 (17)	0.1034 (14)
H20	-0.1371	-0.2595	0.0224	0.124*
C21	0.0496 (5)	-0.2446 (5)	0.02384 (13)	0.0921 (12)
H21	0.0482	-0.3150	-0.0060	0.111*
C22	0.1643 (3)	-0.1750 (4)	0.05112 (11)	0.0680 (8)
H22	0.2398	-0.1996	0.0397	0.082*

Atomic displacement parameters (\AA^2)

	U^{11}	U^{22}	U^{33}	U^{12}	U^{13}	U^{23}
F1	0.1048 (16)	0.1189 (18)	0.1105 (16)	0.0546 (14)	0.0040 (13)	0.0346 (14)
S1	0.1037 (6)	0.0484 (4)	0.0428 (4)	-0.0095 (4)	0.0210 (4)	0.0047 (3)
O1	0.0912 (15)	0.0817 (15)	0.0405 (10)	0.0164 (12)	-0.0058 (10)	-0.0154 (10)
N1	0.0710 (14)	0.0460 (12)	0.0316 (10)	0.0060 (11)	0.0050 (10)	-0.0024 (9)
N2	0.0605 (13)	0.0358 (11)	0.0379 (11)	0.0003 (10)	0.0055 (10)	0.0029 (8)
C1	0.0627 (18)	0.0552 (17)	0.0482 (15)	0.0041 (15)	-0.0013 (13)	0.0088 (13)
C2	0.075 (2)	0.063 (2)	0.0670 (19)	0.0095 (17)	0.0033 (16)	0.0058 (15)
C3	0.064 (2)	0.087 (2)	0.072 (2)	0.0230 (18)	0.0127 (17)	0.0279 (19)
C4	0.064 (2)	0.121 (3)	0.0510 (18)	0.020 (2)	-0.0025 (15)	0.020 (2)
C5	0.0630 (18)	0.099 (3)	0.0424 (15)	0.0080 (19)	0.0008 (14)	0.0019 (15)
C6	0.0500 (15)	0.0624 (17)	0.0360 (13)	-0.0023 (14)	0.0080 (11)	0.0092 (12)
C7	0.0567 (16)	0.0580 (17)	0.0350 (13)	-0.0042 (14)	0.0027 (12)	-0.0001 (12)
C8	0.0644 (17)	0.0389 (14)	0.0309 (12)	-0.0067 (13)	-0.0007 (12)	-0.0060 (10)
C9	0.0685 (18)	0.0413 (15)	0.0468 (14)	0.0062 (14)	0.0038 (13)	0.0037 (11)
C10	0.0657 (17)	0.0346 (13)	0.0452 (14)	-0.0024 (13)	0.0125 (13)	0.0041 (11)
C11	0.072 (2)	0.0539 (18)	0.0614 (17)	-0.0025 (15)	0.0122 (15)	-0.0097 (14)
C12	0.101 (3)	0.065 (2)	0.0594 (19)	-0.012 (2)	0.0029 (18)	-0.0140 (16)
C13	0.143 (3)	0.0475 (19)	0.0521 (18)	-0.005 (2)	0.029 (2)	-0.0051 (14)
C14	0.118 (3)	0.0539 (19)	0.074 (2)	0.018 (2)	0.051 (2)	0.0056 (16)
C15	0.076 (2)	0.0505 (17)	0.0659 (18)	0.0077 (16)	0.0252 (16)	0.0086 (14)
C16	0.0652 (17)	0.0459 (15)	0.0415 (13)	0.0018 (13)	0.0126 (12)	0.0053 (11)
C17	0.0682 (18)	0.0428 (15)	0.0455 (14)	-0.0060 (14)	0.0114 (13)	0.0071 (12)
C18	0.074 (2)	0.078 (2)	0.0655 (19)	-0.0069 (18)	0.0196 (17)	0.0071 (16)
C19	0.067 (2)	0.127 (4)	0.098 (3)	-0.021 (2)	0.014 (2)	0.022 (3)
C20	0.100 (3)	0.116 (3)	0.083 (3)	-0.051 (3)	-0.011 (2)	0.022 (2)
C21	0.135 (4)	0.074 (2)	0.060 (2)	-0.038 (3)	0.001 (2)	-0.0021 (17)
C22	0.092 (2)	0.0541 (17)	0.0564 (17)	-0.0092 (17)	0.0109 (16)	-0.0003 (14)

Geometric parameters (\AA , $^\circ$)

F1—C3	1.360 (4)	C10—C11	1.376 (4)
S1—C8	1.678 (3)	C11—C12	1.375 (4)
O1—C7	1.213 (3)	C11—H11	0.9300
N1—C7	1.385 (3)	C12—C13	1.365 (5)
N1—C8	1.401 (3)	C12—H12	0.9300
N1—H1	0.8600	C13—C14	1.367 (5)
N2—C8	1.323 (3)	C13—H13	0.9300

N2—C9	1.467 (3)	C14—C15	1.391 (4)
N2—C16	1.475 (3)	C14—H14	0.9300
C1—C2	1.378 (4)	C15—H15	0.9300
C1—C6	1.383 (4)	C16—C17	1.508 (4)
C1—H1A	0.9300	C16—H16A	0.9700
C2—C3	1.361 (4)	C16—H16B	0.9700
C2—H2	0.9300	C17—C18	1.372 (4)
C3—C4	1.360 (5)	C17—C22	1.385 (4)
C4—C5	1.376 (4)	C18—C19	1.383 (5)
C4—H4	0.9300	C18—H18	0.9300
C5—C6	1.389 (4)	C19—C20	1.364 (6)
C5—H5	0.9300	C19—H19	0.9300
C6—C7	1.482 (4)	C20—C21	1.356 (6)
C9—C10	1.507 (3)	C20—H20	0.9300
C9—H9A	0.9700	C21—C22	1.386 (5)
C9—H9B	0.9700	C21—H21	0.9300
C10—C15	1.375 (4)	C22—H22	0.9300
C7—N1—C8	122.6 (2)	C12—C11—C10	121.2 (3)
C7—N1—H1	118.7	C12—C11—H11	119.4
C8—N1—H1	118.7	C10—C11—H11	119.4
C8—N2—C9	122.0 (2)	C13—C12—C11	120.2 (3)
C8—N2—C16	123.8 (2)	C13—C12—H12	119.9
C9—N2—C16	113.9 (2)	C11—C12—H12	119.9
C2—C1—C6	121.1 (3)	C12—C13—C14	119.6 (3)
C2—C1—H1A	119.4	C12—C13—H13	120.2
C6—C1—H1A	119.4	C14—C13—H13	120.2
C3—C2—C1	118.2 (3)	C13—C14—C15	120.2 (3)
C3—C2—H2	120.9	C13—C14—H14	119.9
C1—C2—H2	120.9	C15—C14—H14	119.9
C4—C3—F1	118.4 (3)	C10—C15—C14	120.4 (3)
C4—C3—C2	123.0 (3)	C10—C15—H15	119.8
F1—C3—C2	118.5 (3)	C14—C15—H15	119.8
C3—C4—C5	118.3 (3)	N2—C16—C17	110.32 (19)
C3—C4—H4	120.8	N2—C16—H16A	109.6
C5—C4—H4	120.8	C17—C16—H16A	109.6
C4—C5—C6	120.9 (3)	N2—C16—H16B	109.6
C4—C5—H5	119.5	C17—C16—H16B	109.6
C6—C5—H5	119.5	H16A—C16—H16B	108.1
C1—C6—C5	118.4 (3)	C18—C17—C22	118.8 (3)
C1—C6—C7	123.7 (2)	C18—C17—C16	121.1 (3)
C5—C6—C7	117.8 (3)	C22—C17—C16	120.1 (3)
O1—C7—N1	121.1 (2)	C17—C18—C19	120.5 (3)
O1—C7—C6	122.2 (2)	C17—C18—H18	119.7
N1—C7—C6	116.7 (2)	C19—C18—H18	119.7
N2—C8—N1	116.7 (2)	C20—C19—C18	120.1 (4)
N2—C8—S1	124.9 (2)	C20—C19—H19	120.0
N1—C8—S1	118.3 (2)	C18—C19—H19	120.0
N2—C9—C10	112.4 (2)	C21—C20—C19	120.2 (4)
N2—C9—H9A	109.1	C21—C20—H20	119.9

supplementary materials

C10—C9—H9A	109.1	C19—C20—H20	119.9
N2—C9—H9B	109.1	C20—C21—C22	120.3 (4)
C10—C9—H9B	109.1	C20—C21—H21	119.8
H9A—C9—H9B	107.8	C22—C21—H21	119.8
C15—C10—C11	118.4 (3)	C17—C22—C21	120.0 (3)
C15—C10—C9	120.1 (3)	C17—C22—H22	120.0
C11—C10—C9	121.5 (3)	C21—C22—H22	120.0

Hydrogen-bond geometry (Å, °)

<i>D</i> —H \cdots <i>A</i>	<i>D</i> —H	H \cdots <i>A</i>	<i>D</i> \cdots <i>A</i>	<i>D</i> —H \cdots <i>A</i>
C9—H9B \cdots S1	0.97	2.55	3.076 (3)	114
N1—H1 \cdots S1 ⁱ	0.86	2.61	3.422 (2)	159
C1—H1A \cdots S1 ⁱ	0.93	2.87	3.727 (3)	154
C4—H4 \cdots O1 ⁱⁱ	0.93	2.50	3.322 (4)	147

Symmetry codes: (i) $-x+1, -y, -z$; (ii) $-x+3/2, y-1/2, -z+1/2$.

Fig. 1




Fig. 2

