

2,2-Diphenylacetamide

Jerry P. Jasinski,^{a*} James A. Golen,^a M. S. Siddegowda,^b
H. S. Yathirajan^b and M. T. Swamy^c

^aDepartment of Chemistry, Keene State College, 229 Main Street, Keene, NH 03435-2001, USA, ^bDepartment of Studies in Chemistry, University of Mysore, Manasagangotri, Mysore 570 006, India, and ^cDepartment of Chemistry, Sambhram Institute of Technology, Bangalore 560 097, India

Correspondence e-mail: jjasinski@keene.edu


Received 3 July 2011; accepted 5 July 2011

Key indicators: single-crystal X-ray study; $T = 173$ K; mean $\sigma(\text{C}-\text{C}) = 0.003$ Å; R factor = 0.037; wR factor = 0.107; data-to-parameter ratio = 13.1.

In the title compound, $\text{C}_{14}\text{H}_{13}\text{NO}$, which has two molecules in the asymmetric unit, the dihedral angles between the mean planes of the benzene rings are 84.6 (7) and 85.0 (6)°. $\text{N}-\text{H}\cdots\text{O}$ hydrogen bonds [forming $R_2^2(8)$ ring motifs] and $\text{C}-\text{H}\cdots\text{O}$ hydrogen bonds dominate the crystal packing, forming zigzag chains parallel to the a axis. In addition, weak intermolecular $\text{C}-\text{H}\cdots\pi$ interactions are observed.

Related literature

For the synthesis and antimycobacterial activity of 2,2-diphenylacetamide derivatives, see: Guzel *et al.* (2006). For related structures, see: Akkurt *et al.* (2007); Dutkiewicz *et al.* (2010); Gerkin (1998); Krigbaum *et al.* (1968); Narasegowda *et al.* (2005); Yathirajan *et al.* (2005). For standard bond lengths, see: Allen *et al.* (1987).



Experimental

Crystal data

$\text{C}_{14}\text{H}_{13}\text{NO}$	$V = 1139.52$ (10) Å ³
$M_r = 211.25$	$Z = 4$
Monoclinic, $P2_1$	Cu $K\alpha$ radiation
$a = 5.1687$ (3) Å	$\mu = 0.61$ mm ⁻¹
$b = 28.5511$ (13) Å	$T = 173$ K
$c = 7.8006$ (4) Å	$0.40 \times 0.25 \times 0.20$ mm
$\beta = 98.152$ (5)°	

Data collection

Oxford Diffraction Xcalibur Eos Gemini diffractometer	6490 measured reflections
Absorption correction: multi-scan (<i>CrysAlis RED</i> ; Oxford Diffraction, 2010)	3978 independent reflections
$T_{\min} = 0.792$, $T_{\max} = 0.887$	3869 reflections with $I > 2\sigma(I)$
	$R_{\text{int}} = 0.017$

Refinement

$R[F^2 > 2\sigma(F^2)] = 0.037$	H atoms treated by a mixture of independent and constrained refinement
$wR(F^2) = 0.107$	$\Delta\rho_{\text{max}} = 0.21$ e Å ⁻³
$S = 1.07$	$\Delta\rho_{\text{min}} = -0.20$ e Å ⁻³
3978 reflections	
303 parameters	
9 restraints	

Table 1

Hydrogen-bond geometry (Å, °).

Cg1 and Cg4 are the centroids of the $\text{C3}-\text{C8}$ and $\text{C23}-\text{C28}$ rings, respectively.

$D-\text{H}\cdots A$	$D-\text{H}$	$\text{H}\cdots A$	$D\cdots A$	$D-\text{H}\cdots A$
$\text{N1}-\text{H1B}\cdots\text{O1}^{\text{i}}$	0.89 (1)	2.20 (2)	2.9409 (17)	140 (2)
$\text{N1}-\text{H1A}\cdots\text{O2}^{\text{ii}}$	0.87 (1)	2.09 (1)	2.9575 (19)	177 (2)
$\text{N2}-\text{H2B}\cdots\text{O1}^{\text{iii}}$	0.88 (1)	2.07 (1)	2.9526 (19)	176 (2)
$\text{N2}-\text{H2A}\cdots\text{O2}^{\text{iv}}$	0.89 (1)	2.17 (2)	2.9407 (18)	145 (2)
$\text{N1}-\text{H1A}\cdots\text{N2}^{\text{ii}}$	0.87 (1)	3.06 (2)	3.7246 (18)	134 (2)
$\text{N2}-\text{H2B}\cdots\text{N1}^{\text{iii}}$	0.88 (1)	3.10 (2)	3.7246 (18)	130 (2)
$\text{C10}-\text{H10A}\cdots\text{O1}$	0.95	2.50	3.093 (2)	120
$\text{C18}-\text{H18A}\cdots\text{O2}$	0.95	2.51	3.099 (2)	120
$\text{C2}-\text{H2C}\cdots\text{Cg1}^{\text{i}}$	1.00	2.96	3.9379 (18)	165
$\text{C16}-\text{H16A}\cdots\text{Cg4}^{\text{iv}}$	1.00	2.95	3.9263 (18)	166

Symmetry codes: (i) $x + 1, y, z$; (ii) $x, y, z + 1$; (iii) $x, y, z - 1$; (iv) $x - 1, y, z$.

Data collection: *CrysAlis PRO* (Oxford Diffraction, 2010); cell refinement: *CrysAlis PRO*; data reduction: *CrysAlis RED* (Oxford Diffraction, 2010); program(s) used to solve structure: *SHELXS97* (Sheldrick, 2008); program(s) used to refine structure: *SHELXL97* (Sheldrick, 2008); molecular graphics: *SHELXTL* (Sheldrick, 2008); software used to prepare material for publication: *SHELXTL*.

JPJ acknowledges the NSF-MRI program (grant No. CHE1039027) for funds to purchase the X-ray diffractometer. MSS thanks the University of Mysore for the research facilities and R. L. Fine Chem, Bangalore, India, for the gift sample.

Supplementary data and figures for this paper are available from the IUCr electronic archives (Reference: JH2307).

References

- Akkurt, M., Karaca, S., Şahin, E., Güzel, Ö., Salman, A. & İlhan, E. (2007). *Acta Cryst.* **E63**, o3379–o3380.
- Allen, F. H., Kennard, O., Watson, D. G., Brammer, L., Orpen, A. G. & Taylor, R. (1987). *J. Chem. Soc. Perkin Trans. 2*, pp. S1–19.
- Dutkiewicz, G., Siddaraju, B. P., Yathirajan, H. S., Narayana, B. & Kubicki, M. (2010). *Acta Cryst.* **E66**, o499.
- Gerkin, R. E. (1998). *Acta Cryst.* **C54**, 1887–1889.
- Guzel, O., İlhan, E. & Salman, A. (2006). *Monatsh. Chem.* **137**, 795–801.
- Krigbaum, W. R., Roe, R.-J. & Woods, J. D. (1968). *Acta Cryst.* **B24**, 1304–1312.
- Narasegowda, R. S., Yathirajan, H. S. & Bolte, M. (2005). *Acta Cryst.* **E61**, o939–o940.

Oxford Diffraction (2010). *CrysAlis PRO* and *CrysAlis RED*. Oxford Diffraction Ltd, Yarnton, England.
Sheldrick, G. M. (2008). *Acta Cryst.* **A64**, 112–122.

Yathirajan, H. S., Nagaraj, B., Narasegowda, R. S., Nagaraja, P. & Bolte, M. (2005). *Acta Cryst.* **E61**, o1193–o1195.

supplementary materials

Acta Cryst. (2011). E67, o1992-o1993 [doi:10.1107/S1600536811026717]

2,2-Diphenylacetamide

J. P. Jasinski, J. A. Golen, M. S. Siddegowda, H. S. Yathirajan and M. T. Swamy

Comment

The synthesis and antimycobacterial activity of some new 2,2-diphenylacetamide derivatives is described (Guzel *et al.*, 2006). The title compound is used to synthesize various biologically active and pharmaceutical compounds viz., loperamide, darifenacin, fempiverine, etc. The crystal structures of N,N-diphenylacetamide (Krigbaum *et al.*, 1968), 4,4'-dimethylbiphenyl-2,2'-dicarboxylic acid (Gerkin, 1998), 4'-methylbiphenyl-2-carboxylic acid (Narasegowda *et al.*, 2005) and 4'-(2-butyl-4-chloro-5-formylimidazol-1-ylmethyl)biphenyl-2-carbonitrile (Yathirajan *et al.*, 2005), 2-hydroxy-N-(3-oxo-1-thia-4-azaspiro[4.5]dec-4-yl)-2,2-diphenylacetamide (Akkurt *et al.*, 2007) and 2-Chloro-N-[4-chloro-2-(2-chlorobenzoyl)phenyl]acetamide (Dutkiewicz *et al.*, 2010) have been reported. In view of the importance of the title compound, (I), C₁₄H₁₃NO, and in order to determine the conformation of this molecule, a crystal structure determination has been carried out.

In the title compound, (I), with two molecules in the asymmetric unit, the dihedral angle between the mean planes of the benzene rings is 84.6 (7)° or 85.0 (6)°, respectively (Fig. 1). Extensive N—H···O (forming an R₂²(8) ring-motif) intermolecular and C—H···O intramolecular hydrogen bonds and weak C—H···Cg π -ring intermolecular interactions (dominate the crystal packing forming a zigzag chain along [010] (Table 1, Fig. 2).

Experimental

The title compound was obtained as a gift sample from R. L. Fine Chem, Bangalore. X-ray quality crystals were obtained by slow evaporation of 1:1 acetone:methanol solution (m.p.: 430-433 K).

Refinement

The N—H atoms were located by Fourier analysis and refined isotropically with DFIX = 0.87Å. All of the remaining H atoms were placed in their calculated positions and then refined using the riding model with Atom—H lengths of 0.95Å or 1.00Å (CH). Isotropic displacement parameters for these atoms were set to 1.19-1.20 (CH), times U_{eq} of the parent atom.

Figures




Fig. 1. Molecular structure of the title compound showing the atom labeling scheme and 50% probability displacement ellipsoids.




Fig. 2. Packing diagram of the title compound viewed down the *a* axis. Dashed lines represent N—H...O hydrogen bonds.

2,2-Diphenylacetamide

Crystal data

$C_{14}H_{13}NO$

$M_r = 211.25$

Monoclinic, $P2_1$

Hall symbol: P 2yb

$a = 5.1687$ (3) Å

$b = 28.5511$ (13) Å

$c = 7.8006$ (4) Å

$\beta = 98.152$ (5)°

$V = 1139.52$ (10) Å³

$Z = 4$

$F(000) = 448$

$D_x = 1.231$ Mg m⁻³

Cu $K\alpha$ radiation, $\lambda = 1.54178$ Å

Cell parameters from 5082 reflections

$\theta = 4.6$ – 70.4 °

$\mu = 0.61$ mm⁻¹

$T = 173$ K

Block, colorless

$0.40 \times 0.25 \times 0.20$ mm

Data collection

Oxford Diffraction Xcalibur Eos Gemini diffractometer

Radiation source: Enhance (Cu) X-ray Source graphite

Detector resolution: 16.1500 pixels mm⁻¹

ω scans

Absorption correction: multi-scan (*CrysAlis RED*; Oxford Diffraction, 2010)

$T_{\min} = 0.792$, $T_{\max} = 0.887$

6490 measured reflections

3978 independent reflections

3869 reflections with $I > 2\sigma(I)$

$R_{\text{int}} = 0.017$

$\theta_{\max} = 70.5$ °, $\theta_{\min} = 5.7$ °

$h = -6$ → 6

$k = -34$ → 34

$l = -9$ → 8

Refinement

Refinement on F^2

Least-squares matrix: full

$R[F^2 > 2\sigma(F^2)] = 0.037$

$wR(F^2) = 0.107$

$S = 1.07$

Secondary atom site location: difference Fourier map

Hydrogen site location: inferred from neighbouring sites

H atoms treated by a mixture of independent and constrained refinement

$w = 1/[\sigma^2(F_o^2) + (0.0695P)^2 + 0.1057P]$

where $P = (F_o^2 + 2F_c^2)/3$

$(\Delta/\sigma)_{\max} = 0.027$

3978 reflections	$\Delta\rho_{\max} = 0.21 \text{ e } \text{\AA}^{-3}$
303 parameters	$\Delta\rho_{\min} = -0.20 \text{ e } \text{\AA}^{-3}$
9 restraints	Extinction correction: <i>SHELXL97</i> (Sheldrick, 2008), $F_c^* = kFc[1+0.001xFc^2\lambda^3/\sin(2\theta)]^{-1/4}$
Primary atom site location: structure-invariant direct methods	Extinction coefficient: 0.0118 (13)

Special details

Geometry. All esds (except the esd in the dihedral angle between two l.s. planes) are estimated using the full covariance matrix. The cell esds are taken into account individually in the estimation of esds in distances, angles and torsion angles; correlations between esds in cell parameters are only used when they are defined by crystal symmetry. An approximate (isotropic) treatment of cell esds is used for estimating esds involving l.s. planes.

Refinement. Refinement of F^2 against ALL reflections. The weighted R -factor wR and goodness of fit S are based on F^2 , conventional R -factors R are based on F , with F set to zero for negative F^2 . The threshold expression of $F^2 > \sigma(F^2)$ is used only for calculating R -factors(gt) etc. and is not relevant to the choice of reflections for refinement. R -factors based on F^2 are statistically about twice as large as those based on F , and R -factors based on ALL data will be even larger.

Fractional atomic coordinates and isotropic or equivalent isotropic displacement parameters (\AA^2)

	<i>x</i>	<i>y</i>	<i>z</i>	$U_{\text{iso}}^*/U_{\text{eq}}$
N1	0.9797 (3)	0.54261 (6)	0.6365 (2)	0.0459 (3)
H1B	1.121 (3)	0.5307 (8)	0.600 (3)	0.055*
H1A	0.984 (5)	0.5667 (6)	0.706 (3)	0.055*
O2	1.0081 (2)	0.62240 (4)	-0.11977 (17)	0.0465 (3)
O1	0.5496 (2)	0.53784 (5)	0.64407 (16)	0.0459 (3)
N2	0.5779 (3)	0.61736 (6)	-0.1120 (2)	0.0469 (4)
H2B	0.561 (5)	0.5935 (6)	-0.185 (3)	0.056*
H2A	0.431 (3)	0.6281 (9)	-0.080 (3)	0.056*
C1	0.7494 (3)	0.52226 (5)	0.5965 (2)	0.0359 (3)
C2	0.7440 (3)	0.47718 (6)	0.4902 (2)	0.0377 (3)
H2C	0.9097	0.4758	0.4381	0.045*
C3	0.5167 (3)	0.47744 (6)	0.3416 (2)	0.0406 (4)
C4	0.4550 (5)	0.51789 (8)	0.2469 (3)	0.0678 (6)
H4A	0.5534	0.5456	0.2759	0.081*
C5	0.2520 (6)	0.51860 (10)	0.1106 (4)	0.0824 (8)
H5A	0.2130	0.5467	0.0469	0.099*
C6	0.1062 (5)	0.47893 (10)	0.0666 (3)	0.0683 (6)
H6A	-0.0353	0.4796	-0.0254	0.082*
C7	0.1684 (4)	0.43851 (9)	0.1575 (3)	0.0577 (5)
H7A	0.0707	0.4108	0.1272	0.069*
C8	0.3724 (4)	0.43764 (7)	0.2933 (2)	0.0461 (4)
H8A	0.4138	0.4092	0.3543	0.055*
C9	0.7419 (3)	0.43519 (5)	0.6105 (2)	0.0375 (3)
C10	0.5760 (4)	0.43225 (7)	0.7334 (3)	0.0518 (4)
H10A	0.4577	0.4571	0.7451	0.062*
C11	0.5795 (5)	0.39352 (9)	0.8402 (3)	0.0606 (5)

supplementary materials

H11A	0.4630	0.3921	0.9239	0.073*
C12	0.7488 (5)	0.35715 (8)	0.8270 (3)	0.0607 (5)
H12A	0.7521	0.3309	0.9020	0.073*
C13	0.9135 (5)	0.35925 (9)	0.7036 (4)	0.0738 (7)
H13A	1.0288	0.3340	0.6910	0.089*
C14	0.9117 (4)	0.39807 (8)	0.5981 (3)	0.0607 (5)
H14A	1.0294	0.3994	0.5151	0.073*
C15	0.8083 (3)	0.63780 (6)	-0.0712 (2)	0.0355 (3)
C16	0.8143 (3)	0.68289 (5)	0.0360 (2)	0.0367 (3)
H16A	0.6495	0.6841	0.0892	0.044*
C17	0.8148 (3)	0.72535 (6)	-0.0840 (2)	0.0373 (3)
C18	0.9825 (4)	0.72830 (7)	-0.2076 (2)	0.0505 (4)
H18A	1.1005	0.7034	-0.2201	0.061*
C19	0.9779 (4)	0.76758 (8)	-0.3129 (3)	0.0594 (5)
H19A	1.0941	0.7694	-0.3967	0.071*
C20	0.8089 (5)	0.80352 (7)	-0.2976 (3)	0.0585 (5)
H20A	0.8061	0.8301	-0.3710	0.070*
C21	0.6427 (5)	0.80100 (8)	-0.1751 (3)	0.0654 (6)
H21A	0.5257	0.8260	-0.1627	0.078*
C22	0.6461 (4)	0.76191 (7)	-0.0699 (3)	0.0533 (5)
H22A	0.5294	0.7603	0.0137	0.064*
C23	1.0422 (3)	0.68233 (6)	0.1827 (2)	0.0404 (3)
C24	1.1076 (5)	0.64165 (8)	0.2764 (3)	0.0702 (6)
H24A	1.0119	0.6137	0.2462	0.084*
C25	1.3098 (6)	0.64110 (11)	0.4128 (3)	0.0848 (8)
H25A	1.3508	0.6129	0.4757	0.102*
C26	1.4513 (5)	0.68078 (11)	0.4580 (3)	0.0696 (6)
H26A	1.5922	0.6801	0.5505	0.084*
C27	1.3884 (4)	0.72136 (9)	0.3690 (3)	0.0608 (5)
H27A	1.4847	0.7491	0.4008	0.073*
C28	1.1853 (4)	0.72226 (7)	0.2329 (2)	0.0468 (4)
H28A	1.1432	0.7508	0.1727	0.056*

Atomic displacement parameters (\AA^2)

	U^{11}	U^{22}	U^{33}	U^{12}	U^{13}	U^{23}
N1	0.0328 (6)	0.0411 (8)	0.0651 (9)	-0.0033 (6)	0.0110 (6)	-0.0140 (7)
O2	0.0316 (5)	0.0435 (7)	0.0654 (7)	-0.0009 (5)	0.0106 (5)	-0.0159 (5)
O1	0.0321 (5)	0.0424 (6)	0.0639 (7)	-0.0001 (5)	0.0093 (5)	-0.0150 (6)
N2	0.0325 (6)	0.0435 (8)	0.0652 (9)	-0.0021 (6)	0.0086 (6)	-0.0147 (7)
C1	0.0322 (7)	0.0325 (8)	0.0430 (8)	0.0009 (5)	0.0051 (6)	-0.0007 (6)
C2	0.0361 (7)	0.0352 (8)	0.0436 (8)	-0.0012 (6)	0.0116 (6)	-0.0032 (7)
C3	0.0436 (8)	0.0388 (9)	0.0407 (8)	-0.0019 (6)	0.0107 (6)	-0.0039 (7)
C4	0.0871 (15)	0.0441 (11)	0.0663 (13)	-0.0108 (10)	-0.0092 (11)	0.0077 (9)
C5	0.107 (2)	0.0650 (15)	0.0660 (14)	-0.0016 (14)	-0.0202 (14)	0.0137 (12)
C6	0.0713 (13)	0.0847 (16)	0.0442 (10)	0.0011 (12)	-0.0085 (9)	-0.0065 (10)
C7	0.0624 (11)	0.0665 (13)	0.0438 (9)	-0.0133 (10)	0.0061 (8)	-0.0125 (9)
C8	0.0545 (9)	0.0452 (10)	0.0398 (8)	-0.0061 (8)	0.0109 (7)	-0.0068 (7)

C9	0.0368 (7)	0.0332 (8)	0.0423 (8)	-0.0013 (6)	0.0048 (6)	-0.0052 (6)
C10	0.0547 (10)	0.0487 (11)	0.0557 (10)	0.0113 (8)	0.0204 (8)	0.0036 (8)
C11	0.0734 (13)	0.0576 (11)	0.0565 (11)	0.0015 (10)	0.0284 (10)	0.0067 (10)
C12	0.0769 (13)	0.0434 (11)	0.0636 (12)	0.0004 (10)	0.0164 (10)	0.0106 (9)
C13	0.0837 (15)	0.0440 (11)	0.1012 (18)	0.0193 (11)	0.0389 (14)	0.0163 (11)
C14	0.0657 (12)	0.0439 (10)	0.0795 (14)	0.0133 (9)	0.0346 (10)	0.0054 (10)
C15	0.0327 (7)	0.0324 (7)	0.0418 (7)	-0.0002 (6)	0.0066 (6)	-0.0006 (6)
C16	0.0357 (7)	0.0337 (8)	0.0430 (8)	-0.0020 (6)	0.0129 (6)	-0.0041 (7)
C17	0.0363 (7)	0.0358 (8)	0.0399 (7)	-0.0021 (6)	0.0055 (6)	-0.0057 (6)
C18	0.0531 (10)	0.0481 (10)	0.0534 (10)	0.0080 (8)	0.0186 (8)	0.0042 (8)
C19	0.0686 (12)	0.0591 (12)	0.0550 (11)	0.0016 (10)	0.0245 (9)	0.0069 (10)
C20	0.0752 (13)	0.0429 (11)	0.0583 (11)	-0.0010 (9)	0.0123 (10)	0.0104 (8)
C21	0.0786 (15)	0.0409 (10)	0.0818 (15)	0.0171 (9)	0.0294 (12)	0.0066 (10)
C22	0.0597 (10)	0.0404 (9)	0.0655 (11)	0.0073 (8)	0.0291 (9)	0.0050 (8)
C23	0.0451 (8)	0.0399 (9)	0.0381 (7)	-0.0005 (6)	0.0123 (6)	-0.0036 (7)
C24	0.0878 (16)	0.0478 (12)	0.0684 (13)	-0.0088 (11)	-0.0116 (12)	0.0077 (10)
C25	0.109 (2)	0.0685 (16)	0.0669 (14)	0.0064 (15)	-0.0228 (14)	0.0127 (12)
C26	0.0741 (13)	0.0856 (16)	0.0441 (10)	0.0001 (12)	-0.0091 (9)	-0.0043 (11)
C27	0.0643 (12)	0.0720 (14)	0.0462 (9)	-0.0175 (10)	0.0081 (9)	-0.0141 (10)
C28	0.0553 (10)	0.0452 (10)	0.0411 (8)	-0.0090 (8)	0.0112 (7)	-0.0057 (7)

Geometric parameters (Å, °)

N1—C1	1.322 (2)	C12—H12A	0.9500
N1—H1B	0.889 (13)	C13—C14	1.380 (3)
N1—H1A	0.873 (13)	C13—H13A	0.9500
O2—C15	1.2306 (19)	C14—H14A	0.9500
O1—C1	1.2292 (19)	C15—C16	1.533 (2)
N2—C15	1.324 (2)	C16—C23	1.521 (2)
N2—H2B	0.883 (13)	C16—C17	1.532 (2)
N2—H2A	0.889 (13)	C16—H16A	1.0000
C1—C2	1.529 (2)	C17—C22	1.375 (2)
C2—C9	1.523 (2)	C17—C18	1.387 (2)
C2—C3	1.529 (2)	C18—C19	1.388 (3)
C2—H2C	1.0000	C18—H18A	0.9500
C3—C8	1.382 (2)	C19—C20	1.364 (3)
C3—C4	1.384 (3)	C19—H19A	0.9500
C4—C5	1.384 (3)	C20—C21	1.374 (4)
C4—H4A	0.9500	C20—H20A	0.9500
C5—C6	1.377 (4)	C21—C22	1.384 (3)
C5—H5A	0.9500	C21—H21A	0.9500
C6—C7	1.369 (4)	C22—H22A	0.9500
C6—H6A	0.9500	C23—C28	1.385 (2)
C7—C8	1.385 (3)	C23—C24	1.389 (3)
C7—H7A	0.9500	C24—C25	1.382 (3)
C8—H8A	0.9500	C24—H24A	0.9500
C9—C10	1.377 (3)	C25—C26	1.367 (4)
C9—C14	1.388 (3)	C25—H25A	0.9500
C10—C11	1.383 (3)	C26—C27	1.366 (4)

supplementary materials

C10—H10A	0.9500	C26—H26A	0.9500
C11—C12	1.371 (3)	C27—C28	1.384 (3)
C11—H11A	0.9500	C27—H27A	0.9500
C12—C13	1.373 (4)	C28—H28A	0.9500
C1—N1—H1B	120.8 (16)	C13—C14—C9	121.52 (19)
C1—N1—H1A	115.9 (16)	C13—C14—H14A	119.2
H1B—N1—H1A	123 (2)	C9—C14—H14A	119.2
C15—N2—H2B	119.8 (16)	O2—C15—N2	122.28 (16)
C15—N2—H2A	123.8 (16)	O2—C15—C16	120.98 (13)
H2B—N2—H2A	116 (2)	N2—C15—C16	116.72 (13)
O1—C1—N1	122.46 (15)	C23—C16—C17	113.57 (13)
O1—C1—C2	121.11 (14)	C23—C16—C15	110.82 (13)
N1—C1—C2	116.42 (14)	C17—C16—C15	109.44 (12)
C9—C2—C1	109.27 (13)	C23—C16—H16A	107.6
C9—C2—C3	113.52 (13)	C17—C16—H16A	107.6
C1—C2—C3	111.11 (13)	C15—C16—H16A	107.6
C9—C2—H2C	107.6	C22—C17—C18	118.40 (16)
C1—C2—H2C	107.6	C22—C17—C16	119.72 (14)
C3—C2—H2C	107.6	C18—C17—C16	121.88 (14)
C8—C3—C4	117.79 (17)	C17—C18—C19	120.03 (18)
C8—C3—C2	121.97 (15)	C17—C18—H18A	120.0
C4—C3—C2	120.21 (16)	C19—C18—H18A	120.0
C5—C4—C3	121.0 (2)	C20—C19—C18	120.89 (19)
C5—C4—H4A	119.5	C20—C19—H19A	119.6
C3—C4—H4A	119.5	C18—C19—H19A	119.6
C4—C5—C6	120.5 (2)	C19—C20—C21	119.51 (19)
C4—C5—H5A	119.7	C19—C20—H20A	120.2
C6—C5—H5A	119.7	C21—C20—H20A	120.2
C7—C6—C5	119.0 (2)	C20—C21—C22	119.9 (2)
C7—C6—H6A	120.5	C20—C21—H21A	120.1
C5—C6—H6A	120.5	C22—C21—H21A	120.1
C6—C7—C8	120.59 (19)	C17—C22—C21	121.31 (19)
C6—C7—H7A	119.7	C17—C22—H22A	119.3
C8—C7—H7A	119.7	C21—C22—H22A	119.3
C3—C8—C7	121.11 (18)	C28—C23—C24	117.35 (17)
C3—C8—H8A	119.4	C28—C23—C16	121.97 (15)
C7—C8—H8A	119.4	C24—C23—C16	120.64 (16)
C10—C9—C14	117.65 (17)	C25—C24—C23	121.1 (2)
C10—C9—C2	122.44 (15)	C25—C24—H24A	119.5
C14—C9—C2	119.91 (15)	C23—C24—H24A	119.5
C9—C10—C11	120.79 (18)	C26—C25—C24	120.5 (2)
C9—C10—H10A	119.6	C26—C25—H25A	119.7
C11—C10—H10A	119.6	C24—C25—H25A	119.7
C12—C11—C10	120.97 (19)	C25—C26—C27	119.4 (2)
C12—C11—H11A	119.5	C25—C26—H26A	120.3
C10—C11—H11A	119.5	C27—C26—H26A	120.3
C11—C12—C13	119.0 (2)	C26—C27—C28	120.4 (2)
C11—C12—H12A	120.5	C26—C27—H27A	119.8
C13—C12—H12A	120.5	C28—C27—H27A	119.8

C12—C13—C14	120.0 (2)	C27—C28—C23	121.22 (18)
C12—C13—H13A	120.0	C27—C28—H28A	119.4
C14—C13—H13A	120.0	C23—C28—H28A	119.4
O1—C1—C2—C9	-79.23 (19)	O2—C15—C16—C23	-46.8 (2)
N1—C1—C2—C9	99.42 (17)	N2—C15—C16—C23	135.00 (15)
O1—C1—C2—C3	46.8 (2)	O2—C15—C16—C17	79.24 (18)
N1—C1—C2—C3	-134.55 (16)	N2—C15—C16—C17	-99.00 (16)
C9—C2—C3—C8	-17.6 (2)	C23—C16—C17—C22	-103.20 (18)
C1—C2—C3—C8	-141.21 (16)	C15—C16—C17—C22	132.39 (17)
C9—C2—C3—C4	164.54 (19)	C23—C16—C17—C18	76.54 (19)
C1—C2—C3—C4	40.9 (2)	C15—C16—C17—C18	-47.87 (19)
C8—C3—C4—C5	1.2 (4)	C22—C17—C18—C19	0.2 (3)
C2—C3—C4—C5	179.2 (2)	C16—C17—C18—C19	-179.50 (18)
C3—C4—C5—C6	0.2 (5)	C17—C18—C19—C20	-0.4 (3)
C4—C5—C6—C7	-1.3 (5)	C18—C19—C20—C21	0.6 (4)
C5—C6—C7—C8	0.9 (4)	C19—C20—C21—C22	-0.7 (4)
C4—C3—C8—C7	-1.6 (3)	C18—C17—C22—C21	-0.3 (3)
C2—C3—C8—C7	-179.55 (17)	C16—C17—C22—C21	179.4 (2)
C6—C7—C8—C3	0.6 (3)	C20—C21—C22—C17	0.6 (4)
C1—C2—C9—C10	47.5 (2)	C17—C16—C23—C28	18.5 (2)
C3—C2—C9—C10	-77.1 (2)	C15—C16—C23—C28	142.15 (16)
C1—C2—C9—C14	-132.47 (18)	C17—C16—C23—C24	-164.12 (19)
C3—C2—C9—C14	102.9 (2)	C15—C16—C23—C24	-40.5 (2)
C14—C9—C10—C11	-0.1 (3)	C28—C23—C24—C25	-0.8 (4)
C2—C9—C10—C11	179.91 (18)	C16—C23—C24—C25	-178.3 (2)
C9—C10—C11—C12	0.3 (3)	C23—C24—C25—C26	-0.4 (5)
C10—C11—C12—C13	-1.0 (4)	C24—C25—C26—C27	1.1 (5)
C11—C12—C13—C14	1.6 (4)	C25—C26—C27—C28	-0.8 (4)
C12—C13—C14—C9	-1.4 (4)	C26—C27—C28—C23	-0.4 (3)
C10—C9—C14—C13	0.7 (3)	C24—C23—C28—C27	1.1 (3)
C2—C9—C14—C13	-179.3 (2)	C16—C23—C28—C27	178.60 (17)

Hydrogen-bond geometry (\AA , $^\circ$)

Cg1 and Cg4 are the centroids of the C3—C8 and C23—C28 rings, respectively.

$D-H\cdots A$	$D-H$	$H\cdots A$	$D\cdots A$	$D-H\cdots A$
N1—H1B \cdots O1 ⁱ	0.89 (1)	2.20 (2)	2.9409 (17)	140 (2)
N1—H1A \cdots O2 ⁱⁱ	0.87 (1)	2.09 (1)	2.9575 (19)	177 (2)
N2—H2B \cdots O1 ⁱⁱⁱ	0.88 (1)	2.07 (1)	2.9526 (19)	176 (2)
N2—H2A \cdots O2 ^{iv}	0.89 (1)	2.17 (2)	2.9407 (18)	145 (2)
N1—H1A \cdots N2 ⁱⁱ	0.87 (1)	3.06 (2)	3.7246 (18)	134.(2)
N2—H2B \cdots N1 ⁱⁱⁱ	0.88 (1)	3.10 (2)	3.7246 (18)	130.(2)
C10—H10A \cdots O1	0.95	2.50	3.093 (2)	120.
C18—H18A \cdots O2	0.95	2.51	3.099 (2)	120
C2—H2C \cdots Cg1 ⁱ	1.00	2.96	3.9379 (18)	165
C16—H16A \cdots Cg4 ^{iv}	1.00	2.95	3.9263 (18)	166

Symmetry codes: (i) $x+1, y, z$; (ii) $x, y, z+1$; (iii) $x, y, z-1$; (iv) $x-1, y, z$.

Fig. 1




Fig. 2

