

## 2-Amino-3-carboxypyridinium nitrate

Fadila Berrah,<sup>a</sup> Sofiane Bouacida<sup>b,\*</sup> and Thierry Roisnel<sup>c</sup>

<sup>a</sup>Laboratoire de Chimie Appliquée et Technologie des Matériaux LCATM, Université Larbi Ben M'hidi, 04000 Oum El Bouaghi, Algeria, <sup>b</sup>Unité de Recherche de Chimie de l'Environnement et Moléculaire Structurale, CHEMS, Faculté des Sciences Exactes, Université Mentouri Constantine 25000, Algeria., and <sup>c</sup>Centre de Difractométrie X, UMR 6226 CNRS Unité Sciences Chimiques de Rennes, Université de Rennes I, 263 Avenue du Général Leclerc, 35042 Rennes, France  
Correspondence e-mail: bouacida\_sofiane@yahoo.fr


Received 7 July 2011; accepted 12 July 2011

Key indicators: single-crystal X-ray study;  $T = 150$  K; mean  $\sigma(\text{C}-\text{C}) = 0.006$  Å;  $R$  factor = 0.059;  $wR$  factor = 0.158; data-to-parameter ratio = 11.8.

In the crystal structure of the title compound,  $\text{C}_6\text{H}_7\text{N}_2\text{O}_2^+\cdot\text{NO}_3^-$ , the cations are linked *via*  $\text{C}-\text{H}\cdots\text{O}$  hydrogen bonds, forming infinite chains running along the  $b$  axis. These chains are further linked through  $\text{N}-\text{H}\cdots\text{O}$ ,  $\text{O}-\text{H}\cdots\text{O}$  and  $\text{C}-\text{H}\cdots\text{O}$  hydrogen bonds to the nitrate anions, forming well-separated infinite planar layers parallel to (001).

## Related literature

For hybrid compounds based on nicotinic acid, see: Athimoolam et al. (2005); Athimoolam & Rajaram (2005*a,b*); Chen (2009); Slouf (2001); Ye et al. (2010). For hybrid compounds based on amino-nicotinic acid derivatives, see: Akriche & Rzaigui (2007); Berrah et al. (2011*a*); Giantsidis & Turnbull (2000). For related nitrate compounds, see: Berrah et al. (2011*b*); Jebas et al. (2006).



## Experimental

## Crystal data

$\text{C}_6\text{H}_7\text{N}_2\text{O}_2^+\cdot\text{NO}_3^-$   
 $M_r = 201.15$   
Tetragonal,  $I4_1cd$   
 $a = 16.122$  (2) Å  
 $c = 12.446$  (3) Å  
 $V = 3235.0$  (11) Å<sup>3</sup>

$Z = 16$   
Mo  $K\alpha$  radiation  
 $\mu = 0.15$  mm<sup>-1</sup>  
 $T = 150$  K  
 $0.39 \times 0.07 \times 0.05$  mm

\* Current address: Département Sciences de la Matière, Faculté des Sciences Exactes et Sciences de la Nature et de la Vie, Université Larbi Ben M'hidi, 04000 Oum El Bouaghi, Algeria.

## Data collection

Bruker APEXII diffractometer  
Absorption correction: multi-scan  
(*SADABS*; Sheldrick, 2002)  
 $T_{\min} = 0.476$ ,  $T_{\max} = 0.993$

5438 measured reflections  
1509 independent reflections  
921 reflections with  $I > 2\sigma(I)$   
 $R_{\text{int}} = 0.112$

## Refinement

$R[F^2 > 2\sigma(F^2)] = 0.059$   
 $wR(F^2) = 0.158$   
 $S = 0.97$   
1509 reflections  
128 parameters

1 restraint  
H-atom parameters constrained  
 $\Delta\rho_{\max} = 0.41$  e Å<sup>-3</sup>  
 $\Delta\rho_{\min} = -0.28$  e Å<sup>-3</sup>

Table 1

Hydrogen-bond geometry (Å, °).

$D-H\cdots A$	$D-H$	$H\cdots A$	$D\cdots A$	$D-H\cdots A$
$\text{N2}-\text{H2}\cdots\text{O3}^{\text{i}}$	0.86	1.97	2.803 (4)	162
$\text{N3}-\text{H3A}\cdots\text{O2}^{\text{j}}$	0.86	2.18	3.017 (4)	165
$\text{N3}-\text{H3A}\cdots\text{O2}^{\text{ii}}$	0.86	2.43	2.967 (4)	121
$\text{N3}-\text{H3B}\cdots\text{O4}$	0.86	2.10	2.716 (5)	128
$\text{O5}-\text{H51}\cdots\text{O3}^{\text{iii}}$	0.82	1.85	2.670 (4)	180
$\text{C4}-\text{H4}\cdots\text{O1}^{\text{iv}}$	0.93	2.42	3.197 (6)	141
$\text{C5}-\text{H5}\cdots\text{O4}^{\text{v}}$	0.93	2.30	3.216 (6)	167

Symmetry codes: (i)  $y + \frac{1}{2}, -x, z - \frac{1}{4}$ ; (ii)  $-y + \frac{1}{2}, x, z - \frac{1}{4}$ ; (iii)  $-y, -x + \frac{1}{2}, z - \frac{1}{4}$ ; (iv)  $y, x - \frac{1}{2}, z - \frac{1}{4}$ ; (v)  $-x + \frac{1}{2}, y - \frac{1}{2}, z$ .

Data collection: *APEX2* (Bruker, 2006); cell refinement: *SAINT* (Bruker, 2006); data reduction: *SAINT*; program(s) used to solve structure: *SIR2002* (Burla et al., 2005); program(s) used to refine structure: *SHELXL97* (Sheldrick, 2008); molecular graphics: *ORTEP-3 for Windows* (Farrugia, 1997) and *DIAMOND* (Brandenburg & Berndt, 2001); software used to prepare material for publication: *WinGX* (Farrugia, 1999).

We are grateful to the LCATM Laboratory, Université Larbi Ben M'hidi, Oum El Bouaghi, Algeria, for financial support.

Supplementary data and figures for this paper are available from the IUCr electronic archives (Reference: LX2193).

## References

- Akriche, S. & Rzaigui, M. (2007). *Acta Cryst.* **E63**, o3460.  
Athimoolam, S., Anitha, K. & Rajaram, R. K. (2005). *Acta Cryst.* **E61**, o2553–o2555.  
Athimoolam, S. & Rajaram, R. K. (2005*a*). *Acta Cryst.* **E61**, o2674–o2676.  
Athimoolam, S. & Rajaram, R. K. (2005*b*). *Acta Cryst.* **E61**, o2764–o2767.  
Berrah, F., Ouakkaf, A., Bouacida, S. & Roisnel, T. (2011*a*). *Acta Cryst.* **E67**, o953–o954.  
Berrah, F., Ouakkaf, A., Bouacida, S. & Roisnel, T. (2011*b*). *Acta Cryst.* **E67**, o525–o526.  
Brandenburg, K. & Berndt, M. (2001). *DIAMOND*. Crystal Impact, Bonn, Germany.  
Bruker (2006). *APEX2* and *SAINT*. Bruker AXS Inc., Madison, Wisconsin, USA.  
Burla, M. C., Caliandro, R., Camalli, M., Carrozzini, B., Casciaro, G. L., De Caro, L., Giacovazzo, C., Polidori, G. & Spagna, R. (2005). *J. Appl. Cryst.* **38**, 381–388.  
Chen, L.-Z. (2009). *Acta Cryst.* **E65**, o2350.  
Farrugia, L. J. (1997). *J. Appl. Cryst.* **30**, 565.  
Farrugia, L. J. (1999). *J. Appl. Cryst.* **32**, 837–838.

Giantsidis, J. & Turnbull, M. M. (2000). *Acta Cryst.* **C56**, 334–335.

Jebas, S. R., Balasubramanian, T. & Light, M. E. (2006). *Acta Cryst.* **E62**, o3481–o3482.

Sheldrick, G. M. (2002). *SADABS*. Bruker AXS Inc., Madison, Wisconsin, USA

Sheldrick, G. M. (2008). *Acta Cryst.* **A64**, 112–122.

Slouf, M. (2001). *Acta Cryst.* **E57**, o61–o62

Ye, H.-Y., Chen, L.-Z. & Xiong, R.-G. (2010). *Acta Cryst.* **B66**, 387–395.

**supplementary materials**

*Acta Cryst.* (2011). E67, o2057-o2058 [ doi:10.1107/S1600536811027978 ]

## 2-Amino-3-carboxypyridinium nitrate

F. Berrah, S. Bouacida and T. Roisnel

### Comment

Crystal structures of hybrid compounds based on nicotinic acid or its amino derivatives and inorganic acids have been reported (Athimoolam *et al.*, 2005; Athimoolam & Rajaram, 2005*a, b*; Chen, 2009; Giantsidis & Turnbull, 2000; Jebas *et al.*, 2006; Slouf, 2001; Ye *et al.*, 2010) showing interesting structural diversity governed mainly by hydrogen bonds. Anion substitution seems to have an important influence on hydrogen bond patterns. In attempt to elucidate this influence and as part of our search for new hybrid compounds based on protonated N-heterocycle, we report in this paper the new structure of 2-aminonicotinium nitrate; its homologues obtained with phosphate and sulfate anions have been described previously (Akriche & Rzaigui, 2007; Berrah *et al.*, 2011*a*).

The asymmetric unit of the title compound (Fig. 1) contains one cation and one anion with geometry similar to that observed in similar compounds (Akriche & Rzaigui, 2007; Berrah *et al.*, 2011*a, b*; Jebas *et al.*, 2006). However in this structure, cations do not form dimers *via* N–H $\cdots$ O hydrogen bonds as observed in the structures obtained with phosphate and sulfate anions but they are linked to each other *via* C–H $\cdots$ O hydrogen bonds to form infinite chains running along the *b* axis (Table 1 & Fig. 2.). These chains are further linked, through N–H $\cdots$ O, O–H $\cdots$ O and C–H $\cdots$ O hydrogen contacts, to nitrate anions to form well separated infinite planar layers parallel to (001) (Fig. 3). A such two-dimensional network have been already observed in compounds including nicotinium entities (Giantsidis & Turnbull, 2000; Slouf, 2001; Ye *et al.*, 2010).

### Experimental

The title compound was synthesized by reacting 3-amino-pyridine-2-carboxylic acid (0.138 mg, 1 mmol) with nitric acid (1 mmol) in a solution of equal volume of H<sub>2</sub>O and CH<sub>3</sub>OH. Slow evaporation leads to well crystallized colourless needles.

### Refinement

All the Friedel pairs were merged. All non-H atoms were refined with anisotropic atomic displacement parameters. The remaining H atoms were localized on Fourier maps but introduced in calculated positions and treated as riding on their parent atoms (C, N or O) with C–H = 0.93 Å, O–H = 0.82 Å and N–H = 0.86 Å with  $U_{\text{iso}}(\text{H}) = 1.2 U_{\text{eq}}(\text{C or N})$  and  $U_{\text{iso}}(\text{H}) = 1.5 U_{\text{eq}}(\text{O})$ .

Figures




Fig. 1. The structure of the title compound with the atomic labelling scheme. Displacement are drawn at the 50% probability level.





Fig. 2. A part of crystal packing showing cationic infinite chains linked to nitrate anions *via* [N–H···O, O–H···O and C–H···O] hydrogen contacts. Hydrogen bonds are shown as dashed lines. [Symmetry codes: (i)  $y + 1/2, -x, z - 1/4$ ; (ii)  $-y + 1/2, x, z - 1/4$ ; (iii)  $-y, -x + 1/2, z - 1/4$ ; (iv)  $y, x - 1/2, z - 1/4$ ; (v)  $-x + 1/2, y - 1/2, z$ .]



Fig. 3. Layered packing of the structure viewed down the *b* axis.

**2-Amino-3-carboxypyridinium nitrate**

*Crystal data*



$M_r = 201.15$

Tetragonal,  $I4_1cd$

Hall symbol: I 4bw -2c

$a = 16.122 (2) \text{ \AA}$

$c = 12.446 (3) \text{ \AA}$

$V = 3235.0 (11) \text{ \AA}^3$

$Z = 16$

$F(000) = 1664$

$D_x = 1.652 \text{ Mg m}^{-3}$

Mo  $K\alpha$  radiation,  $\lambda = 0.71073 \text{ \AA}$

Cell parameters from 491 reflections

$\theta = 3.3\text{--}20.3^\circ$

$\mu = 0.15 \text{ mm}^{-1}$

$T = 150 \text{ K}$

Needle, colourless

$0.39 \times 0.07 \times 0.05 \text{ mm}$

*Data collection*

Bruker APEXII  
diffractometer

graphite

CCD rotation images, thin slices scans

Absorption correction: multi-scan  
(*SADABS*; Sheldrick, 2002)

$T_{\min} = 0.476, T_{\max} = 0.993$

5438 measured reflections

1509 independent reflections

921 reflections with  $I > 2\sigma(I)$

$R_{\text{int}} = 0.112$

$\theta_{\max} = 27.5^\circ, \theta_{\min} = 3.3^\circ$

$h = -16 \rightarrow 20$

$k = -13 \rightarrow 20$

$l = -16 \rightarrow 10$

Refinement

Refinement on $F^2$	Primary atom site location: structure-invariant direct methods
Least-squares matrix: full	Secondary atom site location: difference Fourier map
$R[F^2 > 2\sigma(F^2)] = 0.059$	Hydrogen site location: difference Fourier map
$wR(F^2) = 0.158$	H-atom parameters constrained
$S = 0.97$	$w = 1/[\sigma^2(F_o^2) + (0.076P)^2]$
1509 reflections	where $P = (F_o^2 + 2F_c^2)/3$
128 parameters	$(\Delta/\sigma)_{\max} < 0.001$
1 restraint	$\Delta\rho_{\max} = 0.41 \text{ e } \text{\AA}^{-3}$
	$\Delta\rho_{\min} = -0.28 \text{ e } \text{\AA}^{-3}$

Special details

**Geometry.** All e.s.d.'s (except the e.s.d. in the dihedral angle between two l.s. planes) are estimated using the full covariance matrix. The cell e.s.d.'s are taken into account individually in the estimation of e.s.d.'s in distances, angles and torsion angles; correlations between e.s.d.'s in cell parameters are only used when they are defined by crystal symmetry. An approximate (isotropic) treatment of cell e.s.d.'s is used for estimating e.s.d.'s involving l.s. planes.

**Refinement.** Refinement of  $F^2$  against ALL reflections. The weighted  $R$ -factor  $wR$  and goodness of fit  $S$  are based on  $F^2$ , conventional  $R$ -factors  $R$  are based on  $F$ , with  $F$  set to zero for negative  $F^2$ . The threshold expression of  $F^2 > \sigma(F^2)$  is used only for calculating  $R$ -factors(gt) *etc.* and is not relevant to the choice of reflections for refinement.  $R$ -factors based on  $F^2$  are statistically about twice as large as those based on  $F$ , and  $R$ -factors based on ALL data will be even larger.

Fractional atomic coordinates and isotropic or equivalent isotropic displacement parameters ( $\text{\AA}^2$ )

	$x$	$y$	$z$	$U_{\text{iso}}^*/U_{\text{eq}}$
C1	0.1754 (3)	0.1137 (3)	0.0309 (6)	0.0288 (11)
C2	0.1875 (3)	0.0223 (2)	0.0276 (6)	0.0229 (10)
C3	0.1218 (3)	-0.0300 (3)	0.0259 (6)	0.0295 (12)
H3	0.0686	-0.0078	0.0264	0.035*
C4	0.1311 (3)	-0.1174 (3)	0.0233 (7)	0.0309 (11)
H4	0.0854	-0.1526	0.0215	0.037*
C5	0.2092 (3)	-0.1469 (3)	0.0235 (7)	0.0302 (12)
H5	0.2179	-0.2039	0.0221	0.036*
C6	0.2702 (2)	-0.0099 (3)	0.0279 (10)	0.0234 (10)
N1	0.1623 (2)	-0.0149 (2)	0.2777 (8)	0.0247 (8)
N2	0.2755 (2)	-0.0950 (2)	0.0256 (5)	0.0269 (9)
H2	0.3242	-0.1168	0.0256	0.032*
N3	0.3388 (2)	0.0327 (2)	0.0312 (5)	0.0308 (10)
H3A	0.3858	0.0075	0.032	0.037*
H3B	0.3371	0.086	0.0326	0.037*
O1	0.20156 (19)	0.05016 (19)	0.2755 (4)	0.0313 (8)
O2	0.08451 (16)	-0.01536 (18)	0.2757 (5)	0.0293 (8)
O3	0.19907 (18)	-0.08530 (18)	0.2798 (5)	0.0309 (8)
O5	0.09671 (19)	0.13575 (18)	0.0242 (4)	0.0342 (9)

## supplementary materials

---

H51	0.0932	0.1865	0.0259	0.051*
O4	0.2318 (2)	0.1628 (2)	0.0360 (4)	0.0355 (10)

### Atomic displacement parameters ( $\text{\AA}^2$ )

	$U^{11}$	$U^{22}$	$U^{33}$	$U^{12}$	$U^{13}$	$U^{23}$
C1	0.035 (3)	0.017 (2)	0.035 (3)	-0.002 (2)	-0.001 (4)	-0.004 (3)
C2	0.020 (2)	0.021 (3)	0.028 (2)	0.0022 (16)	0.001 (3)	0.002 (4)
C3	0.025 (3)	0.021 (2)	0.042 (3)	-0.0068 (19)	0.000 (4)	0.003 (3)
C4	0.042 (3)	0.016 (3)	0.035 (3)	-0.001 (2)	-0.003 (5)	0.000 (3)
C5	0.040 (3)	0.014 (2)	0.036 (3)	-0.014 (2)	-0.004 (4)	-0.001 (4)
C6	0.022 (2)	0.026 (3)	0.022 (2)	-0.0025 (18)	-0.006 (5)	0.003 (3)
N1	0.0198 (18)	0.020 (2)	0.0341 (18)	-0.0058 (16)	0.006 (4)	-0.001 (4)
N2	0.024 (2)	0.022 (2)	0.035 (2)	0.0103 (15)	-0.005 (3)	-0.002 (3)
N3	0.0137 (19)	0.029 (2)	0.050 (2)	-0.0035 (15)	0.003 (3)	0.002 (3)
O1	0.0253 (18)	0.0256 (17)	0.043 (2)	-0.0062 (15)	0.000 (3)	-0.006 (3)
O2	0.0124 (15)	0.0231 (18)	0.052 (2)	0.0030 (13)	0.004 (3)	0.004 (3)
O3	0.0130 (16)	0.0233 (17)	0.056 (2)	0.0049 (13)	-0.002 (3)	-0.007 (3)
O5	0.0233 (17)	0.0205 (19)	0.059 (2)	0.0042 (14)	0.004 (3)	0.000 (3)
O4	0.0253 (19)	0.0212 (18)	0.060 (3)	0.0027 (15)	-0.003 (3)	-0.005 (3)

### Geometric parameters ( $\text{\AA}$ , $^\circ$ )

C1—O4	1.207 (6)	C5—H5	0.93
C1—O5	1.319 (6)	C6—N3	1.303 (5)
C1—C2	1.487 (6)	C6—N2	1.375 (6)
C2—C3	1.354 (6)	N1—O1	1.225 (4)
C2—C6	1.431 (5)	N1—O2	1.255 (4)
C3—C4	1.417 (6)	N1—O3	1.281 (4)
C3—H3	0.93	N2—H2	0.86
C4—C5	1.347 (8)	N3—H3A	0.86
C4—H4	0.93	N3—H3B	0.86
C5—N2	1.359 (6)	O5—H51	0.82
O4—C1—O5	123.4 (4)	N2—C5—H5	119.4
O4—C1—C2	123.5 (4)	N3—C6—N2	118.2 (4)
O5—C1—C2	113.0 (4)	N3—C6—C2	126.9 (5)
C3—C2—C6	120.2 (4)	N2—C6—C2	114.8 (4)
C3—C2—C1	121.0 (4)	O1—N1—O2	121.4 (4)
C6—C2—C1	118.8 (4)	O1—N1—O3	121.4 (3)
C2—C3—C4	122.5 (4)	O2—N1—O3	117.2 (3)
C2—C3—H3	118.8	C5—N2—C6	124.5 (4)
C4—C3—H3	118.8	C5—N2—H2	117.8
C5—C4—C3	116.8 (4)	C6—N2—H2	117.8
C5—C4—H4	121.6	C6—N3—H3A	120
C3—C4—H4	121.6	C6—N3—H3B	120
C4—C5—N2	121.2 (4)	H3A—N3—H3B	120
C4—C5—H5	119.4	C1—O5—H51	109.5
O4—C1—C2—C3	177.5 (7)	C3—C2—C6—N3	-178.8 (9)

O5—C1—C2—C3	-4.4 (11)	C1—C2—C6—N3	0.4 (17)
O4—C1—C2—C6	-1.6 (13)	C3—C2—C6—N2	0.3 (14)
O5—C1—C2—C6	176.4 (9)	C1—C2—C6—N2	179.5 (7)
C6—C2—C3—C4	-0.6 (12)	C4—C5—N2—C6	0.0 (14)
C1—C2—C3—C4	-179.7 (7)	N3—C6—N2—C5	179.1 (8)
C2—C3—C4—C5	0.5 (11)	C2—C6—N2—C5	-0.1 (15)
C3—C4—C5—N2	-0.3 (12)		

*Hydrogen-bond geometry (Å, °)*

<i>D</i> —H... <i>A</i>	<i>D</i> —H	H... <i>A</i>	<i>D</i> ... <i>A</i>	<i>D</i> —H... <i>A</i>
N2—H2...O3 <sup>i</sup>	0.86	1.97	2.803 (4)	162
N3—H3A...O2 <sup>i</sup>	0.86	2.18	3.017 (4)	165
N3—H3A...O2 <sup>ii</sup>	0.86	2.43	2.967 (4)	121
N3—H3B...O4	0.86	2.10	2.716 (5)	128
O5—H51...O3 <sup>iii</sup>	0.82	1.85	2.670 (4)	180
C4—H4...O1 <sup>iv</sup>	0.93	2.42	3.197 (6)	141
C5—H5...O4 <sup>v</sup>	0.93	2.30	3.216 (6)	167

Symmetry codes: (i)  $y+1/2, -x, z-1/4$ ; (ii)  $-y+1/2, x, z-1/4$ ; (iii)  $-y, -x+1/2, z-1/4$ ; (iv)  $y, x-1/2, z-1/4$ ; (v)  $-x+1/2, y-1/2, z$ .



Fig. 1




Fig. 2




Fig. 3

