

(2Z)-3-(4-Chloroanilino)-1-(5-hydroxy-3-methyl-1-phenyl-1*H*-pyrazol-4-yl)but-2-en-1-one

Abdullah M. Asiri,^{a,b}‡ Abdulrahman O. Al-Youbi,^a
Khalid A. Alamry,^a Hassan M. Faidallah,^a Seik Weng Ng^{c,a}
and Edward R. T. Tiekkink^{c,*}

^aChemistry Department, Faculty of Science, King Abdulaziz University, PO Box 80203, Jeddah, Saudi Arabia, ^bThe Center of Excellence for Advanced Materials Research, King Abdulaziz University, Jeddah, PO Box 80203, Saudi Arabia, and

^cDepartment of Chemistry, University of Malaya, 50603 Kuala Lumpur, Malaysia
Correspondence e-mail: edward.tiekkink@gmail.com


Received 21 July 2011; accepted 21 July 2011

Key indicators: single-crystal X-ray study; $T = 100\text{ K}$; mean $\sigma(\text{C}-\text{C}) = 0.002\text{ \AA}$; R factor = 0.039; wR factor = 0.103; data-to-parameter ratio = 15.8.

With the exception of the terminal benzene rings, the atoms in the title compound, $\text{C}_{20}\text{H}_{18}\text{ClN}_3\text{O}_2$, are approximately coplanar (r.m.s. deviation = 0.0495 Å). The benzene/chlorobenzene rings form dihedral angles of 3.02 (4) and 41.59 (5)°, respectively, with this plane. The hydroxy, amino and carbonyl groups all lie to the same side of the molecule, enabling the formation of intramolecular O–H···O and N–H···O hydrogen bonds that close *S*(6) rings. The configuration about the 2-butene bond is *Z*. Supramolecular chains mediated by C–H···Cl interactions and aligned along the *c* axis are found in the crystal packing. These assemble into layers that are connected by weak π – π interactions between centrosymmetrically related chlorobenzene rings [3.8156 (9) Å].

Related literature

For background to the synthesis, see: Gelin *et al.* (1983); Bendaas *et al.* (1999).



Experimental

Crystal data

$\text{C}_{20}\text{H}_{18}\text{ClN}_3\text{O}_2$

$M_r = 367.82$

‡ Additional correspondence author, e-mail: aasiri2@kau.edu.sa.

Monoclinic, $P2_1/n$
 $a = 10.7782 (3)\text{ \AA}$
 $b = 12.6349 (4)\text{ \AA}$
 $c = 12.9071 (4)\text{ \AA}$
 $\beta = 100.956 (3)^\circ$
 $V = 1725.67 (9)\text{ \AA}^3$

$Z = 4$
Mo $K\alpha$ radiation
 $\mu = 0.24\text{ mm}^{-1}$
 $T = 100\text{ K}$
 $0.30 \times 0.25 \times 0.20\text{ mm}$

Data collection

Agilent SuperNova Dual diffractometer with an Atlas detector
Absorption correction: multi-scan (*CrysAlis PRO*; Agilent, 2010)
 $T_{\min} = 0.931$, $T_{\max} = 0.953$

8785 measured reflections
3860 independent reflections
3199 reflections with $I > 2\sigma(I)$
 $R_{\text{int}} = 0.025$

Refinement

$R[F^2 > 2\sigma(F^2)] = 0.039$
 $wR(F^2) = 0.103$
 $S = 1.01$
3860 reflections
245 parameters

H atoms treated by a mixture of independent and constrained refinement
 $\Delta\rho_{\max} = 0.27\text{ e \AA}^{-3}$
 $\Delta\rho_{\min} = -0.30\text{ e \AA}^{-3}$

Table 1
Hydrogen-bond geometry (Å, °).

$D-\text{H}\cdots A$	$D-\text{H}$	$\text{H}\cdots A$	$D\cdots A$	$D-\text{H}\cdots A$
O1–H1···O2	0.96 (3)	1.64 (3)	2.5283 (16)	153 (3)
N3–H3···O2	0.91 (2)	1.93 (2)	2.6678 (18)	136.9 (18)
C4–H4···Cl1 ⁱ	0.95	2.81	3.6217 (18)	144

Symmetry code: (i) $x - \frac{1}{2}, -y + \frac{3}{2}, z - \frac{3}{2}$.

Data collection: *CrysAlis PRO* (Agilent, 2010); cell refinement: *CrysAlis PRO*; data reduction: *CrysAlis PRO*; program(s) used to solve structure: *SHELXS97* (Sheldrick, 2008); program(s) used to refine structure: *SHELXL97* (Sheldrick, 2008); molecular graphics: *ORTEP-3* (Farrugia, 1997) and *DIAMOND* (Brandenburg, 2006); software used to prepare material for publication: *publCIF* (Westrip, 2010).

The authors are thankful to the Center of Excellence for Advanced Materials Research and the Chemistry Department at King Abdulaziz University for providing the research facilities. They also thank the University of Malaya for supporting this study.

Supplementary data and figures for this paper are available from the IUCr electronic archives (Reference: HG5069).

References

- Agilent (2010). *CrysAlis PRO*. Agilent Technologies, Yarnton, England.
- Bendaas, A., Hamdi, M. & Sellier, N. (1999). *J. Heterocycl. Chem.* **36**, 1291–1294.
- Brandenburg, K. (2006). *DIAMOND*. Crystal Impact GbR, Bonn, Germany.
- Farrugia, L. J. (1997). *J. Appl. Cryst.* **30**, 565.
- Gelin, S., Chantegrel, B. & Nadi, A. I. (1983). *J. Org. Chem.* **48**, 4078–4082.
- Sheldrick, G. M. (2008). *Acta Cryst. A* **64**, 112–122.
- Westrip, S. P. (2010). *J. Appl. Cryst.* **43**, 920–925.

supplementary materials

Acta Cryst. (2011). E67, o2157 [doi:10.1107/S1600536811029473]

(2Z)-3-(4-Chloroanilino)-1-(5-hydroxy-3-methyl-1-phenyl-1*H*-pyrazol-4-yl)but-2-en-1-one

A. M. Asiri, A. O. Al-Youbi, K. A. Alamry, H. M. Faidallah, S. W. Ng and E. R. T. Tiekink

Comment

The title compound (**I**) was isolated during an investigation of reactions between pyrazoles and aniline derivatives following literature precedents (Gelin *et al.*, 1983; Bendaas *et al.*, 1999). The molecular structure of (**I**), Fig. 1, features a *Z* configuration about the C12—C13 [1.376 (2) Å] bond. The hydroxy and amino groups are *syn* to the central carbonyl group and each forms a hydrogen bond to close a *S*(6) ring (Table 1). A result of this feature of the structure is that the central residue is planar; the values of the C10—C9—C11—O2, C9—C11—C12—C13 and C11—C12—C13—N3 torsion angles are 2.5 (2), -177.93 (14) and -0.4 (2) °, respectively. Indeed, the r.m.s. deviation for the non-hydrogen atoms comprising the entire molecule excluding the terminal benzene rings is 0.0495 Å. The benzene and chlorobenzene rings form dihedral angles of 3.02 (4) and 41.59 (5) °, respectively, with the central plane.

The most prominent feature of the crystal packing is the formation of supramolecular chains mediated by C—H···Cl interactions, Table 1. The chains assemble into layers in the *bc* plane, Fig. 2. The closest interactions between layers stacking along the *a* direction are weak π–π contacts between centrosymmetrically related chlorobenzene rings [3.8156 (9) Å for symmetry operation: 2 - *x*, 1 - *y*, 2 - *z*].

Experimental

A solution of 4-acetoacetyl-5-hydroxy-3-methyl-1-*p*-sulfamylphenypyrazole (1.7 g, 0.005 mol) and 4-chloroaniline (0.63 g, 0.005 mole) in ethanol (25 ml) was refluxed for 2 h. The precipitate, obtained from the hot solution, was collected, washed with methanol, and recrystallized from ethanol-benzene as orange crystals; *M.pt* 505–507 K.

Refinement

Carbon-bound H-atoms were placed in calculated positions [C—H 0.95 to 0.98 Å, $U_{\text{iso}}(\text{H})$ 1.2 to 1.5 $U_{\text{eq}}(\text{C})$] and were included in the refinement in the riding model approximation. The hydroxy- and amino- H-atoms were located in a difference Fourier map, and subsequently refined freely.

Figures




Fig. 1. The molecular structure of (**I**) showing the atom-labelling scheme and displacement ellipsoids at the 50% probability level.

supplementary materials




Fig. 2. Assembly of supramolecular chains aligned along the c axis in (I) mediated by C—H \cdots Cl interactions shown as orange dashed lines.

(2Z)-3-(4-Chloroanilino)-1-(5-hydroxy-3-methyl-1-phenyl-1*H*-pyrazol-4-yl)but-2-en-1-one

Crystal data

C ₂₀ H ₁₈ ClN ₃ O ₂	$F(000) = 768$
$M_r = 367.82$	$D_x = 1.416 \text{ Mg m}^{-3}$
Monoclinic, $P2_1/n$	Mo $K\alpha$ radiation, $\lambda = 0.71073 \text{ \AA}$
Hall symbol: -P 2yn	Cell parameters from 4068 reflections
$a = 10.7782 (3) \text{ \AA}$	$\theta = 2.5\text{--}29.3^\circ$
$b = 12.6349 (4) \text{ \AA}$	$\mu = 0.24 \text{ mm}^{-1}$
$c = 12.9071 (4) \text{ \AA}$	$T = 100 \text{ K}$
$\beta = 100.956 (3)^\circ$	Block, orange
$V = 1725.67 (9) \text{ \AA}^3$	$0.30 \times 0.25 \times 0.20 \text{ mm}$
$Z = 4$	

Data collection

Agilent SuperNova Dual diffractometer with an Atlas detector	3860 independent reflections
Radiation source: SuperNova (Mo) X-ray Source	3199 reflections with $I > 2\sigma(I)$
Mirror	$R_{\text{int}} = 0.025$
Detector resolution: 10.4041 pixels mm^{-1}	$\theta_{\text{max}} = 27.5^\circ, \theta_{\text{min}} = 2.5^\circ$
ω scans	$h = -13 \rightarrow 11$
Absorption correction: multi-scan (<i>CrysAlis PRO</i> ; Agilent, 2010)	$k = -13 \rightarrow 16$
$T_{\text{min}} = 0.931, T_{\text{max}} = 0.953$	$l = -15 \rightarrow 16$
8785 measured reflections	

Refinement

Refinement on F^2	Primary atom site location: structure-invariant direct methods
Least-squares matrix: full	Secondary atom site location: difference Fourier map
$R[F^2 > 2\sigma(F^2)] = 0.039$	Hydrogen site location: inferred from neighbouring sites
$wR(F^2) = 0.103$	H atoms treated by a mixture of independent and constrained refinement
$S = 1.01$	$w = 1/[\sigma^2(F_o^2) + (0.0495P)^2 + 0.5858P]$ where $P = (F_o^2 + 2F_c^2)/3$
3860 reflections	$(\Delta/\sigma)_{\text{max}} < 0.001$
245 parameters	$\Delta\rho_{\text{max}} = 0.27 \text{ e \AA}^{-3}$

0 restraints

 $\Delta\rho_{\min} = -0.30 \text{ e } \text{\AA}^{-3}$ *Special details*

Geometry. All e.s.d.'s (except the e.s.d. in the dihedral angle between two l.s. planes) are estimated using the full covariance matrix. The cell e.s.d.'s are taken into account individually in the estimation of e.s.d.'s in distances, angles and torsion angles; correlations between e.s.d.'s in cell parameters are only used when they are defined by crystal symmetry. An approximate (isotropic) treatment of cell e.s.d.'s is used for estimating e.s.d.'s involving l.s. planes.

Refinement. Refinement of F^2 against ALL reflections. The weighted R -factor wR and goodness of fit S are based on F^2 , conventional R -factors R are based on F , with F set to zero for negative F^2 . The threshold expression of $F^2 > \sigma(F^2)$ is used only for calculating R -factors(gt) etc. and is not relevant to the choice of reflections for refinement. R -factors based on F^2 are statistically about twice as large as those based on F , and R -factors based on ALL data will be even larger.

Fractional atomic coordinates and isotropic or equivalent isotropic displacement parameters (\AA^2)

	x	y	z	$U_{\text{iso}}^*/U_{\text{eq}}$
Cl1	0.91036 (4)	0.59737 (3)	1.20821 (3)	0.02517 (13)
O1	0.69340 (10)	0.75724 (9)	0.38953 (9)	0.0208 (3)
O2	0.73186 (10)	0.65748 (9)	0.56249 (9)	0.0216 (3)
N1	0.59886 (11)	0.64664 (11)	0.24798 (10)	0.0184 (3)
N2	0.56236 (12)	0.54063 (11)	0.23365 (10)	0.0201 (3)
N3	0.77747 (12)	0.55053 (12)	0.74398 (11)	0.0200 (3)
C1	0.57853 (13)	0.71508 (14)	0.15919 (12)	0.0195 (3)
C2	0.60600 (15)	0.82241 (14)	0.17012 (13)	0.0238 (4)
H2	0.6377	0.8518	0.2377	0.029*
C3	0.58657 (16)	0.88614 (15)	0.08098 (14)	0.0290 (4)
H3A	0.6056	0.9595	0.0878	0.035*
C4	0.53986 (16)	0.84422 (16)	-0.01787 (14)	0.0291 (4)
H4	0.5276	0.8885	-0.0785	0.035*
C5	0.51121 (15)	0.73772 (16)	-0.02783 (13)	0.0282 (4)
H5	0.4780	0.7090	-0.0954	0.034*
C6	0.53058 (14)	0.67263 (15)	0.06000 (13)	0.0236 (4)
H6	0.5113	0.5993	0.0527	0.028*
C7	0.56011 (16)	0.38241 (13)	0.33884 (14)	0.0241 (4)
H7A	0.5145	0.3555	0.2710	0.036*
H7B	0.5070	0.3749	0.3922	0.036*
H7C	0.6383	0.3420	0.3606	0.036*
C8	0.59131 (13)	0.49623 (13)	0.32786 (12)	0.0193 (3)
C9	0.64770 (14)	0.57033 (13)	0.40580 (12)	0.0185 (3)
C10	0.64989 (13)	0.66471 (13)	0.35046 (12)	0.0175 (3)
C11	0.69441 (14)	0.56817 (13)	0.51866 (12)	0.0189 (3)
C12	0.69884 (14)	0.47308 (13)	0.57726 (13)	0.0210 (3)
H12	0.6726	0.4100	0.5393	0.025*
C13	0.73809 (14)	0.46454 (13)	0.68479 (13)	0.0196 (3)
C14	0.74065 (15)	0.35747 (13)	0.73563 (13)	0.0223 (4)
H14A	0.7539	0.3030	0.6848	0.033*
H14B	0.6601	0.3448	0.7581	0.033*

supplementary materials

H14C	0.8097	0.3548	0.7971	0.033*
C15	0.80969 (14)	0.55777 (13)	0.85538 (12)	0.0187 (3)
C16	0.73868 (14)	0.50806 (14)	0.92039 (13)	0.0223 (4)
H16	0.6675	0.4665	0.8902	0.027*
C17	0.77113 (14)	0.51871 (14)	1.02897 (13)	0.0220 (4)
H17	0.7244	0.4827	1.0736	0.026*
C18	0.87226 (14)	0.58229 (13)	1.07155 (12)	0.0192 (3)
C19	0.94267 (14)	0.63411 (13)	1.00821 (13)	0.0199 (3)
H19	1.0115	0.6781	1.0386	0.024*
C20	0.91164 (14)	0.62108 (13)	0.89951 (13)	0.0192 (3)
H20	0.9601	0.6555	0.8552	0.023*
H1	0.721 (3)	0.739 (2)	0.462 (2)	0.077 (9)*
H3	0.7859 (18)	0.6100 (17)	0.7060 (17)	0.035 (6)*

Atomic displacement parameters (\AA^2)

	U^{11}	U^{22}	U^{33}	U^{12}	U^{13}	U^{23}
Cl1	0.0307 (2)	0.0284 (2)	0.0149 (2)	0.00130 (16)	0.00068 (15)	-0.00084 (17)
O1	0.0259 (6)	0.0178 (6)	0.0170 (6)	-0.0017 (4)	0.0000 (4)	-0.0009 (5)
O2	0.0283 (6)	0.0188 (6)	0.0167 (6)	-0.0025 (5)	0.0019 (4)	-0.0017 (5)
N1	0.0190 (6)	0.0191 (7)	0.0162 (7)	-0.0017 (5)	0.0013 (5)	-0.0003 (6)
N2	0.0225 (7)	0.0184 (7)	0.0186 (7)	-0.0019 (5)	0.0019 (5)	-0.0015 (6)
N3	0.0245 (7)	0.0193 (7)	0.0150 (7)	-0.0012 (5)	0.0009 (5)	0.0018 (6)
C1	0.0151 (7)	0.0269 (9)	0.0164 (8)	0.0016 (6)	0.0031 (6)	0.0032 (7)
C2	0.0229 (8)	0.0279 (10)	0.0200 (8)	-0.0024 (7)	0.0029 (6)	0.0012 (7)
C3	0.0313 (9)	0.0291 (10)	0.0264 (9)	-0.0025 (7)	0.0051 (7)	0.0066 (8)
C4	0.0290 (9)	0.0385 (11)	0.0195 (9)	0.0003 (8)	0.0039 (7)	0.0096 (8)
C5	0.0255 (8)	0.0433 (12)	0.0152 (8)	0.0010 (8)	0.0021 (6)	0.0013 (8)
C6	0.0226 (8)	0.0293 (10)	0.0187 (8)	-0.0006 (7)	0.0037 (6)	-0.0018 (7)
C7	0.0285 (8)	0.0201 (9)	0.0226 (9)	-0.0031 (7)	0.0019 (7)	-0.0038 (7)
C8	0.0187 (7)	0.0207 (9)	0.0185 (8)	0.0010 (6)	0.0033 (6)	-0.0015 (7)
C9	0.0195 (7)	0.0190 (8)	0.0169 (8)	0.0004 (6)	0.0030 (6)	-0.0008 (7)
C10	0.0152 (7)	0.0204 (8)	0.0167 (8)	0.0003 (6)	0.0023 (6)	-0.0022 (7)
C11	0.0187 (7)	0.0215 (8)	0.0166 (8)	0.0008 (6)	0.0043 (6)	-0.0016 (7)
C12	0.0236 (8)	0.0200 (9)	0.0190 (8)	0.0001 (6)	0.0032 (6)	-0.0017 (7)
C13	0.0173 (7)	0.0202 (8)	0.0214 (8)	0.0005 (6)	0.0037 (6)	-0.0007 (7)
C14	0.0244 (8)	0.0200 (9)	0.0210 (9)	-0.0009 (6)	0.0009 (6)	0.0010 (7)
C15	0.0206 (7)	0.0187 (8)	0.0156 (8)	0.0036 (6)	0.0006 (6)	-0.0001 (6)
C16	0.0189 (7)	0.0261 (9)	0.0209 (8)	-0.0038 (6)	0.0012 (6)	-0.0004 (7)
C17	0.0229 (8)	0.0239 (9)	0.0198 (8)	0.0002 (6)	0.0052 (6)	0.0016 (7)
C18	0.0221 (7)	0.0195 (8)	0.0145 (8)	0.0052 (6)	-0.0002 (6)	0.0007 (6)
C19	0.0212 (8)	0.0164 (8)	0.0204 (8)	0.0008 (6)	0.0000 (6)	-0.0015 (7)
C20	0.0225 (8)	0.0155 (8)	0.0196 (8)	0.0018 (6)	0.0041 (6)	0.0016 (6)

Geometric parameters (\AA , $^\circ$)

Cl1—C18	1.7438 (16)	C7—H7A	0.9800
O1—C10	1.3233 (19)	C7—H7B	0.9800
O1—H1	0.96 (3)	C7—H7C	0.9800

O2—C11	1.2921 (19)	C8—C9	1.423 (2)
N1—C10	1.3519 (19)	C9—C10	1.392 (2)
N1—N2	1.3984 (19)	C9—C11	1.448 (2)
N1—C1	1.419 (2)	C11—C12	1.416 (2)
N2—C8	1.321 (2)	C12—C13	1.376 (2)
N3—C13	1.349 (2)	C12—H12	0.9500
N3—C15	1.417 (2)	C13—C14	1.502 (2)
N3—H3	0.91 (2)	C14—H14A	0.9800
C1—C2	1.389 (2)	C14—H14B	0.9800
C1—C6	1.394 (2)	C14—H14C	0.9800
C2—C3	1.387 (2)	C15—C20	1.391 (2)
C2—H2	0.9500	C15—C16	1.389 (2)
C3—C4	1.385 (3)	C16—C17	1.385 (2)
C3—H3A	0.9500	C16—H16	0.9500
C4—C5	1.381 (3)	C17—C18	1.381 (2)
C4—H4	0.9500	C17—H17	0.9500
C5—C6	1.384 (2)	C18—C19	1.382 (2)
C5—H5	0.9500	C19—C20	1.389 (2)
C6—H6	0.9500	C19—H19	0.9500
C7—C8	1.490 (2)	C20—H20	0.9500
C10—O1—H1	100.3 (17)	C8—C9—C11	135.94 (15)
C10—N1—N2	110.06 (13)	O1—C10—N1	124.81 (14)
C10—N1—C1	131.22 (14)	O1—C10—C9	126.90 (14)
N2—N1—C1	118.72 (13)	N1—C10—C9	108.29 (14)
C8—N2—N1	105.88 (12)	O2—C11—C12	122.09 (14)
C13—N3—C15	128.07 (15)	O2—C11—C9	116.35 (14)
C13—N3—H3	114.3 (13)	C12—C11—C9	121.56 (15)
C15—N3—H3	117.6 (13)	C13—C12—C11	125.31 (16)
C2—C1—C6	120.24 (15)	C13—C12—H12	117.3
C2—C1—N1	121.03 (15)	C11—C12—H12	117.3
C6—C1—N1	118.73 (15)	N3—C13—C12	120.74 (15)
C1—C2—C3	119.12 (16)	N3—C13—C14	120.02 (14)
C1—C2—H2	120.4	C12—C13—C14	119.21 (15)
C3—C2—H2	120.4	C13—C14—H14A	109.5
C4—C3—C2	120.88 (18)	C13—C14—H14B	109.5
C4—C3—H3A	119.6	H14A—C14—H14B	109.5
C2—C3—H3A	119.6	C13—C14—H14C	109.5
C5—C4—C3	119.61 (17)	H14A—C14—H14C	109.5
C5—C4—H4	120.2	H14B—C14—H14C	109.5
C3—C4—H4	120.2	C20—C15—C16	119.74 (15)
C4—C5—C6	120.43 (17)	C20—C15—N3	118.36 (14)
C4—C5—H5	119.8	C16—C15—N3	121.82 (14)
C6—C5—H5	119.8	C17—C16—C15	120.35 (15)
C5—C6—C1	119.71 (17)	C17—C16—H16	119.8
C5—C6—H6	120.1	C15—C16—H16	119.8
C1—C6—H6	120.1	C18—C17—C16	119.18 (15)
C8—C7—H7A	109.5	C18—C17—H17	120.4
C8—C7—H7B	109.5	C16—C17—H17	120.4
H7A—C7—H7B	109.5	C19—C18—C17	121.39 (15)

supplementary materials


C8—C7—H7C	109.5	C19—C18—Cl1	119.77 (12)
H7A—C7—H7C	109.5	C17—C18—Cl1	118.84 (13)
H7B—C7—H7C	109.5	C18—C19—C20	119.19 (15)
N2—C8—C9	111.44 (14)	C18—C19—H19	120.4
N2—C8—C7	118.63 (14)	C20—C19—H19	120.4
C9—C8—C7	129.91 (15)	C19—C20—C15	120.12 (15)
C10—C9—C8	104.32 (14)	C19—C20—H20	119.9
C10—C9—C11	119.70 (15)	C15—C20—H20	119.9
C10—N1—N2—C8	0.30 (16)	C11—C9—C10—O1	1.3 (2)
C1—N1—N2—C8	-179.71 (12)	C8—C9—C10—N1	-0.15 (16)
C10—N1—C1—C2	-3.9 (2)	C11—C9—C10—N1	-178.50 (13)
N2—N1—C1—C2	176.08 (13)	C10—C9—C11—O2	2.5 (2)
C10—N1—C1—C6	175.87 (14)	C8—C9—C11—O2	-175.18 (16)
N2—N1—C1—C6	-4.1 (2)	C10—C9—C11—C12	-177.32 (14)
C6—C1—C2—C3	-0.8 (2)	C8—C9—C11—C12	5.0 (3)
N1—C1—C2—C3	178.97 (14)	O2—C11—C12—C13	2.2 (2)
C1—C2—C3—C4	0.3 (2)	C9—C11—C12—C13	-177.93 (14)
C2—C3—C4—C5	0.5 (3)	C15—N3—C13—C12	173.70 (14)
C3—C4—C5—C6	-0.9 (3)	C15—N3—C13—C14	-8.2 (2)
C4—C5—C6—C1	0.4 (2)	C11—C12—C13—N3	-0.4 (2)
C2—C1—C6—C5	0.4 (2)	C11—C12—C13—C14	-178.47 (14)
N1—C1—C6—C5	-179.35 (14)	C13—N3—C15—C20	141.37 (16)
N1—N2—C8—C9	-0.40 (16)	C13—N3—C15—C16	-42.0 (2)
N1—N2—C8—C7	178.35 (13)	C20—C15—C16—C17	-1.9 (2)
N2—C8—C9—C10	0.35 (17)	N3—C15—C16—C17	-178.53 (15)
C7—C8—C9—C10	-178.22 (15)	C15—C16—C17—C18	2.2 (2)
N2—C8—C9—C11	178.29 (16)	C16—C17—C18—C19	-1.0 (2)
C7—C8—C9—C11	-0.3 (3)	C16—C17—C18—Cl1	178.69 (12)
N2—N1—C10—O1	-179.86 (13)	C17—C18—C19—C20	-0.6 (2)
C1—N1—C10—O1	0.1 (2)	Cl1—C18—C19—C20	179.74 (12)
N2—N1—C10—C9	-0.08 (16)	C18—C19—C20—C15	0.9 (2)
C1—N1—C10—C9	179.92 (14)	C16—C15—C20—C19	0.3 (2)
C8—C9—C10—O1	179.62 (14)	N3—C15—C20—C19	177.05 (14)

Hydrogen-bond geometry (\AA , $^\circ$)

$D\cdots H$	D	$H\cdots A$	$D\cdots A$	$D\cdots H$
O1—H1 \cdots O2	0.96 (3)	1.64 (3)	2.5283 (16)	153 (3)
N3—H3 \cdots O2	0.91 (2)	1.93 (2)	2.6678 (18)	136.9 (18)
C4—H4 \cdots Cl1 ⁱ	0.95	2.81	3.6217 (18)	144

Symmetry codes: (i) $x-1/2, -y+3/2, z-3/2$.

Fig. 1



supplementary materials

Fig. 2

