

Acta Crystallographica Section E

Structure Reports

Online

ISSN 1600-5368

catena-Poly[[tetrakis(μ -pentafluorobenzoato- κ^2 O:O')dimolybdenum(II)]- μ -4,4'-bipyridine- κ^2 N:N']

Li-Juan Han

Department of Chemistry, Tongji University, Shanghai 200092, People's Republic of China

Correspondence e-mail: 08hanlij@tongji.edu.cn


Received 14 July 2011; accepted 5 August 2011

Key indicators: single-crystal X-ray study; $T = 293$ K; mean $\sigma(\text{C}-\text{C}) = 0.004$ Å; R factor = 0.024; wR factor = 0.061; data-to-parameter ratio = 10.4.

In the title compound, $[\text{Mo}_2(\text{C}_7\text{F}_5\text{O}_2)_4(\text{C}_{10}\text{H}_8\text{N}_2)]_n$, the molecule forms a paddle-wheel-type structure. Each Mo_2^{4+} unit is equatorially coordinated by four pentafluorobenzoate groups, while the axial positions are occupied by two 4,4'-bipyridine molecules. The Mo—Mo bond length of 2.1227 (4) Å is representative of a dimolybdenum quadruple bond. An infinite linear chain parallel to [110] is formed by the Mo_2^{4+} unit coordinating axially to the two N atoms of the 4,4'-bipyridine ligand [Mo—N = 2.594 (2) Å]. The crystal packing shows molecules linked together into a three-dimensional network *via* Mo—N coordination interactions and weak π – π stacking interactions between perfluorophenyl rings [centroid–centroid distance = 3.7280 (3) Å and centroid-to-plane distance = 3.6103 (12) Å between two pentafluorophenyl rings].

Related literature

For background to coordination polymers, see: Batten (2002); Kumar *et al.* (2004). For torsion angles about the pentafluorobenzoate anion, see: Reddy *et al.* (2004); Bach *et al.* (2001); For Mo—Mo quadruple bond lengths, see: Cotton *et al.* (2005).



Experimental

Crystal data

$[\text{Mo}_2(\text{C}_7\text{F}_5\text{O}_2)_4(\text{C}_{10}\text{H}_8\text{N}_2)]$
 $M_r = 1192.34$
 Triclinic, $P\bar{1}$
 $a = 8.8858$ (8) Å
 $b = 9.9311$ (9) Å
 $c = 11.1978$ (10) Å
 $\alpha = 101.158$ (1)°
 $\beta = 94.697$ (1)°

$\gamma = 99.092$ (1)°
 $V = 950.83$ (15) Å³
 $Z = 1$
 Mo $K\alpha$ radiation
 $\mu = 0.82$ mm⁻¹
 $T = 293$ K
 $0.20 \times 0.18 \times 0.15$ mm

Data collection

Bruker APEXII CCD
 diffractometer
 Absorption correction: multi-scan
 (SADABS; Sheldrick, 2004)
 $T_{\min} = 0.849$, $T_{\max} = 0.884$

4937 measured reflections
 3294 independent reflections
 3072 reflections with $I > 2\sigma(I)$
 $R_{\text{int}} = 0.013$

Refinement

$R[F^2 > 2\sigma(F^2)] = 0.024$
 $wR(F^2) = 0.061$
 $S = 1.01$
 3294 reflections

316 parameters
 H-atom parameters constrained
 $\Delta\rho_{\text{max}} = 0.32$ e Å⁻³
 $\Delta\rho_{\text{min}} = -0.47$ e Å⁻³

Table 1

Selected bond lengths (Å).

Mo1—O1	2.1124 (17)	Mo1—O5	2.1427 (16)
Mo1—O6	2.1155 (16)		

Table 2

Hydrogen-bond geometry (Å, °).

$D-H\cdots A$	$D-H$	$H\cdots A$	$D\cdots A$	$D-H\cdots A$
C55—H55A \cdots F35	0.93	2.55	3.152 (2)	122
C51—H51A \cdots F33 ⁱⁱ	0.93	2.78	2.987 (3)	94

Symmetry code: (ii) $-x + 1, -y + 1, -z + 2$.

Data collection: APEX2 (Bruker, 2004); cell refinement: SAINT-Plus (Bruker, 2001); data reduction: SAINT-Plus; program(s) used to solve structure: SHELXS97 (Sheldrick, 2008); program(s) used to refine structure: SHELXL97 (Sheldrick, 2008); molecular graphics: XP in SHELXTL (Sheldrick, 2008); software used to prepare material for publication: SHELXL97.

This work was supported by the National Natural Scientific Foundation of China (No. 20741004/B010303).

Supplementary data and figures for this paper are available from the IUCr electronic archives (Reference: JJ2094).

References

Bach, A., Lentz, D. & Luger, P. (2001). *J. Phys. Chem. A*, **105**, 7405–7412.

Batten, S. R. (2002). *Curr. Opin. Solid State Mater. Sci.* **5**, 107–114.
Bruker (2001). *SAINT-Plus*. Bruker AXS Inc., Madison, Wisconsin, USA.
Bruker (2004). *APEX2*. Bruker AXS Inc., Madison, Wisconsin, USA.
Cotton, F. A., Murillo, C. A. & Walton, R. A. (2005). *Multiple Bonds Between Metal Atoms*, 3rd ed. New York: Springer Science and Business Media Inc.
Kumar, T., Uemura, K., Chang, H.-C., Matsuda, R. & Kitagawa, S. (2004). *Angew. Chem. Int. Ed.* **43**, 3269–3272.
Reddy, L. S., Nangia, A. & Lynch, V. M. (2004). *Cryst. Growth Des.* **4**, 89–94.
Sheldrick, G. M. (2004). *SADABS*. University of Göttingen, Germany.
Sheldrick, G. M. (2008). *Acta Cryst.* **A64**, 112–122.

supplementary materials

Acta Cryst. (2011). E67, m1235-m1236 [doi:10.1107/S1600536811031734]

***catena*-Poly[[tetrakis(μ -pentafluorobenzoato- $\kappa^2 O:O'$)dimolybdenum(II)]- μ -4,4'-bipyridine- $\kappa^2 N:N'$]**

L.-J. Han

Comment

The design and construction of coordination polymers are of great interest due to their structural topologies and potential application as functional materials (Batten, 2002; Kumar, *et al.*, 2004). Here, we report the synthesis and crystal structure of the coordination polymer [Mo₂(OOC₆F₅)₄(C₁₀H₈N₂)]_n.

In the title compound, Mo₂(C₇F₅O₂)₄(C₁₀H₈N₂)_n (**I**), the molecule forms a paddle-wheel-type structure. Each quadruply bonded Mo₂⁴⁺ unit is equatorially coordinated by four pentafluoro-benzoate (OOC₆F₅) groups and the axial positions have associated with them two 4, 4-bipyridine molecules (Fig. 1). The Mo–Mo bond length of 2.1227 (4) Å is representative for dimolybdenum quadruple bonds (Cotton *et al.*, 2005). The torsion angles between the C₆F₅ group and the connected chelating ring (Mo₂OCO) are 72.081 (3)°, 75.537 (3)°, 22.059 (3)° and 22.422 (3)°, respectively, and relate to the O···F repulsion within the pentafluoro-benzonate anion (Reddy *et al.*, 2004; Bach *et al.*, 2001). Weak π – π stacking interactions between perfluorophenyl rings also affect the 72.081 (3)° and 75.537 (3)° torsion angles.

An infinite, linear chain coordination polymer is formed by the Mo₂⁴⁺ unit coordinating axially to the two N atoms of the 4, 4-bipyridine ligand [Mo–N distance = 2.5938 (2) Å] (Fig. 2). A one-dimensional linear chain is generated by the π – π stacking between perfluorophenyl rings as viewed along the equatorial position of the Mo–Mo quadruple bonds (Fig. 3), [the center-to-center distance = 3.7280 (3) Å and center-to-plane distance = 3.6103 (0) Å between two pentafluorophenyl rings]. Crystal packing shows molecules linked together into a three-dimensional network (Fig. 4) *via* Mo–N coordination interactions and perfluorophenyl rings *via* weak π – π stacking interactions. Weak C–H···F intermolecular interactions further stabilize the crystal structure [(F···H distances = 2.7879 (15) Å].

Experimental

4, 4-bipyridine (0.312 g, 2 mmol) was dissolved in dichloromethane (30 ml), and the solution was filtered to a Schlenk tube. Mo₂(OOC₆F₅)₄ (0.207 g, 0.2 mmol) was dissolved in ethanol (10 ml), resulting in a clear yellow solution. The yellow solution was carefully layered on the top of the Schlenk tube. Solution diffusion at low temperature (in refrigerator) afforded yellow X-ray quality crystals after three days. Yield: 0.190 g (80%). Anal. Calcd. for C₃₈H₈N₂O₈F₂₀Mo₂: C, 38.28; H, 0.68; N, 2.35. Found: C, 38.13; H, 0.53; N, 2.37.

Refinement

The H atoms were positioned geometrically and refined using the riding model with C–H = 0.93 Å for aromatic H, 0.96 Å for methyl H atoms. The U_{iso} parameters for H atoms were constrained to be 1.5 U_{eq} of the carrier atom for the methyl H atoms and 1.2 U_{eq} of the carrier atom for the remaining H atoms.

Figures




Fig. 1. Molecular structure of the title compound drawn with displacement ellipsoids at the 30% probability level. All hydrogen atoms have been omitted for clarity.




Fig. 2. Part of a one-dimensional linear chain formed by a Mo–N coordination bond viewed along the *a* axis.




Fig. 3. Part of a one-dimensional linear chain formed by the π – π intermolecular stacking interactions.




Fig. 4. Molecular packing diagram of (I) with a view along the *a* axis.

catena-Poly[[tetrakis(μ -pentafluorobenzoato- κ^2 O:O')dimolybdenum(II)]- μ -4,4'-bipyridine- κ^2 N:N']

Crystal data

[Mo₂(C₇F₅O₂)₄(C₁₀H₈N₂)]

$M_r = 1192.34$

Triclinic, $P\bar{1}$

Hall symbol: -P 1

$a = 8.8858(8) \text{ \AA}$

$Z = 1$

$F(000) = 578$

$D_x = 2.082 \text{ Mg m}^{-3}$

Mo $K\alpha$ radiation, $\lambda = 0.71073 \text{ \AA}$

Cell parameters from 4122 reflections

$b = 9.9311 (9) \text{ \AA}$
 $c = 11.1978 (10) \text{ \AA}$
 $\alpha = 101.158 (1)^\circ$
 $\beta = 94.697 (1)^\circ$
 $\gamma = 99.092 (1)^\circ$
 $V = 950.83 (15) \text{ \AA}^3$

$\theta = 2.5\text{--}27.6^\circ$
 $\mu = 0.82 \text{ mm}^{-1}$
 $T = 293 \text{ K}$
 Block, yellow
 $0.20 \times 0.18 \times 0.15 \text{ mm}$

Data collection

Bruker APEXII CCD diffractometer
 Radiation source: fine-focus sealed tube
 graphite
 ω scans
 Absorption correction: multi-scan (SADABS; Sheldrick, 2004)
 $T_{\min} = 0.849$, $T_{\max} = 0.884$
 4937 measured reflections

3294 independent reflections
 3072 reflections with $I > 2\sigma(I)$
 $R_{\text{int}} = 0.013$
 $\theta_{\max} = 25.0^\circ$, $\theta_{\min} = 2.1^\circ$
 $h = -8 \rightarrow 10$
 $k = -11 \rightarrow 11$
 $l = -13 \rightarrow 12$

Refinement

Refinement on F^2
 Least-squares matrix: full
 $R[F^2 > 2\sigma(F^2)] = 0.024$
 $wR(F^2) = 0.061$
 $S = 1.01$
 3294 reflections
 316 parameters
 0 restraints

Primary atom site location: structure-invariant direct methods
 Secondary atom site location: difference Fourier map
 Hydrogen site location: inferred from neighbouring sites
 H-atom parameters constrained
 $w = 1/[\sigma^2(F_o^2) + (0.0323P)^2 + 0.6373P]$
 where $P = (F_o^2 + 2F_c^2)/3$
 $(\Delta/\sigma)_{\max} = 0.001$
 $\Delta\rho_{\max} = 0.32 \text{ e \AA}^{-3}$
 $\Delta\rho_{\min} = -0.47 \text{ e \AA}^{-3}$

Special details

Geometry. All e.s.d.'s (except the e.s.d. in the dihedral angle between two l.s. planes) are estimated using the full covariance matrix. The cell e.s.d.'s are taken into account individually in the estimation of e.s.d.'s in distances, angles and torsion angles; correlations between e.s.d.'s in cell parameters are only used when they are defined by crystal symmetry. An approximate (isotropic) treatment of cell e.s.d.'s is used for estimating e.s.d.'s involving l.s. planes.

Refinement. Refinement of F^2 against ALL reflections. The weighted R -factor wR and goodness of fit S are based on F^2 , conventional R -factors R are based on F , with F set to zero for negative F^2 . The threshold expression of $F^2 > \sigma(F^2)$ is used only for calculating R -factors(gt) etc. and is not relevant to the choice of reflections for refinement. R -factors based on F^2 are statistically about twice as large as those based on F , and R -factors based on ALL data will be even larger.

Fractional atomic coordinates and isotropic or equivalent isotropic displacement parameters (\AA^2)

x y z $U_{\text{iso}}^*/U_{\text{eq}}$

supplementary materials

Mo1	0.94356 (2)	0.58128 (2)	0.480838 (17)	0.02757 (8)
F11	0.46291 (19)	0.34697 (17)	0.29542 (17)	0.0569 (4)
F12	0.2509 (2)	0.1493 (2)	0.1523 (2)	0.0712 (5)
F13	0.3174 (2)	-0.10294 (19)	0.05942 (18)	0.0682 (5)
F14	0.6015 (2)	-0.15799 (17)	0.11139 (18)	0.0662 (5)
F15	0.81465 (19)	0.03236 (17)	0.25713 (17)	0.0568 (4)
F31	1.0522 (2)	0.7439 (2)	0.89717 (17)	0.0701 (5)
F32	0.9848 (3)	0.8073 (2)	1.13037 (18)	0.0825 (6)
F33	0.7539 (2)	0.6435 (2)	1.20533 (14)	0.0708 (6)
F34	0.5986 (3)	0.4110 (2)	1.05142 (19)	0.0823 (6)
F35	0.6643 (2)	0.3488 (2)	0.81764 (17)	0.0770 (6)
O1	0.75702 (18)	0.43333 (16)	0.38004 (15)	0.0332 (4)
O2	0.87528 (19)	0.26377 (17)	0.42436 (15)	0.0345 (4)
O5	0.85431 (19)	0.59500 (18)	0.65400 (15)	0.0355 (4)
O6	1.03083 (19)	0.58024 (17)	0.31053 (15)	0.0352 (4)
C1	0.7665 (3)	0.3051 (3)	0.3681 (2)	0.0330 (5)
C3	0.8955 (3)	0.5167 (3)	0.7219 (2)	0.0325 (5)
C11	0.6483 (3)	0.1984 (3)	0.2857 (2)	0.0341 (5)
C12	0.5018 (3)	0.2234 (3)	0.2528 (2)	0.0391 (6)
C13	0.3913 (3)	0.1235 (3)	0.1796 (3)	0.0456 (7)
C14	0.4231 (3)	-0.0065 (3)	0.1325 (3)	0.0467 (7)
C15	0.5677 (3)	-0.0340 (3)	0.1609 (2)	0.0437 (6)
C16	0.6762 (3)	0.0656 (3)	0.2357 (2)	0.0382 (6)
C31	0.8594 (3)	0.5460 (3)	0.8527 (2)	0.0346 (5)
C32	0.9392 (3)	0.6610 (3)	0.9340 (2)	0.0428 (6)
C33	0.9070 (4)	0.6945 (3)	1.0538 (2)	0.0497 (7)
C34	0.7909 (3)	0.6100 (3)	1.0917 (2)	0.0480 (7)
C35	0.7115 (3)	0.4929 (3)	1.0133 (3)	0.0502 (7)
C36	0.7458 (3)	0.4611 (3)	0.8941 (2)	0.0437 (6)
N1	0.2434 (3)	0.2426 (2)	0.5169 (2)	0.0434 (5)
C51	0.3883 (4)	0.2800 (3)	0.5640 (2)	0.0498 (7)
H51A	0.4226	0.3727	0.6027	0.060*
C52	0.1981 (3)	0.1097 (3)	0.4617 (3)	0.0555 (8)
H52A	0.0966	0.0810	0.4270	0.067*
C53	0.2934 (3)	0.0113 (3)	0.4527 (3)	0.0540 (8)
H53A	0.2554	-0.0805	0.4130	0.065*
C54	0.4463 (3)	0.0507 (3)	0.5034 (2)	0.0377 (6)
C55	0.4927 (3)	0.1900 (3)	0.5594 (2)	0.0467 (7)
H55A	0.5938	0.2230	0.5938	0.056*

Atomic displacement parameters (\AA^2)

	U^{11}	U^{22}	U^{33}	U^{12}	U^{13}	U^{23}
Mo1	0.03074 (12)	0.03149 (12)	0.02725 (12)	0.01909 (8)	0.00822 (8)	0.00964 (8)
F11	0.0419 (9)	0.0543 (10)	0.0761 (12)	0.0260 (8)	0.0033 (8)	0.0049 (9)
F12	0.0397 (9)	0.0805 (13)	0.0925 (14)	0.0211 (9)	-0.0114 (9)	0.0154 (11)
F13	0.0585 (11)	0.0614 (11)	0.0735 (12)	-0.0033 (9)	-0.0153 (10)	0.0079 (9)
F14	0.0698 (12)	0.0416 (9)	0.0795 (13)	0.0163 (9)	-0.0058 (10)	-0.0049 (9)

F15	0.0458 (9)	0.0479 (9)	0.0739 (12)	0.0251 (8)	-0.0035 (8)	-0.0034 (8)
F31	0.0668 (12)	0.0753 (13)	0.0585 (11)	-0.0139 (10)	0.0155 (9)	0.0085 (9)
F32	0.1018 (17)	0.0801 (14)	0.0504 (11)	0.0059 (12)	0.0050 (11)	-0.0135 (10)
F33	0.0872 (14)	0.1141 (16)	0.0306 (8)	0.0598 (12)	0.0245 (9)	0.0213 (9)
F34	0.0797 (14)	0.1097 (17)	0.0669 (12)	0.0041 (12)	0.0374 (11)	0.0393 (12)
F35	0.0870 (15)	0.0773 (13)	0.0534 (11)	-0.0205 (11)	0.0149 (10)	0.0069 (10)
O1	0.0319 (9)	0.0335 (9)	0.0390 (9)	0.0169 (7)	0.0043 (7)	0.0098 (7)
O2	0.0357 (9)	0.0349 (9)	0.0375 (9)	0.0162 (7)	0.0044 (7)	0.0108 (7)
O5	0.0391 (9)	0.0449 (10)	0.0318 (9)	0.0260 (8)	0.0126 (7)	0.0124 (7)
O6	0.0403 (10)	0.0412 (9)	0.0334 (9)	0.0225 (8)	0.0121 (7)	0.0153 (7)
C1	0.0331 (13)	0.0395 (13)	0.0328 (12)	0.0160 (11)	0.0105 (10)	0.0123 (10)
C3	0.0317 (12)	0.0401 (13)	0.0312 (12)	0.0150 (11)	0.0092 (10)	0.0116 (10)
C11	0.0344 (13)	0.0394 (13)	0.0339 (12)	0.0131 (11)	0.0097 (10)	0.0131 (10)
C12	0.0380 (14)	0.0440 (14)	0.0426 (14)	0.0178 (12)	0.0109 (11)	0.0162 (12)
C13	0.0342 (14)	0.0573 (17)	0.0504 (16)	0.0141 (13)	0.0022 (12)	0.0196 (14)
C14	0.0457 (16)	0.0500 (16)	0.0420 (15)	0.0011 (13)	-0.0020 (13)	0.0126 (13)
C15	0.0505 (16)	0.0375 (14)	0.0441 (15)	0.0117 (12)	0.0056 (13)	0.0079 (12)
C16	0.0351 (13)	0.0423 (14)	0.0419 (14)	0.0151 (11)	0.0053 (11)	0.0132 (11)
C31	0.0365 (13)	0.0452 (14)	0.0298 (12)	0.0206 (11)	0.0109 (10)	0.0128 (11)
C32	0.0408 (15)	0.0538 (16)	0.0387 (14)	0.0140 (13)	0.0112 (12)	0.0141 (12)
C33	0.0576 (18)	0.0579 (17)	0.0334 (14)	0.0234 (15)	0.0003 (13)	0.0008 (13)
C34	0.0540 (17)	0.076 (2)	0.0283 (13)	0.0404 (16)	0.0159 (12)	0.0177 (14)
C35	0.0463 (16)	0.073 (2)	0.0427 (16)	0.0201 (15)	0.0188 (13)	0.0262 (15)
C36	0.0443 (15)	0.0543 (16)	0.0357 (14)	0.0124 (13)	0.0101 (12)	0.0117 (12)
N1	0.0503 (14)	0.0533 (14)	0.0405 (12)	0.0343 (11)	0.0176 (11)	0.0187 (10)
C51	0.0609 (19)	0.0551 (17)	0.0396 (15)	0.0382 (15)	0.0052 (13)	0.0025 (13)
C52	0.0399 (16)	0.0552 (18)	0.084 (2)	0.0259 (14)	0.0165 (15)	0.0281 (17)
C53	0.0416 (16)	0.0402 (15)	0.090 (2)	0.0202 (13)	0.0163 (15)	0.0234 (15)
C54	0.0446 (15)	0.0458 (14)	0.0363 (13)	0.0264 (12)	0.0179 (11)	0.0211 (11)
C55	0.0491 (16)	0.0586 (17)	0.0360 (14)	0.0320 (14)	0.0019 (12)	0.0018 (12)

Geometric parameters (Å, °)

Mo1—O2 ⁱ	2.0955 (17)	C11—C12	1.397 (3)
Mo1—O1	2.1124 (17)	C11—C16	1.398 (3)
Mo1—O6	2.1155 (16)	C12—C13	1.366 (4)
Mo1—Mo1 ⁱ	2.1227 (4)	C13—C14	1.379 (4)
Mo1—O5	2.1427 (16)	C14—C15	1.379 (4)
Mo1—N1 ⁱⁱ	2.594 (2)	C15—C16	1.362 (4)
F11—C12	1.337 (3)	C31—C32	1.368 (4)
F12—C13	1.335 (3)	C31—C36	1.379 (4)
F13—C14	1.325 (3)	C32—C33	1.383 (4)
F14—C15	1.340 (3)	C33—C34	1.373 (4)
F15—C16	1.338 (3)	C34—C35	1.365 (4)
F31—C32	1.338 (3)	C35—C36	1.381 (4)
F32—C33	1.325 (3)	N1—C51	1.318 (4)
F33—C34	1.333 (3)	N1—C52	1.327 (4)
F34—C35	1.341 (3)	C51—C55	1.383 (4)

supplementary materials

F35—C36	1.333 (3)	C51—H51A	0.9300
O1—C1	1.272 (3)	C52—C53	1.385 (4)
O2—C1	1.274 (3)	C52—H52A	0.9300
O2—Mo1 ⁱ	2.0955 (17)	C53—C54	1.393 (4)
O5—C3	1.261 (3)	C53—H53A	0.9300
O6—C3 ⁱ	1.261 (3)	C54—C55	1.385 (4)
C1—C11	1.480 (3)	C54—C54 ⁱⁱⁱ	1.489 (5)
C3—O6 ⁱ	1.261 (3)	C55—H55A	0.9300
C3—C31	1.508 (3)		
O2 ⁱ —Mo1—O1	176.82 (6)	F15—C16—C11	120.6 (2)
O2 ⁱ —Mo1—O6	92.90 (7)	C15—C16—C11	122.5 (2)
O1—Mo1—O6	85.79 (7)	C32—C31—C36	118.0 (2)
O2 ⁱ —Mo1—Mo1 ⁱ	92.41 (4)	C32—C31—C3	119.8 (2)
O1—Mo1—Mo1 ⁱ	90.57 (4)	C36—C31—C3	122.2 (2)
O6—Mo1—Mo1 ⁱ	93.70 (4)	F31—C32—C31	119.4 (2)
O2 ⁱ —Mo1—O5	85.43 (7)	F31—C32—C33	118.7 (3)
O1—Mo1—O5	95.73 (7)	C31—C32—C33	121.9 (3)
O6—Mo1—O5	176.63 (6)	F32—C33—C34	120.4 (3)
Mo1 ⁱ —Mo1—O5	89.29 (4)	F32—C33—C32	120.8 (3)
C1—O1—Mo1	117.52 (15)	C34—C33—C32	118.9 (3)
C1—O2—Mo1 ⁱ	116.51 (15)	F33—C34—C35	119.8 (3)
C3—O5—Mo1	117.24 (14)	F33—C34—C33	119.8 (3)
C3 ⁱ —O6—Mo1	114.32 (14)	C35—C34—C33	120.5 (2)
O1—C1—O2	122.5 (2)	F34—C35—C34	119.9 (3)
O1—C1—C11	119.5 (2)	F34—C35—C36	120.3 (3)
O2—C1—C11	118.0 (2)	C34—C35—C36	119.8 (3)
O5—C3—O6 ⁱ	124.7 (2)	F35—C36—C31	119.9 (2)
O5—C3—C31	117.2 (2)	F35—C36—C35	119.0 (3)
O6 ⁱ —C3—C31	118.1 (2)	C31—C36—C35	121.0 (3)
C12—C11—C16	115.6 (2)	C51—N1—C52	116.5 (2)
C12—C11—C1	122.4 (2)	N1—C51—C55	124.1 (3)
C16—C11—C1	122.0 (2)	N1—C51—H51A	117.9
F11—C12—C13	117.1 (2)	C55—C51—H51A	117.9
F11—C12—C11	120.6 (2)	N1—C52—C53	123.8 (3)
C13—C12—C11	122.3 (2)	N1—C52—H52A	118.1
F12—C13—C12	121.0 (3)	C53—C52—H52A	118.1
F12—C13—C14	118.7 (3)	C52—C53—C54	119.7 (3)
C12—C13—C14	120.3 (2)	C52—C53—H53A	120.2
F13—C14—C15	120.5 (3)	C54—C53—H53A	120.2
F13—C14—C13	120.6 (3)	C55—C54—C53	116.0 (2)
C15—C14—C13	119.0 (3)	C55—C54—C54 ⁱⁱⁱ	122.0 (3)
F14—C15—C16	120.4 (2)	C53—C54—C54 ⁱⁱⁱ	122.0 (3)
F14—C15—C14	119.3 (3)	C51—C55—C54	119.9 (3)
C16—C15—C14	120.3 (2)	C51—C55—H55A	120.1
F15—C16—C15	116.8 (2)	C54—C55—H55A	120.1

O6—Mo1—O1—C1	91.22 (16)	C1—C11—C16—F15	-2.7 (4)
Mo1 ⁱ —Mo1—O1—C1	-2.45 (16)	C12—C11—C16—C15	-0.8 (4)
O5—Mo1—O1—C1	-91.80 (16)	C1—C11—C16—C15	179.2 (2)
O2 ⁱ —Mo1—O5—C3	-89.66 (18)	O5—C3—C31—C32	-72.1 (3)
O1—Mo1—O5—C3	93.32 (18)	O6 ⁱ —C3—C31—C32	105.4 (3)
Mo1 ⁱ —Mo1—O5—C3	2.82 (18)	O5—C3—C31—C36	107.0 (3)
O2 ⁱ —Mo1—O6—C3 ⁱ	98.99 (17)	O6 ⁱ —C3—C31—C36	-75.6 (3)
O1—Mo1—O6—C3 ⁱ	-83.91 (17)	C36—C31—C32—F31	178.7 (2)
Mo1 ⁱ —Mo1—O6—C3 ⁱ	6.38 (17)	C3—C31—C32—F31	-2.2 (4)
Mo1—O1—C1—O2	7.3 (3)	C36—C31—C32—C33	-1.2 (4)
Mo1—O1—C1—C11	-172.39 (15)	C3—C31—C32—C33	177.9 (2)
Mo1 ⁱ —O2—C1—O1	-8.4 (3)	F31—C32—C33—F32	0.9 (4)
Mo1 ⁱ —O2—C1—C11	171.27 (15)	C31—C32—C33—F32	-179.3 (3)
Mo1—O5—C3—O6 ⁱ	-9.2 (3)	F31—C32—C33—C34	179.8 (2)
Mo1—O5—C3—C31	168.03 (16)	C31—C32—C33—C34	-0.3 (4)
O1—C1—C11—C12	-22.4 (3)	F32—C33—C34—F33	1.7 (4)
O2—C1—C11—C12	157.9 (2)	C32—C33—C34—F33	-177.3 (2)
O1—C1—C11—C16	157.6 (2)	F32—C33—C34—C35	-179.4 (3)
O2—C1—C11—C16	-22.1 (3)	C32—C33—C34—C35	1.7 (4)
C16—C11—C12—F11	-179.9 (2)	F33—C34—C35—F34	-1.1 (4)
C1—C11—C12—F11	0.0 (4)	C33—C34—C35—F34	179.9 (3)
C16—C11—C12—C13	2.0 (4)	F33—C34—C35—C36	177.4 (3)
C1—C11—C12—C13	-178.1 (2)	C33—C34—C35—C36	-1.5 (4)
F11—C12—C13—F12	0.0 (4)	C32—C31—C36—F35	179.4 (2)
C11—C12—C13—F12	178.2 (2)	C3—C31—C36—F35	0.3 (4)
F11—C12—C13—C14	-179.9 (2)	C32—C31—C36—C35	1.3 (4)
C11—C12—C13—C14	-1.8 (4)	C3—C31—C36—C35	-177.8 (2)
F12—C13—C14—F13	1.4 (4)	F34—C35—C36—F35	0.5 (4)
C12—C13—C14—F13	-178.6 (2)	C34—C35—C36—F35	-178.0 (3)
F12—C13—C14—C15	-179.6 (3)	F34—C35—C36—C31	178.6 (3)
C12—C13—C14—C15	0.3 (4)	C34—C35—C36—C31	0.0 (4)
F13—C14—C15—F14	1.7 (4)	C52—N1—C51—C55	-0.3 (4)
C13—C14—C15—F14	-177.2 (3)	C51—N1—C52—C53	0.6 (4)
F13—C14—C15—C16	179.7 (2)	N1—C52—C53—C54	-0.1 (5)
C13—C14—C15—C16	0.8 (4)	C52—C53—C54—C55	-0.7 (4)
F14—C15—C16—F15	-0.7 (4)	C52—C53—C54—C54 ⁱⁱⁱ	179.9 (3)
C14—C15—C16—F15	-178.7 (2)	N1—C51—C55—C54	-0.5 (4)
F14—C15—C16—C11	177.5 (2)	C53—C54—C55—C51	1.0 (4)
C14—C15—C16—C11	-0.5 (4)	C54 ⁱⁱⁱ —C54—C55—C51	-179.6 (3)
C12—C11—C16—F15	177.3 (2)		

Symmetry codes: (i) $-x+2, -y+1, -z+1$; (ii) $-x+1, -y+1, -z+1$; (iii) $-x+1, -y, -z+1$.

Hydrogen-bond geometry ($\text{\AA}, ^\circ$)

<i>D</i> —H \cdots <i>A</i>	<i>D</i> —H	H \cdots <i>A</i>	<i>D</i> \cdots <i>A</i>	<i>D</i> —H \cdots <i>A</i>
C55—H55A \cdots F35	0.93	2.55	3.152 (2)	122.

supplementary materials

C51—H51A···F33^{iv}

0.93

2.78

2.987 (3)

94.

Symmetry codes: (iv) $-x+1, -y+1, -z+2$.

Fig. 1




Fig. 2




Fig. 3




Fig. 4

