

## Tetra- $\mu$ -acetato- $\kappa^8$ O:O'-bis[(3,5-di-methyl-1H-pyrazole- $\kappa N^2$ )copper(II)]

Juanita van Wyk,<sup>a</sup> Bernard Omondi<sup>b\*</sup> and James Darkwa<sup>a</sup>

<sup>a</sup>Department of Chemistry, University of Johannesburg, PO Box 524 Auckland Park, Johannesburg, 2006, South Africa, and <sup>b</sup>School of Chemistry, University of KwaZulu-Natal, Westville Campus, Private Bag X54001, Durban 4000, South Africa  
Correspondence e-mail: owaga@ukzn.ac.za


Received 5 August 2011; accepted 11 August 2011

Key indicators: single-crystal X-ray study;  $T = 100$  K; mean  $\sigma(C-C) = 0.003$  Å; disorder in main residue;  $R$  factor = 0.031;  $wR$  factor = 0.089; data-to-parameter ratio = 21.2.

The dinuclear centrosymmetric title compound,  $[Cu_2(CH_3CO_2)_4(C_5H_8N_2)_2]$ , has a distorted square-pyramidal coordination geometry around each Cu<sup>II</sup> atom in which four O atoms from the bridging acetate ligands form the basal plane while two N atoms from the pyrazole ligands occupy the apical positions. The crystal has two half molecules in the asymmetric unit with a Cu···Cu distance of 2.6762 (4) Å. Disorder was found for two O atoms and two C atoms of one acetate ligand and refined with occupancies of 0.265 (7) and 0.735 (7). The crystal also features molecules linked through two N—H···O hydrogen bonds resulting in one-dimensional chains extending along the crystallographic  $b$  axis.

### Related literature

For the properties and applications of 1H-pyrazolyl-3,5-substituted ligands, see: Deka *et al.* (2006); Guzei *et al.* (2003); Mohlala *et al.* (2005); Nelana *et al.* (2008); Ojwach *et al.* (2005).



### Experimental

#### Crystal data

$[Cu_2(CH_3CO_2)_4(C_5H_8N_2)_2]$	$\gamma = 82.354 (1)^\circ$
$M_r = 555.52$	$V = 1328.55 (12) \text{ \AA}^3$
Triclinic, $P\bar{1}$	$Z = 2$
$a = 8.1125 (4) \text{ \AA}$	Mo $K\alpha$ radiation
$b = 13.6429 (7) \text{ \AA}$	$\mu = 1.64 \text{ mm}^{-1}$
$c = 13.7755 (7) \text{ \AA}$	$T = 100 \text{ K}$
$\alpha = 61.571 (1)^\circ$	$0.39 \times 0.16 \times 0.10 \text{ mm}$
$\beta = 87.449 (1)^\circ$	

#### Data collection

Bruker X8 APEXII 4K Kappa CCD diffractometer	17190 measured reflections
Absorption correction: multi-scan ( <i>SADABS</i> ; Bruker, 2007)	6578 independent reflections
$T_{\min} = 0.566$ , $T_{\max} = 0.853$	5934 reflections with $I > 2\sigma(I)$
	$R_{\text{int}} = 0.022$

#### Refinement

$R[F^2 > 2\sigma(F^2)] = 0.031$	17 restraints
$wR(F^2) = 0.089$	H-atom parameters constrained
$S = 1.06$	$\Delta\rho_{\max} = 1.28 \text{ e \AA}^{-3}$
6578 reflections	$\Delta\rho_{\min} = -0.59 \text{ e \AA}^{-3}$
310 parameters	

**Table 1**  
Hydrogen-bond geometry (Å, °).

D—H···A	D—H	H···A	D···A	D—H···A
N2—H2···O5	0.88	1.93	2.785 (2)	163
N3—H3···O3	0.88	2	2.847 (2)	163

Data collection: *APEX2* (Bruker, 2007); cell refinement: *SAINT-Plus* (Bruker, 2007); data reduction: *SAINT-Plus* and *XPREP* (Bruker, 2007); program(s) used to solve structure: *SHELXS97* (Sheldrick, 2008); program(s) used to refine structure: *SHELXL97* (Sheldrick, 2008) and *PLATON* (Spek, 2009); molecular graphics: *DIAMOND* (Brandenburg & Putz, 2005), *ORTEP-3* (Farrugia, 1997); software used to prepare material for publication: *WinGX* (Farrugia, 1999).

The authors gratefully acknowledge the University of Johannesburg for funding and Dr Ilia Guzei for help with the refinement of the structure.

Supplementary data and figures for this paper are available from the IUCr electronic archives (Reference: HG5077).

### References

- Brandenburg, K. & Putz, H. (2005). *DIAMOND*. Crystal Impact GbR, Bonn, Germany.
- Bruker (2007). *APEX2*, *SADABS*, *SAINT-Plus* and *XPREP*. Bruker AXS Inc., Madison, Wisconsin, USA.
- Deka, K., Laskar, M. & Baruah, J. B. (2006). *Polyhedron*, **25**, 2525–2529.
- Farrugia, L. J. (1997). *J. Appl. Cryst.* **30**, 565.
- Farrugia, L. J. (1999). *J. Appl. Cryst.* **32**, 837–838.
- Guzei, I. A., Li, K., Bikhazanova, G. A., Darkwa, J. & Mapolie, S. F. (2003). *Dalton Trans.* pp. 715–722.
- Mohlala, M. S., Guzei, I. A., Darkwa, J. & Mapolie, S. F. (2005). *J. Mol. Catal. A Chem.* **241**, 93–100.
- Nelana, S. M., Kruger, G. J. & Darkwa, J. (2008). *Acta Cryst. E64*, m206–m207.
- Ojwach, S. O., Tshivhase, M. G., Guzei, I. A., Darkwa, J. & Mapolie, S. F. (2005). *Can. J. Chem.* **83**, 843–853.
- Sheldrick, G. M. (2008). *Acta Cryst. A64*, 112–122.
- Spek, A. L. (2009). *Acta Cryst. D65*, 148–155.

## **supplementary materials**

*Acta Cryst.* (2011). E67, m1299 [doi:10.1107/S1600536811032600]

## Tetra- $\mu$ -acetato- $\kappa^8$ O: $O'$ -bis[(3,5-dimethyl-1*H*-pyrazole- $\kappa N^2$ )copper(II)]

J. van Wyk, B. Omondi and J. Darkwa

### Comment

Pyrazolyl ligands containing a carbonyl linker has been utilized to prepare a number of coordination compounds with palladium salts (Guzei *et al.*, 2003; Mohlala *et al.*, 2005, Ojwach *et al.*, 2005). In these compounds the pyrazolyl carbonyl moiety appear to be robust enough to avoid hydrolysis. However in a few instances the presence of metal ions like Cu(II) (Deka *et al.*, 2006) and Pd(II) (Nelana *et al.*, 2008) appear to catalyze the hydrolysis of the benzoyl fragments. We have observed similar hydrolysis when reacting copper(II) acetate with (3,5-dimethyl-pyrazol-1-yl)-*o*-benzoyl-methane. The title compound formed from this reaction is the subject of this report. The half "solvent" molecule excluded from the structure had a total number of 30.7 electrons which is approximately half the total number of electrons that acetophenone has.

Compound (I) crystallizes with two half molecules in the assymetric unit. The compound is dinuclear with each of the Cu atoms coordinated to four O atoms and a N atom from the pyrazole ligand. The O atoms are from acetate ions, all in the equatorial positions of a slightly distorted octahedral geometry around the Cu atoms. The N atom is bound *trans* to the Cu—Cu vector completing a the distorted octahedral geometry as axial ligands.

The crystal structure of (I) is composed of two N—H $\cdots$ O hydrogen bonded chains (Table 1) that extend in the crystallographic *b* axis (Fig. 2).

### Experimental

A mixture of copper(II) acetate monohydrate (0.20 g, 1 mmol) and (3,5-dimethyl-pyrazol-1-yl)-*o*-benzyl-methane (0.20 g, 1 mmol) was refluxed in methanol (20 ml) for 4 h. The bluish-green mixture turned deep green during the course of the reaction and upon removal of the solvent a green solid residue was obtained. Recrystallization from a methanol:ethylacetate (1:2) mixture produced X-ray quality crystals after several days. Yield = 0.36 g, 59%

### Refinement

The methyl, methine and aromatic H atoms were placed in geometrically idealized positions and constrained to ride on their parent atoms, with C—H = 0.95 Å and  $U_{\text{iso}}(\text{H}) = 1.2U_{\text{eq}}(\text{C})$  for aromatic, C—H = 0.98 Å and  $U_{\text{iso}}(\text{H}) = 1.2U_{\text{eq}}(\text{C})$  for CH<sub>3</sub> and N—H = 0.88 Å and  $U_{\text{iso}}(\text{H}) = 1.2U_{\text{eq}}(\text{C})$  for NH. Half a molecule of acetophenone that resided on an inversion center was grossly disordered and was excluded using the SQUEEZE subroutine in PLATON (Spek (2009)).

# supplementary materials

---

## Figures




Fig. 1. View of (I) (50% probability displacement ellipsoids) with H atoms presented as small spheres of arbitrary radii.




Fig. 2. N—H···O hydrogen bond interactions in the crystal structure of (I). [Symmetry operators: (i) = 1 -  $x$ , 1 -  $y$ , 1 -  $z$  and (ii) = 1 -  $x$ , - $y$ , 1 -  $z$ ]

## Tetra- $\mu$ -acetato- $\kappa^8$ O: $O'$ -bis[(3,5-dimethyl-1*H*-pyrazole- $\kappa^2$ N<sup>2</sup>)copper(II)]

### Crystal data

[Cu <sub>2</sub> (C <sub>2</sub> H <sub>3</sub> O <sub>2</sub> ) <sub>4</sub> (C <sub>5</sub> H <sub>8</sub> N <sub>2</sub> ) <sub>2</sub> ]	$Z = 2$
$M_r = 555.52$	$F(000) = 572$
Triclinic, $P\bar{1}$	$D_x = 1.389 \text{ Mg m}^{-3}$
Hall symbol: -P 1	Mo $K\alpha$ radiation, $\lambda = 0.71073 \text{ \AA}$
$a = 8.1125 (4) \text{ \AA}$	Cell parameters from 17682 reflections
$b = 13.6429 (7) \text{ \AA}$	$\theta = 2.5\text{--}28.4^\circ$
$c = 13.7755 (7) \text{ \AA}$	$\mu = 1.64 \text{ mm}^{-1}$
$\alpha = 61.571 (1)^\circ$	$T = 100 \text{ K}$
$\beta = 87.449 (1)^\circ$	Block, green
$\gamma = 82.354 (1)^\circ$	$0.39 \times 0.16 \times 0.1 \text{ mm}$
$V = 1328.55 (12) \text{ \AA}^3$	

### Data collection

Bruker X8 APEXII 4K Kappa CCD diffractometer	5934 reflections with $I > 2\sigma(I)$
graphite	$R_{\text{int}} = 0.022$
$\varphi$ and $\omega$ scans	$\theta_{\text{max}} = 28.4^\circ, \theta_{\text{min}} = 2.5^\circ$
Absorption correction: multi-scan ( <i>SADABS</i> ; Bruker, 2007)	$h = -10 \rightarrow 10$
$T_{\text{min}} = 0.566, T_{\text{max}} = 0.853$	$k = -18 \rightarrow 17$
17190 measured reflections	$l = -18 \rightarrow 18$
6578 independent reflections	

### Refinement

Refinement on $F^2$	Primary atom site location: structure-invariant direct methods
Least-squares matrix: full	Secondary atom site location: difference Fourier map
$R[F^2 > 2\sigma(F^2)] = 0.031$	Hydrogen site location: inferred from neighbouring sites

$wR(F^2) = 0.089$	H-atom parameters constrained
$S = 1.06$	$w = 1/[\sigma^2(F_o^2) + (0.0503P)^2 + 0.7634P]$
6578 reflections	where $P = (F_o^2 + 2F_c^2)/3$
310 parameters	$(\Delta/\sigma)_{\text{max}} = 0.012$
17 restraints	$\Delta\rho_{\text{max}} = 1.28 \text{ e \AA}^{-3}$
	$\Delta\rho_{\text{min}} = -0.59 \text{ e \AA}^{-3}$

### *Special details*

**Experimental.** Diorder: Disorder was found for two O atoms and two C atoms of one acetate ligand in which is not an uncommon situation. The disorder was modelled for two O-, two C- and H-atoms using distance restraints and PART instructions and the total occupancy at each atom site was kept as 1 during the refinement. DELU and SIMU constraints and restraints were used on the disordered atoms. All carbon atoms involved in disorder were modelled with anisotropic thermal parameters and refined with occupancies of 0.265 (7) and 0.735 (7). The "solvent" molecule resided in a special position and looked disordered as well. Modelling the disorder only destabilized the refinement. As a result, the molecule was removed using the SQUEEZE subroutine in *PLATON* giving an *R* factor of 3.0%. H-atom Placement: All H-atoms were placed in idealized locations and refined as riding with appropriate thermal displacement coefficients  $U_{\text{iso}}(\text{H}) = 1.2$  or 1.5 times  $U_{\text{eq}}$ (bearing atom).

**Geometry.** All e.s.d.'s (except the e.s.d. in the dihedral angle between two l.s. planes) are estimated using the full covariance matrix. The cell e.s.d.'s are taken into account individually in the estimation of e.s.d.'s in distances, angles and torsion angles; correlations between e.s.d.'s in cell parameters are only used when they are defined by crystal symmetry. An approximate (isotropic) treatment of cell e.s.d.'s is used for estimating e.s.d.'s involving l.s. planes.

**Refinement.** Refinement of  $F^2$  against ALL reflections. The weighted  $R$ -factor  $wR$  and goodness of fit  $S$  are based on  $F^2$ , conventional  $R$ -factors  $R$  are based on  $F$ , with  $F$  set to zero for negative  $F^2$ . The threshold expression of  $F^2 > \sigma(F^2)$  is used only for calculating  $R$ -factors(gt) etc. and is not relevant to the choice of reflections for refinement.  $R$ -factors based on  $F^2$  are statistically about twice as large as those based on  $F$ , and  $R$ -factors based on ALL data will be even larger.

The following ALERTS were generated. Each ALERT has the format test-name\_ALERT\_alert-type\_alert-level.

PLAT094\_ALERT\_2\_C Ratio of Maximum / Minimum Residual Density ... 2.18

The solvent molecule was excluded. The remaining electron density peaks are very close to other atoms and make no chemical sense. PLAT910\_ALERT\_3\_C Missing # of FCF Reflections Below Th(Min) .... 3 PLAT911\_ALERT\_3\_C Missing # FCF Refl Between THmin & STh/L= 0.600 33 PLAT912\_ALERT\_4\_C Missing # of FCF Reflections Above STh/L= 0.600 39 PLAT913\_ALERT\_3\_C Missing # of Very Strong Reflections in FCF .... 1 PLAT002\_ALERT\_2\_G Number of Distance or Angle Restraints on AtSite 10 PLAT003\_ALERT\_2\_G Number of  $U_{iso}$  or  $U^{ij}$  Restrained Atom Sites .... 4 PLAT154\_ALERT\_1\_G The su's on the Cell Angles are Equal ..... 0.00100 Deg.

These are noted PLAT301\_ALERT\_3\_G Note: Main Residue Disorder ..... 13 Perc.

See disorder explanation above. PLAT380\_ALERT\_4\_G Check Incorrectly? Oriented X(sp2)-Methyl Moiety C18  
PLAT605\_ALERT\_4\_G Structure Contains Solvent Accessible VOIDS of. 199 A\*\*3 PLAT869\_ALERT\_4\_G ALERTS Related to  
the use of SQUEEZE Suppressed ! Solvent molecule was excluded as it was grossly disordered. PLAT764\_ALERT\_4\_G Overcom-  
plete CIF Bond List Detected (Rep/Expd). 1.17 Ratio PLAT860\_ALERT\_3\_G Note: Number of Least-Squares Restraints ..... 17  
PLAT961 ALERT 5 G Dataset Contains no Negative Intensities ..... ! Noted.

## supplementary materials

---

*Fractional atomic coordinates and isotropic or equivalent isotropic displacement parameters ( $\text{\AA}^2$ )*

	<i>x</i>	<i>y</i>	<i>z</i>	$U_{\text{iso}}^*/U_{\text{eq}}$	Occ. (<1)
Cu1	0.53478 (2)	0.494657 (16)	0.406632 (17)	0.01851 (7)	
O1	0.40264 (18)	0.64415 (11)	0.33807 (11)	0.0291 (3)	
O2	0.34793 (16)	0.65464 (11)	0.49368 (11)	0.0270 (3)	
C1	0.3431 (2)	0.69462 (15)	0.39072 (16)	0.0237 (3)	
C2	0.2597 (3)	0.81398 (16)	0.32348 (18)	0.0329 (4)	
H2A	0.1426	0.8186	0.3437	0.049*	
H2B	0.2666	0.8358	0.2447	0.049*	
H2C	0.3159	0.8648	0.3385	0.049*	
O3	0.32633 (15)	0.42403 (12)	0.43652 (12)	0.0264 (3)	
O4	0.26852 (16)	0.43494 (12)	0.59184 (11)	0.0272 (3)	
N1	0.59519 (19)	0.48564 (13)	0.25736 (13)	0.0227 (3)	
N2	0.61058 (18)	0.38656 (13)	0.25468 (13)	0.0225 (3)	
H2	0.602	0.3213	0.3131	0.027*	
C3	0.2366 (2)	0.40998 (14)	0.51864 (16)	0.0226 (3)	
C4	0.0775 (2)	0.35986 (17)	0.52824 (19)	0.0308 (4)	
H4A	0.0222	0.3494	0.5966	0.046*	
H4B	0.1034	0.2871	0.5294	0.046*	
H4C	0.0036	0.4106	0.4649	0.046*	
C5	0.6131 (3)	0.68424 (17)	0.12252 (19)	0.0398 (5)	
H5A	0.5011	0.7136	0.134	0.06*	
H5B	0.6415	0.7267	0.0447	0.06*	
H5C	0.6935	0.692	0.1687	0.06*	
C6	0.6178 (2)	0.56309 (16)	0.15334 (16)	0.0269 (4)	
C7	0.6457 (3)	0.51248 (17)	0.08502 (16)	0.0303 (4)	
H7	0.6644	0.5488	0.0079	0.036*	
C8	0.6403 (2)	0.39916 (17)	0.15302 (16)	0.0272 (4)	
C9	0.6625 (3)	0.30197 (19)	0.12979 (19)	0.0363 (5)	
H9A	0.7677	0.2549	0.1634	0.055*	
H9B	0.6644	0.3298	0.0498	0.055*	
H9C	0.5701	0.2574	0.1609	0.055*	
Cu2	0.40853 (2)	0.091442 (18)	0.42346 (2)	0.02307 (7)	
O7A	0.338 (2)	0.1009 (10)	0.5625 (7)	0.0308 (6)	0.265 (7)
O8A	0.500 (3)	-0.0504 (12)	0.6844 (12)	0.0307 (7)	0.265 (7)
C12A	0.390 (2)	0.0326 (10)	0.6618 (7)	0.0265 (7)	0.265 (7)
C13A	0.3283 (13)	0.0421 (10)	0.7642 (8)	0.0357 (8)	0.265 (7)
H13A	0.4128	0.0705	0.7894	0.054*	0.265 (7)
H13B	0.308	-0.0321	0.8231	0.054*	0.265 (7)
H13C	0.2247	0.094	0.7457	0.054*	0.265 (7)
O7	0.3327 (7)	0.1281 (3)	0.5399 (3)	0.0308 (6)	0.735 (7)
O8	0.4827 (8)	-0.0262 (4)	0.6696 (4)	0.0307 (7)	0.735 (7)
C12	0.3826 (6)	0.0633 (3)	0.6357 (3)	0.0265 (7)	0.735 (7)
C13	0.3140 (4)	0.0927 (3)	0.7245 (3)	0.0357 (8)	0.735 (7)
H13D	0.3479	0.1646	0.7103	0.054*	0.735 (7)
H13E	0.3576	0.0338	0.797	0.054*	0.735 (7)
H13F	0.1923	0.0988	0.7234	0.054*	0.735 (7)

O5	0.61101 (16)	0.16201 (11)	0.41197 (12)	0.0290 (3)
O6	0.76410 (15)	0.00869 (11)	0.53847 (12)	0.0264 (3)
N3	0.23162 (18)	0.33734 (13)	0.29863 (14)	0.0241 (3)
H3	0.2743	0.3505	0.3483	0.029*
N4	0.26197 (18)	0.23761 (13)	0.29750 (14)	0.0248 (3)
C10	0.7459 (2)	0.10916 (14)	0.46471 (16)	0.0217 (3)
C11	0.8953 (2)	0.17237 (16)	0.43519 (18)	0.0283 (4)
H11A	0.9293	0.1893	0.3602	0.042*
H11B	0.9871	0.1262	0.4874	0.042*
H11C	0.8665	0.2427	0.4387	0.042*
C14	0.0769 (3)	0.52926 (18)	0.2005 (2)	0.0438 (5)
H14A	-0.0194	0.5283	0.2464	0.066*
H14B	0.0469	0.5806	0.1228	0.066*
H14C	0.1688	0.5548	0.2226	0.066*
C15	0.1293 (2)	0.41377 (17)	0.21551 (17)	0.0295 (4)
C16	0.0898 (3)	0.36072 (18)	0.15710 (17)	0.0325 (4)
H16	0.0191	0.3919	0.0935	0.039*
C17	0.1746 (2)	0.25240 (17)	0.21025 (17)	0.0291 (4)
C18	0.1723 (3)	0.1581 (2)	0.1831 (2)	0.0442 (5)
H18A	0.2449	0.0919	0.2356	0.066*
H18B	0.2121	0.1813	0.108	0.066*
H18C	0.0585	0.1397	0.1881	0.066*

*Atomic displacement parameters ( $\text{\AA}^2$ )*

	$U^{11}$	$U^{22}$	$U^{33}$	$U^{12}$	$U^{13}$	$U^{23}$
Cu1	0.01382 (10)	0.01959 (11)	0.02424 (11)	-0.00497 (7)	-0.00013 (7)	-0.01135 (9)
O1	0.0309 (7)	0.0250 (6)	0.0293 (7)	0.0015 (5)	-0.0050 (5)	-0.0119 (6)
O2	0.0234 (6)	0.0237 (6)	0.0323 (7)	0.0002 (5)	0.0000 (5)	-0.0127 (5)
C1	0.0167 (7)	0.0199 (8)	0.0342 (9)	-0.0051 (6)	-0.0024 (7)	-0.0115 (7)
C2	0.0340 (10)	0.0204 (8)	0.0389 (11)	0.0002 (7)	-0.0056 (8)	-0.0102 (8)
O3	0.0175 (6)	0.0336 (7)	0.0382 (7)	-0.0098 (5)	0.0029 (5)	-0.0238 (6)
O4	0.0212 (6)	0.0345 (7)	0.0314 (7)	-0.0143 (5)	0.0032 (5)	-0.0176 (6)
N1	0.0203 (7)	0.0227 (7)	0.0261 (7)	-0.0055 (5)	0.0016 (5)	-0.0118 (6)
N2	0.0194 (7)	0.0231 (7)	0.0270 (7)	-0.0055 (5)	0.0019 (6)	-0.0129 (6)
C3	0.0142 (7)	0.0206 (8)	0.0342 (9)	-0.0043 (6)	-0.0016 (6)	-0.0132 (7)
C4	0.0189 (8)	0.0351 (10)	0.0491 (12)	-0.0132 (7)	0.0053 (8)	-0.0267 (9)
C5	0.0529 (13)	0.0254 (10)	0.0361 (11)	-0.0120 (9)	0.0088 (10)	-0.0095 (8)
C6	0.0270 (9)	0.0265 (9)	0.0261 (9)	-0.0064 (7)	0.0011 (7)	-0.0108 (7)
C7	0.0313 (10)	0.0348 (10)	0.0241 (9)	-0.0050 (8)	0.0017 (7)	-0.0133 (8)
C8	0.0221 (8)	0.0339 (10)	0.0296 (9)	-0.0051 (7)	0.0004 (7)	-0.0178 (8)
C9	0.0376 (11)	0.0404 (11)	0.0410 (11)	-0.0042 (9)	0.0013 (9)	-0.0276 (10)
Cu2	0.01346 (11)	0.02278 (12)	0.03971 (14)	-0.00579 (8)	0.00376 (9)	-0.01966 (10)
O7A	0.0260 (8)	0.0252 (19)	0.0478 (14)	0.0024 (16)	0.0025 (14)	-0.0243 (13)
O8A	0.0267 (17)	0.029 (2)	0.0418 (16)	-0.0021 (15)	0.0072 (12)	-0.0219 (16)
C12A	0.0203 (10)	0.028 (2)	0.0427 (16)	-0.0108 (16)	0.0121 (15)	-0.0251 (15)
C13A	0.0362 (13)	0.040 (2)	0.0327 (19)	0.0104 (15)	0.0002 (13)	-0.0219 (17)
O7	0.0260 (8)	0.0252 (19)	0.0478 (14)	0.0024 (16)	0.0025 (14)	-0.0243 (13)

## supplementary materials

---

O8	0.0267 (17)	0.029 (2)	0.0418 (16)	-0.0021 (15)	0.0072 (12)	-0.0219 (16)
C12	0.0203 (10)	0.028 (2)	0.0427 (16)	-0.0108 (16)	0.0121 (15)	-0.0251 (15)
C13	0.0362 (13)	0.040 (2)	0.0327 (19)	0.0104 (15)	0.0002 (13)	-0.0219 (17)
O5	0.0178 (6)	0.0226 (6)	0.0441 (8)	-0.0066 (5)	-0.0034 (5)	-0.0127 (6)
O6	0.0147 (5)	0.0223 (6)	0.0437 (8)	-0.0053 (5)	0.0016 (5)	-0.0162 (6)
N3	0.0181 (7)	0.0247 (7)	0.0330 (8)	-0.0053 (5)	-0.0005 (6)	-0.0157 (6)
N4	0.0186 (7)	0.0278 (8)	0.0349 (8)	-0.0079 (6)	0.0034 (6)	-0.0194 (7)
C10	0.0153 (7)	0.0212 (8)	0.0353 (9)	-0.0058 (6)	0.0042 (6)	-0.0183 (7)
C11	0.0166 (8)	0.0245 (8)	0.0461 (11)	-0.0078 (6)	0.0045 (7)	-0.0175 (8)
C14	0.0451 (13)	0.0270 (10)	0.0538 (14)	0.0009 (9)	-0.0147 (11)	-0.0148 (10)
C15	0.0233 (9)	0.0293 (9)	0.0334 (10)	-0.0073 (7)	0.0005 (7)	-0.0120 (8)
C16	0.0305 (10)	0.0382 (11)	0.0282 (9)	-0.0103 (8)	-0.0002 (8)	-0.0138 (8)
C17	0.0258 (9)	0.0365 (10)	0.0313 (9)	-0.0118 (7)	0.0051 (7)	-0.0197 (8)
C18	0.0488 (13)	0.0505 (14)	0.0502 (13)	-0.0092 (11)	-0.0038 (11)	-0.0363 (12)

### Geometric parameters ( $\text{\AA}$ , $^\circ$ )

Cu1—O2 <sup>i</sup>	1.9691 (13)	Cu2—O7A	2.032 (9)
Cu1—O1	1.9685 (13)	Cu2—N4	2.1585 (17)
Cu1—O4 <sup>i</sup>	1.9747 (12)	Cu2—Cu2 <sup>ii</sup>	2.6755 (5)
Cu1—O3	1.9879 (12)	O7A—C12A	1.283 (8)
Cu1—N1	2.1492 (16)	O8A—C12A	1.265 (9)
Cu1—Cu1 <sup>i</sup>	2.6763 (4)	O8A—Cu2 <sup>ii</sup>	1.912 (15)
O1—C1	1.261 (2)	C12A—C13A	1.534 (9)
O2—C1	1.254 (2)	C13A—H13A	0.98
O2—Cu1 <sup>i</sup>	1.9691 (13)	C13A—H13B	0.98
C1—C2	1.515 (2)	C13A—H13C	0.98
C2—H2A	0.98	O7—C12	1.236 (4)
C2—H2B	0.98	O8—C12	1.263 (4)
C2—H2C	0.98	O8—Cu2 <sup>ii</sup>	1.997 (5)
O3—C3	1.266 (2)	C12—C13	1.522 (4)
O4—C3	1.254 (2)	C13—H13D	0.98
O4—Cu1 <sup>i</sup>	1.9747 (12)	C13—H13E	0.98
N1—C6	1.338 (2)	C13—H13F	0.98
N1—N2	1.359 (2)	O5—C10	1.268 (2)
N2—C8	1.342 (2)	O6—C10	1.252 (2)
N2—H2	0.88	O6—Cu2 <sup>ii</sup>	1.9657 (12)
C3—C4	1.512 (2)	N3—C15	1.344 (2)
C4—H4A	0.98	N3—N4	1.357 (2)
C4—H4B	0.98	N3—H3	0.88
C4—H4C	0.98	N4—C17	1.340 (3)
C5—C6	1.493 (3)	C10—C11	1.509 (2)
C5—H5A	0.98	C11—H11A	0.98
C5—H5B	0.98	C11—H11B	0.98
C5—H5C	0.98	C11—H11C	0.98
C6—C7	1.404 (3)	C14—C15	1.492 (3)
C7—C8	1.380 (3)	C14—H14A	0.98
C7—H7	0.95	C14—H14B	0.98

C8—C9	1.493 (3)	C14—H14C	0.98
C9—H9A	0.98	C15—C16	1.383 (3)
C9—H9B	0.98	C16—C17	1.392 (3)
C9—H9C	0.98	C16—H16	0.95
Cu2—O8A <sup>ii</sup>	1.912 (15)	C17—C18	1.503 (3)
Cu2—O7	1.949 (3)	C18—H18A	0.98
Cu2—O6 <sup>ii</sup>	1.9657 (12)	C18—H18B	0.98
Cu2—O5	1.9756 (13)	C18—H18C	0.98
Cu2—O8 <sup>ii</sup>	1.997 (5)		
O2 <sup>i</sup> —Cu1—O1	167.13 (6)	O5—Cu2—O8 <sup>ii</sup>	88.6 (2)
O2 <sup>i</sup> —Cu1—O4 <sup>i</sup>	90.21 (6)	O8A <sup>ii</sup> —Cu2—O7A	167.1 (4)
O1—Cu1—O4 <sup>i</sup>	89.34 (6)	O7—Cu2—O7A	9.9 (3)
O2 <sup>i</sup> —Cu1—O3	87.95 (6)	O6 <sup>ii</sup> —Cu2—O7A	84.8 (5)
O1—Cu1—O3	89.68 (6)	O5—Cu2—O7A	92.1 (5)
O4 <sup>i</sup> —Cu1—O3	167.33 (6)	O8 <sup>ii</sup> —Cu2—O7A	158.3 (3)
O2 <sup>i</sup> —Cu1—N1	95.41 (6)	O8A <sup>ii</sup> —Cu2—N4	90.8 (3)
O1—Cu1—N1	97.43 (6)	O7—Cu2—N4	93.06 (12)
O4 <sup>i</sup> —Cu1—N1	95.81 (6)	O6 <sup>ii</sup> —Cu2—N4	95.65 (6)
O3—Cu1—N1	96.84 (6)	O5—Cu2—N4	97.26 (6)
O2 <sup>i</sup> —Cu1—Cu1 <sup>i</sup>	84.06 (4)	O8 <sup>ii</sup> —Cu2—N4	99.45 (13)
O1—Cu1—Cu1 <sup>i</sup>	83.11 (4)	O7A—Cu2—N4	101.9 (3)
O4 <sup>i</sup> —Cu1—Cu1 <sup>i</sup>	83.34 (4)	O8A <sup>ii</sup> —Cu2—Cu2 <sup>ii</sup>	88.0 (3)
O3—Cu1—Cu1 <sup>i</sup>	83.99 (4)	O7—Cu2—Cu2 <sup>ii</sup>	88.14 (11)
N1—Cu1—Cu1 <sup>i</sup>	179.00 (4)	O6 <sup>ii</sup> —Cu2—Cu2 <sup>ii</sup>	84.11 (4)
C1—O1—Cu1	124.01 (12)	O5—Cu2—Cu2 <sup>ii</sup>	82.99 (4)
C1—O2—Cu1 <sup>i</sup>	123.04 (12)	O8 <sup>ii</sup> —Cu2—Cu2 <sup>ii</sup>	79.36 (12)
O2—C1—O1	125.59 (17)	O7A—Cu2—Cu2 <sup>ii</sup>	79.3 (3)
O2—C1—C2	117.43 (17)	N4—Cu2—Cu2 <sup>ii</sup>	178.78 (5)
O1—C1—C2	116.98 (17)	C12A—O7A—Cu2	127.1 (9)
C1—C2—H2A	109.5	C12A—O8A—Cu2 <sup>ii</sup>	123.2 (10)
C1—C2—H2B	109.5	O8A—C12A—O7A	122.3 (10)
H2A—C2—H2B	109.5	O8A—C12A—C13A	112.9 (9)
C1—C2—H2C	109.5	O7A—C12A—C13A	124.8 (10)
H2A—C2—H2C	109.5	C12A—C13A—H13A	109.5
H2B—C2—H2C	109.5	C12A—C13A—H13B	109.5
C3—O3—Cu1	122.78 (11)	H13A—C13A—H13B	109.5
C3—O4—Cu1 <sup>i</sup>	124.51 (12)	C12A—C13A—H13C	109.5
C6—N1—N2	105.07 (15)	H13A—C13A—H13C	109.5
C6—N1—Cu1	133.35 (13)	H13B—C13A—H13C	109.5
N2—N1—Cu1	121.51 (11)	C12—O7—Cu2	118.6 (3)
C8—N2—N1	112.54 (15)	C12—O8—Cu2 <sup>ii</sup>	126.0 (3)
C8—N2—H2	123.7	O7—C12—O8	127.8 (4)
N1—N2—H2	123.7	O7—C12—C13	116.7 (3)
O4—C3—O3	125.35 (16)	O8—C12—C13	115.5 (4)

## supplementary materials

---

O4—C3—C4	117.62 (17)	C10—O5—Cu2	124.10 (12)
O3—C3—C4	117.03 (16)	C10—O6—Cu2 <sup>ii</sup>	123.70 (11)
C3—C4—H4A	109.5	C15—N3—N4	112.53 (16)
C3—C4—H4B	109.5	C15—N3—H3	123.7
H4A—C4—H4B	109.5	N4—N3—H3	123.7
C3—C4—H4C	109.5	C17—N4—N3	104.94 (16)
H4A—C4—H4C	109.5	C17—N4—Cu2	131.64 (13)
H4B—C4—H4C	109.5	N3—N4—Cu2	123.27 (12)
C6—C5—H5A	109.5	O6—C10—O5	124.78 (16)
C6—C5—H5B	109.5	O6—C10—C11	117.99 (15)
H5A—C5—H5B	109.5	O5—C10—C11	117.23 (16)
C6—C5—H5C	109.5	C10—C11—H11A	109.5
H5A—C5—H5C	109.5	C10—C11—H11B	109.5
H5B—C5—H5C	109.5	H11A—C11—H11B	109.5
N1—C6—C7	110.27 (17)	C10—C11—H11C	109.5
N1—C6—C5	121.39 (18)	H11A—C11—H11C	109.5
C7—C6—C5	128.34 (18)	H11B—C11—H11C	109.5
C8—C7—C6	105.85 (17)	C15—C14—H14A	109.5
C8—C7—H7	127.1	C15—C14—H14B	109.5
C6—C7—H7	127.1	H14A—C14—H14B	109.5
N2—C8—C7	106.27 (17)	C15—C14—H14C	109.5
N2—C8—C9	122.23 (18)	H14A—C14—H14C	109.5
C7—C8—C9	131.49 (19)	H14B—C14—H14C	109.5
C8—C9—H9A	109.5	N3—C15—C16	105.90 (18)
C8—C9—H9B	109.5	N3—C15—C14	122.15 (19)
H9A—C9—H9B	109.5	C16—C15—C14	131.9 (2)
C8—C9—H9C	109.5	C15—C16—C17	106.16 (18)
H9A—C9—H9C	109.5	C15—C16—H16	126.9
H9B—C9—H9C	109.5	C17—C16—H16	126.9
O8A <sup>ii</sup> —Cu2—O7	175.4 (6)	N4—C17—C16	110.47 (18)
O8A <sup>ii</sup> —Cu2—O6 <sup>ii</sup>	92.2 (7)	N4—C17—C18	120.98 (19)
O7—Cu2—O6 <sup>ii</sup>	89.93 (17)	C16—C17—C18	128.53 (19)
O8A <sup>ii</sup> —Cu2—O5	88.0 (7)	C17—C18—H18A	109.5
O7—Cu2—O5	89.00 (18)	C17—C18—H18B	109.5
O6 <sup>ii</sup> —Cu2—O5	167.08 (6)	H18A—C18—H18B	109.5
O8A <sup>ii</sup> —Cu2—O8 <sup>ii</sup>	8.8 (4)	C17—C18—H18C	109.5
O7—Cu2—O8 <sup>ii</sup>	167.47 (13)	H18A—C18—H18C	109.5
O6 <sup>ii</sup> —Cu2—O8 <sup>ii</sup>	89.6 (2)	H18B—C18—H18C	109.5
O2 <sup>i</sup> —Cu1—O1—C1	8.2 (3)	Cu2—O7A—C12A—O8A	3(3)
O4 <sup>i</sup> —Cu1—O1—C1	-79.85 (15)	Cu2—O7A—C12A—C13A	-178.1 (12)
O3—Cu1—O1—C1	87.51 (15)	O6 <sup>ii</sup> —Cu2—O7—C12	82.6 (4)
N1—Cu1—O1—C1	-175.63 (14)	O5—Cu2—O7—C12	-84.5 (4)
Cu1 <sup>i</sup> —Cu1—O1—C1	3.53 (14)	O8 <sup>ii</sup> —Cu2—O7—C12	-5.3 (15)
Cu1 <sup>i</sup> —O2—C1—O1	4.2 (3)	O7A—Cu2—O7—C12	24 (4)
Cu1 <sup>i</sup> —O2—C1—C2	-175.42 (12)	N4—Cu2—O7—C12	178.3 (4)

Cu1—O1—C1—O2	-5.7 (3)	Cu2 <sup>ii</sup> —Cu2—O7—C12	-1.5 (4)
Cu1—O1—C1—C2	173.89 (12)	Cu2—O7—C12—O8	1.6 (9)
O2 <sup>i</sup> —Cu1—O3—C3	85.70 (14)	Cu2—O7—C12—C13	-177.4 (3)
O1—Cu1—O3—C3	-81.65 (14)	Cu2 <sup>ii</sup> —O8—C12—O7	-0.5 (10)
O4 <sup>i</sup> —Cu1—O3—C3	3.9 (3)	Cu2 <sup>ii</sup> —O8—C12—C13	178.5 (4)
N1—Cu1—O3—C3	-179.09 (14)	O8A <sup>ii</sup> —Cu2—O5—C10	-84.2 (4)
Cu1 <sup>i</sup> —Cu1—O3—C3	1.46 (13)	O7—Cu2—O5—C10	92.26 (18)
O2 <sup>i</sup> —Cu1—N1—C6	-150.52 (17)	O6 <sup>ii</sup> —Cu2—O5—C10	7.0 (4)
O1—Cu1—N1—C6	30.33 (18)	O8 <sup>ii</sup> —Cu2—O5—C10	-75.43 (19)
O4 <sup>i</sup> —Cu1—N1—C6	-59.76 (18)	O7A—Cu2—O5—C10	82.9 (4)
O3—Cu1—N1—C6	120.91 (17)	N4—Cu2—O5—C10	-174.79 (15)
O2 <sup>i</sup> —Cu1—N1—N2	32.91 (13)	Cu2 <sup>ii</sup> —Cu2—O5—C10	4.01 (15)
O1—Cu1—N1—N2	-146.24 (12)	C15—N3—N4—C17	-0.2 (2)
O4 <sup>i</sup> —Cu1—N1—N2	123.67 (13)	C15—N3—N4—Cu2	-176.20 (12)
O3—Cu1—N1—N2	-55.66 (13)	O8A <sup>ii</sup> —Cu2—N4—C17	41.6 (7)
C6—N1—N2—C8	-0.6 (2)	O7—Cu2—N4—C17	-140.9 (2)
Cu1—N1—N2—C8	176.81 (12)	O6 <sup>ii</sup> —Cu2—N4—C17	-50.68 (17)
Cu1 <sup>i</sup> —O4—C3—O3	1.4 (3)	O5—Cu2—N4—C17	129.72 (17)
Cu1 <sup>i</sup> —O4—C3—C4	-178.07 (13)	O8 <sup>ii</sup> —Cu2—N4—C17	39.9 (3)
Cu1—O3—C3—O4	-2.1 (3)	O7A—Cu2—N4—C17	-136.6 (6)
Cu1—O3—C3—C4	177.33 (12)	O8A <sup>ii</sup> —Cu2—N4—N3	-143.5 (7)
N2—N1—C6—C7	0.6 (2)	O7—Cu2—N4—N3	34.0 (2)
Cu1—N1—C6—C7	-176.34 (13)	O6 <sup>ii</sup> —Cu2—N4—N3	124.19 (13)
N2—N1—C6—C5	-178.95 (18)	O5—Cu2—N4—N3	-55.42 (14)
Cu1—N1—C6—C5	4.1 (3)	O8 <sup>ii</sup> —Cu2—N4—N3	-145.3 (3)
N1—C6—C7—C8	-0.5 (2)	O7A—Cu2—N4—N3	38.3 (6)
C5—C6—C7—C8	179.1 (2)	Cu2 <sup>ii</sup> —O6—C10—O5	6.4 (3)
N1—N2—C8—C7	0.3 (2)	Cu2 <sup>ii</sup> —O6—C10—C11	-173.03 (13)
N1—N2—C8—C9	179.72 (17)	Cu2—O5—C10—O6	-7.4 (3)
C6—C7—C8—N2	0.1 (2)	Cu2—O5—C10—C11	172.06 (13)
C6—C7—C8—C9	-179.2 (2)	N4—N3—C15—C16	0.3 (2)
O8A <sup>ii</sup> —Cu2—O7A—C12A	8(5)	N4—N3—C15—C14	179.44 (19)
O7—Cu2—O7A—C12A	-154 (6)	N3—C15—C16—C17	-0.3 (2)
O6 <sup>ii</sup> —Cu2—O7A—C12A	85.3 (17)	C14—C15—C16—C17	-179.3 (2)
O5—Cu2—O7A—C12A	-82.2 (17)	N3—N4—C17—C16	-0.1 (2)
O8 <sup>ii</sup> —Cu2—O7A—C12A	10 (3)	Cu2—N4—C17—C16	175.49 (13)
N4—Cu2—O7A—C12A	180.0 (17)	N3—N4—C17—C18	-178.19 (18)
Cu2 <sup>ii</sup> —Cu2—O7A—C12A	0.3 (17)	Cu2—N4—C17—C18	-2.6 (3)
Cu2 <sup>ii</sup> —O8A—C12A—O7A	-5(3)	C15—C16—C17—N4	0.3 (2)
Cu2 <sup>ii</sup> —O8A—C12A—C13A	175.6 (13)	C15—C16—C17—C18	178.2 (2)

Symmetry codes: (i)  $-x+1, -y+1, -z+1$ ; (ii)  $-x+1, -y, -z+1$ .


## **supplementary materials**

---

*Hydrogen-bond geometry (Å, °)*

$D\text{---H}\cdots A$	$D\text{---H}$	$H\cdots A$	$D\cdots A$	$D\text{---H}\cdots A$
N2—H2···O5	0.88	1.93	2.785 (2)	163.
N3—H3···O3	0.88	2	2.847 (2)	163.

Fig. 1



## supplementary materials

---

Fig. 2

