

Acta Crystallographica Section E

## Structure Reports

Online

ISSN 1600-5368

8-(2-Hydroxyphenyl)-1,3-dimethyl-1*H*-purine-2,6(3*H*,7*H*)-dioneIrvin Booyen,<sup>a</sup> Thulani Hlela,<sup>a</sup> Muhammed Ismail,<sup>a</sup> Thomas Gerber,<sup>b</sup> Eric Hosten<sup>b</sup> and Richard Betz<sup>b\*</sup><sup>a</sup>University of Kwazulu-Natal, School of Chemistry, Private Bag X01, Scottsville 3209, Pietermaritzburg, South Africa, and <sup>b</sup>Nelson Mandela Metropolitan University, Summerstrand Campus, Department of Chemistry, University Way, Summerstrand, PO Box 77000, Port Elizabeth 6031, South Africa

Correspondence e-mail: richard.betz@webmail.co.za


Received 3 August 2011; accepted 10 August 2011

Key indicators: single-crystal X-ray study; *T* = 200 K; mean  $\sigma(\text{C}-\text{C}) = 0.003 \text{ \AA}$ ; *R* factor = 0.049; *wR* factor = 0.124; data-to-parameter ratio = 15.8.

The title compound,  $\text{C}_{13}\text{H}_{12}\text{N}_4\text{O}_3$ , is an imidazole derivative featuring an annealed purine ring system. The benzimidazole-inspired moiety is essentially planar (r.m.s. of all fitted non-H atoms = 0.0205 Å). An intramolecular O—H...N hydrogen bond occurs. In the crystal, intermolecular N—H...O and C—H...O hydrogen bonds are observed, which connect the molecules into chains along [110]. The shortest centroid-centroid distance between two aromatic systems is 3.7771 (11) Å.

## Related literature

For the crystal structure of benzimidazole, see: Krawczyk & Gdaniec (2005). For the crystal structure of hypoxanthinium nitrate monohydrate as an example of an oxopurine compound, see: Schmalte *et al.* (1990). For graph-set analysis of hydrogen bonds, see: Etter *et al.* (1990); Bernstein *et al.* (1995). For puckering analysis, see: Cremer & Pople (1975). For general information about the chelate effect in coordination chemistry, see: Gade (1998).



## Experimental

## Crystal data

 $\text{C}_{13}\text{H}_{12}\text{N}_4\text{O}_3$  $M_r = 272.27$ 

Monoclinic,  $P2_1/c$   
 $a = 8.6418 (5) \text{ \AA}$   
 $b = 5.9415 (3) \text{ \AA}$   
 $c = 23.4475 (10) \text{ \AA}$   
 $\beta = 91.275 (2)^\circ$   
 $V = 1203.62 (11) \text{ \AA}^3$

$Z = 4$   
 Mo  $K\alpha$  radiation  
 $\mu = 0.11 \text{ mm}^{-1}$   
 $T = 200 \text{ K}$   
 $0.41 \times 0.10 \times 0.05 \text{ mm}$

## Data collection

Bruker APEXII CCD  
 diffractometer  
 Absorption correction: multi-scan  
 (*SADABS*; Bruker, 2008)  
 $T_{\min} = 0.860$ ,  $T_{\max} = 1.000$

10068 measured reflections  
 2975 independent reflections  
 1752 reflections with  $I > 2\sigma(I)$   
 $R_{\text{int}} = 0.043$

## Refinement

$R[F^2 > 2\sigma(F^2)] = 0.049$   
 $wR(F^2) = 0.124$   
 $S = 1.01$   
 2975 reflections  
 188 parameters

H atoms treated by a mixture of independent and constrained refinement  
 $\Delta\rho_{\text{max}} = 0.25 \text{ e \AA}^{-3}$   
 $\Delta\rho_{\text{min}} = -0.21 \text{ e \AA}^{-3}$

Table 1

Hydrogen-bond geometry (Å, °).

<i>D</i> —H... <i>A</i>	<i>D</i> —H	H... <i>A</i>	<i>D</i> ... <i>A</i>	<i>D</i> —H... <i>A</i>
O3—H3...N2	0.84	1.86	2.611 (2)	148
N1—H71...O1 <sup>i</sup>	0.97 (2)	1.78 (2)	2.746 (2)	175.4 (19)
C9—H9...O1 <sup>i</sup>	0.95	2.37	3.294 (2)	164
C5—H5A...O3 <sup>ii</sup>	0.98	2.58	3.234 (2)	124

Symmetry codes: (i)  $-x + 1, -y + 1, -z$ ; (ii)  $-x + 2, -y - 1, -z$ .

Data collection: *APEX2* (Bruker, 2010); cell refinement: *SAINT* (Bruker, 2010); data reduction: *SAINT*; program(s) used to solve structure: *SHELXS97* (Sheldrick, 2008); program(s) used to refine structure: *SHELXL97* (Sheldrick, 2008); molecular graphics: *ORTEP-3* (Farrugia, 1997) and *Mercury* (Macrae *et al.*, 2008); software used to prepare material for publication: *SHELXL97* and *PLATON* (Spek, 2009).

The authors thank Mrs Patricia Pono for helpful discussions.

Supplementary data and figures for this paper are available from the IUCr electronic archives (Reference: SI2369).

## References

- Bernstein, J., Davis, R. E., Shimon, L. & Chang, N.-L. (1995). *Angew. Chem. Int. Ed. Engl.* **34**, 1555–1573.  
 Bruker (2008). *SADABS*. Bruker Inc., Madison, Wisconsin, USA.  
 Bruker (2010). *APEX2* and *SAINT*. Bruker AXS Inc., Madison, Wisconsin, USA.  
 Cremer, D. & Pople, J. A. (1975). *J. Am. Chem. Soc.* **97**, 1354–1358.  
 Etter, M. C., MacDonald, J. C. & Bernstein, J. (1990). *Acta Cryst.* **B46**, 256–262.  
 Farrugia, L. J. (1997). *J. Appl. Cryst.* **30**, 565.  
 Gade, L. H. (1998). *Koordinationschemie*, 1. Auflage. Weinheim: Wiley-VCH.  
 Krawczyk, S. & Gdaniec, M. (2005). *Acta Cryst.* **E61**, o4116–o4118.  
 Macrae, C. F., Bruno, I. J., Chisholm, J. A., Edgington, P. R., McCabe, P., Pidcock, E., Rodriguez-Monge, L., Taylor, R., van de Streek, J. & Wood, P. A. (2008). *J. Appl. Cryst.* **41**, 466–470.  
 Schmalte, H. W., Hänggi, G. & Dubler, E. (1990). *Acta Cryst.* **C46**, 340–342.  
 Sheldrick, G. M. (2008). *Acta Cryst.* **A64**, 112–122.  
 Spek, A. L. (2009). *Acta Cryst.* **D65**, 148–155.

**supplementary materials**

*Acta Cryst.* (2011). E67, o2347 [ doi:10.1107/S1600536811032417 ]

## 8-(2-Hydroxyphenyl)-1,3-dimethyl-1*H*-purine-2,6(3*H*,7*H*)-dione

I. Booyesen, T. Hlela, M. Ismail, T. Gerber, E. Hosten and R. Betz

### Comment

Chelate ligands have found widespread use in coordination chemistry due to the enhanced thermodynamic stability of resultant coordination compounds in relation to metal complexes exclusively applying comparable monodentate ligands (Gade 1998). Combining different sets of donor atoms in one chelate ligand molecule, a probe for testing and accommodating metal centers of different Lewis acidities is at hand. To enable comparative studies with envisioned coordination compounds, we determined the crystal structure of the title compound. The crystal structure of benzimidazole has been reported various times in the literature (*e.g.* Krawczyk & Gdaniec, 2005). In addition, several oxopurine derivatives have been the topic of crystal structure determinations (*e.g.* Schmalle *et al.*, 1990).

The molecule features a benzimidazole-inspired backbone comprised of a 6-amino-1,3-dimethylpyrimidine-2,4(1*H*,3*H*)-dione moiety which is annealed to a five-membered aromatic ring. This part of the molecule is essentially planar (r.m.s. of its fitted non-hydrogen atoms = 0.0205 Å). The small puckering amplitude ( $\tau = 1.6^\circ$ ) of the six-membered heterocycle precludes a conformation analysis (Cremer & Pople, 1975). The least-squares planes defined by the atoms of the phenyl ring on the one hand and the benzimidazole-type ring system on the other hand enclose an angle of  $0.45(10)^\circ$  (Fig. 1). Both C–N–C angles in the five-membered heterocycle are similar in value with  $106.25(15)^\circ$  and  $104.41(15)^\circ$ , with the smaller value found on the non-protonated nitrogen atom. However, these angles are smaller in value than the corresponding ones in hypoxanthinium nitrate monohydrate – invariably above  $108^\circ$  – where both nitrogen atoms bear a hydrogen atom (Schmalle *et al.*, 1990).

In the crystal structure, intra- as well as intermolecular hydrogen bonds and C–H $\cdots$ O contacts whose range falls by more than 0.1 Å below the sum of van-der-Waals radii of atoms participating are observed. While the intramolecular hydrogen bonds are exclusively made up by the proton of the hydroxyl group as donor and the non-protonated nitrogen atom of the five-membered heterocycle, intermolecular hydrogen bonds are solely apparent between the amino group and one of the double-bonded oxygen atoms (Table 1). The C–H $\cdots$ O contacts can be separated in two groups: while one of the nitrogen-bound methyl groups forms a C–H $\cdots$ O contact involving the oxygen atom of the hydroxyl group, one of the aromatic C–H groups acts as donor for the double-bonded oxygen atom that is already part of the N–H $\cdots$ O type hydrogen bonds. In terms of graph-set analysis (Etter *et al.*, 1990; Bernstein *et al.*, 1995), the descriptor for the classical hydrogen bonds is  $S(6)R^2_2(10)$  on the unitary level. For the C–H $\cdots$ O contacts, a  $R^2_2(16)R^2_2(18)$  descriptor on the same level is needed for description. In total, the molecules are connected to chains along [1 1 0]. The shortest intercentroid distance between two aromatic systems was measured at 3.7771(11) Å (Fig. 2).

The packing of the title compound in the crystal is shown in Figure 3.

## Experimental

The title compound was prepared by reacting 6-amino-1,3-dimethyl-5-[(*E*)-2-(hydroxy)benzylideneamino]pyrimidine-2,4-(1*H*,3*H*)-dione and  $\text{NH}_4\text{VO}_3$  in methanol. A bright yellow precipitate was filtered, washed with methanol and dried under reduced pressure. Single-crystals suitable for the X-ray diffraction study were obtained by recrystallization from ethanol-dichloromethane (*v:v* = 1:1) mixture which was left in a fridge for several days.

## Refinement

Carbon-bound H atoms were placed in calculated positions ( $\text{C}-\text{H}$  0.95 Å) and were included in the refinement in the riding model approximation, with  $U(\text{H})$  set to  $1.2U_{\text{eq}}(\text{C})$ . The H atoms of the methyl groups were allowed to rotate with a fixed angle around the  $\text{C}-\text{C}$  bond to best fit the experimental electron density (HFIX 137 in the *SHELX* program suite (Sheldrick, 2008)), with  $U(\text{H})$  set to  $1.5U_{\text{eq}}(\text{C})$ . The nitrogen-bound H atom was located on a difference Fourier map and refined freely.

## Figures




Fig. 1. The molecular structure of the title compound, with atom labels and anisotropic displacement ellipsoids (drawn at 50% probability level).




Fig. 2. Intermolecular contacts, viewed along  $[-1\ 0\ 0]$ . Symmetry operators: <sup>i</sup>  $-x + 2, -y - 1, -z$ ; <sup>ii</sup>  $-x + 1, -y + 1, -z$ .




Fig. 3. Molecular packing of the title compound, viewed along  $[0\ 1\ 0]$  (anisotropic displacement ellipsoids drawn at 50% probability level).

**8-(2-Hydroxyphenyl)-1,3-dimethyl-1H-purine-2,6(3H,7H)-dione**

*Crystal data*

$C_{13}H_{12}N_4O_3$	$F(000) = 568$
$M_r = 272.27$	$D_x = 1.502 \text{ Mg m}^{-3}$
Monoclinic, $P2_1/c$	Mo $K\alpha$ radiation, $\lambda = 0.71073 \text{ \AA}$
Hall symbol: -P 2ybc	Cell parameters from 1955 reflections
$a = 8.6418 (5) \text{ \AA}$	$\theta = 2.4\text{--}27.2^\circ$
$b = 5.9415 (3) \text{ \AA}$	$\mu = 0.11 \text{ mm}^{-1}$
$c = 23.4475 (10) \text{ \AA}$	$T = 200 \text{ K}$
$\beta = 91.275 (2)^\circ$	Platelet, yellow
$V = 1203.62 (11) \text{ \AA}^3$	$0.41 \times 0.10 \times 0.05 \text{ mm}$
$Z = 4$	

*Data collection*

Bruker APEXII CCD diffractometer	2975 independent reflections
Radiation source: fine-focus sealed tube graphite	1752 reflections with $I > 2\sigma(I)$
$\varphi$ and $\omega$ scans	$R_{\text{int}} = 0.043$
Absorption correction: multi-scan (SADABS; Bruker, 2008)	$\theta_{\text{max}} = 28.3^\circ$ , $\theta_{\text{min}} = 2.4^\circ$
$T_{\text{min}} = 0.860$ , $T_{\text{max}} = 1.000$	$h = -11 \rightarrow 11$
10068 measured reflections	$k = -7 \rightarrow 6$
	$l = -31 \rightarrow 27$

*Refinement*

Refinement on $F^2$	Primary atom site location: structure-invariant direct methods
Least-squares matrix: full	Secondary atom site location: difference Fourier map
$R[F^2 > 2\sigma(F^2)] = 0.049$	Hydrogen site location: inferred from neighbouring sites
$wR(F^2) = 0.124$	H atoms treated by a mixture of independent and constrained refinement
$S = 1.01$	$w = 1/[\sigma^2(F_o^2) + (0.0584P)^2 + 0.0815P]$
2975 reflections	where $P = (F_o^2 + 2F_c^2)/3$
188 parameters	$(\Delta/\sigma)_{\text{max}} < 0.001$
0 restraints	$\Delta\rho_{\text{max}} = 0.25 \text{ e \AA}^{-3}$
	$\Delta\rho_{\text{min}} = -0.21 \text{ e \AA}^{-3}$

*Fractional atomic coordinates and isotropic or equivalent isotropic displacement parameters ( $\text{\AA}^2$ )*

	<i>x</i>	<i>y</i>	<i>z</i>	$U_{\text{iso}}^*/U_{\text{eq}}$
O1	0.47991 (16)	0.4606 (2)	0.07866 (5)	0.0356 (4)
O2	0.66823 (16)	-0.0720 (2)	0.20267 (6)	0.0390 (4)

## supplementary materials

---

O3	0.94790 (18)	-0.3204 (3)	-0.06725 (6)	0.0454 (4)
H3	0.9105	-0.2923	-0.0353	0.068*
N1	0.65931 (17)	0.2091 (3)	-0.01367 (6)	0.0260 (4)
H71	0.615 (2)	0.330 (4)	-0.0365 (9)	0.049 (7)*
N2	0.79184 (18)	-0.1028 (3)	0.00928 (6)	0.0273 (4)
N3	0.73063 (17)	-0.1106 (3)	0.10978 (6)	0.0269 (4)
N4	0.57335 (17)	0.1913 (3)	0.13977 (6)	0.0259 (4)
C1	0.6392 (2)	0.1685 (3)	0.04385 (7)	0.0247 (4)
C2	0.7221 (2)	-0.0211 (3)	0.05591 (7)	0.0238 (4)
C3	0.6584 (2)	-0.0024 (3)	0.15408 (8)	0.0267 (4)
C4	0.5573 (2)	0.2871 (3)	0.08562 (7)	0.0254 (4)
C5	0.8231 (2)	-0.3115 (3)	0.12198 (8)	0.0323 (5)
H5A	0.8296	-0.4032	0.0874	0.048*
H5B	0.7742	-0.3993	0.1520	0.048*
H5C	0.9274	-0.2668	0.1347	0.048*
C6	0.4923 (2)	0.3034 (4)	0.18658 (8)	0.0342 (5)
H6A	0.5389	0.4513	0.1938	0.051*
H6B	0.5012	0.2112	0.2212	0.051*
H6C	0.3827	0.3221	0.1759	0.051*
C7	0.7514 (2)	0.0418 (3)	-0.03268 (8)	0.0254 (4)
C8	0.8012 (2)	0.0162 (3)	-0.09136 (8)	0.0258 (4)
C9	0.7563 (2)	0.1677 (3)	-0.13407 (8)	0.0307 (5)
H9	0.6913	0.2908	-0.1248	0.037*
C10	0.8043 (2)	0.1426 (4)	-0.18967 (8)	0.0352 (5)
H10	0.7732	0.2476	-0.2182	0.042*
C11	0.8985 (2)	-0.0377 (4)	-0.20303 (8)	0.0383 (5)
H11	0.9316	-0.0568	-0.2411	0.046*
C12	0.9443 (2)	-0.1892 (4)	-0.16178 (8)	0.0391 (5)
H12	1.0093	-0.3115	-0.1716	0.047*
C13	0.8968 (2)	-0.1664 (3)	-0.10562 (8)	0.0316 (5)

### Atomic displacement parameters ( $\text{\AA}^2$ )

	$U^{11}$	$U^{22}$	$U^{33}$	$U^{12}$	$U^{13}$	$U^{23}$
O1	0.0444 (8)	0.0353 (8)	0.0272 (8)	0.0189 (7)	0.0052 (6)	0.0019 (6)
O2	0.0474 (9)	0.0435 (9)	0.0265 (8)	0.0104 (7)	0.0069 (6)	0.0106 (7)
O3	0.0634 (10)	0.0420 (10)	0.0311 (8)	0.0278 (8)	0.0109 (7)	0.0036 (7)
N1	0.0290 (9)	0.0272 (9)	0.0219 (8)	0.0080 (7)	0.0029 (6)	0.0010 (7)
N2	0.0320 (9)	0.0256 (9)	0.0245 (8)	0.0059 (7)	0.0016 (7)	-0.0002 (7)
N3	0.0296 (9)	0.0270 (9)	0.0240 (8)	0.0059 (7)	0.0022 (6)	0.0037 (7)
N4	0.0284 (8)	0.0283 (9)	0.0212 (8)	0.0062 (7)	0.0047 (6)	0.0010 (7)
C1	0.0278 (10)	0.0251 (10)	0.0212 (9)	0.0041 (8)	0.0024 (7)	0.0008 (8)
C2	0.0248 (9)	0.0243 (10)	0.0224 (9)	0.0006 (8)	0.0029 (7)	-0.0011 (8)
C3	0.0261 (10)	0.0269 (11)	0.0270 (10)	0.0011 (8)	0.0028 (8)	0.0032 (8)
C4	0.0252 (10)	0.0270 (11)	0.0241 (10)	0.0034 (9)	0.0015 (8)	0.0006 (8)
C5	0.0356 (11)	0.0274 (11)	0.0339 (11)	0.0090 (9)	0.0015 (8)	0.0052 (9)
C6	0.0396 (11)	0.0399 (13)	0.0233 (10)	0.0084 (10)	0.0088 (8)	0.0003 (9)
C7	0.0246 (10)	0.0257 (10)	0.0257 (10)	0.0038 (8)	0.0012 (7)	-0.0019 (8)

C8	0.0274 (10)	0.0282 (10)	0.0219 (9)	0.0036 (8)	0.0031 (7)	-0.0025 (8)
C9	0.0309 (10)	0.0341 (12)	0.0271 (10)	0.0063 (9)	0.0028 (8)	-0.0014 (9)
C10	0.0370 (11)	0.0425 (13)	0.0261 (10)	0.0046 (10)	0.0019 (9)	0.0018 (10)
C11	0.0439 (12)	0.0475 (14)	0.0239 (11)	0.0060 (11)	0.0067 (9)	-0.0053 (10)
C12	0.0458 (13)	0.0388 (13)	0.0330 (12)	0.0124 (11)	0.0090 (9)	-0.0070 (10)
C13	0.0352 (11)	0.0308 (11)	0.0289 (11)	0.0058 (9)	0.0026 (8)	-0.0012 (9)

*Geometric parameters (Å, °)*

O1—C4	1.237 (2)	C5—H5A	0.9800
O2—C3	1.213 (2)	C5—H5B	0.9800
O3—C13	1.351 (2)	C5—H5C	0.9800
O3—H3	0.8400	C6—H6A	0.9800
N1—C7	1.355 (2)	C6—H6B	0.9800
N1—C1	1.385 (2)	C6—H6C	0.9800
N1—H71	0.97 (2)	C7—C8	1.459 (2)
N2—C7	1.346 (2)	C8—C9	1.395 (3)
N2—C2	1.351 (2)	C8—C13	1.409 (3)
N3—C2	1.371 (2)	C9—C10	1.385 (3)
N3—C3	1.382 (2)	C9—H9	0.9500
N3—C5	1.461 (2)	C10—C11	1.385 (3)
N4—C4	1.396 (2)	C10—H10	0.9500
N4—C3	1.402 (2)	C11—C12	1.373 (3)
N4—C6	1.474 (2)	C11—H11	0.9500
C1—C2	1.361 (3)	C12—C13	1.394 (3)
C1—C4	1.410 (3)	C12—H12	0.9500
C13—O3—H3	109.5	H5B—C5—H5C	109.5
C7—N1—C1	106.25 (15)	N4—C6—H6A	109.5
C7—N1—H71	126.1 (13)	N4—C6—H6B	109.5
C1—N1—H71	127.6 (13)	H6A—C6—H6B	109.5
C7—N2—C2	104.41 (15)	N4—C6—H6C	109.5
C2—N3—C3	119.72 (16)	H6A—C6—H6C	109.5
C2—N3—C5	121.03 (15)	H6B—C6—H6C	109.5
C3—N3—C5	119.14 (15)	N2—C7—N1	111.87 (15)
C4—N4—C3	126.42 (15)	N2—C7—C8	123.05 (16)
C4—N4—C6	116.97 (15)	N1—C7—C8	125.08 (16)
C3—N4—C6	116.61 (15)	C9—C8—C13	118.81 (17)
C2—C1—N1	105.66 (16)	C9—C8—C7	121.77 (17)
C2—C1—C4	122.63 (17)	C13—C8—C7	119.42 (17)
N1—C1—C4	131.69 (17)	C10—C9—C8	121.44 (18)
N2—C2—C1	111.80 (16)	C10—C9—H9	119.3
N2—C2—N3	126.19 (17)	C8—C9—H9	119.3
C1—C2—N3	122.00 (16)	C9—C10—C11	119.09 (19)
O2—C3—N3	121.57 (17)	C9—C10—H10	120.5
O2—C3—N4	122.01 (17)	C11—C10—H10	120.5
N3—C3—N4	116.43 (16)	C12—C11—C10	120.61 (18)
O1—C4—N4	120.16 (16)	C12—C11—H11	119.7
O1—C4—C1	127.08 (17)	C10—C11—H11	119.7
N4—C4—C1	112.74 (16)	C11—C12—C13	121.01 (19)

## supplementary materials

N3—C5—H5A	109.5	C11—C12—H12	119.5
N3—C5—H5B	109.5	C13—C12—H12	119.5
H5A—C5—H5B	109.5	O3—C13—C12	117.67 (18)
N3—C5—H5C	109.5	O3—C13—C8	123.27 (17)
H5A—C5—H5C	109.5	C12—C13—C8	119.05 (18)
C7—N1—C1—C2	0.44 (19)	C2—C1—C4—O1	179.62 (19)
C7—N1—C1—C4	179.0 (2)	N1—C1—C4—O1	1.2 (3)
C7—N2—C2—C1	0.3 (2)	C2—C1—C4—N4	1.1 (3)
C7—N2—C2—N3	-179.25 (17)	N1—C1—C4—N4	-177.26 (18)
N1—C1—C2—N2	-0.5 (2)	C2—N2—C7—N1	0.0 (2)
C4—C1—C2—N2	-179.25 (17)	C2—N2—C7—C8	-179.54 (17)
N1—C1—C2—N3	179.11 (16)	C1—N1—C7—N2	-0.3 (2)
C4—C1—C2—N3	0.4 (3)	C1—N1—C7—C8	179.23 (17)
C3—N3—C2—N2	177.08 (17)	N2—C7—C8—C9	179.44 (17)
C5—N3—C2—N2	0.9 (3)	N1—C7—C8—C9	0.0 (3)
C3—N3—C2—C1	-2.5 (3)	N2—C7—C8—C13	-0.4 (3)
C5—N3—C2—C1	-178.64 (17)	N1—C7—C8—C13	-179.78 (18)
C2—N3—C3—O2	-177.22 (18)	C13—C8—C9—C10	-0.4 (3)
C5—N3—C3—O2	-1.0 (3)	C7—C8—C9—C10	179.85 (18)
C2—N3—C3—N4	2.8 (2)	C8—C9—C10—C11	0.3 (3)
C5—N3—C3—N4	179.07 (16)	C9—C10—C11—C12	-0.3 (3)
C4—N4—C3—O2	178.72 (17)	C10—C11—C12—C13	0.3 (3)
C6—N4—C3—O2	-1.7 (3)	C11—C12—C13—O3	-179.3 (2)
C4—N4—C3—N3	-1.3 (3)	C11—C12—C13—C8	-0.4 (3)
C6—N4—C3—N3	178.23 (16)	C9—C8—C13—O3	179.20 (18)
C3—N4—C4—O1	-179.24 (17)	C7—C8—C13—O3	-1.0 (3)
C6—N4—C4—O1	1.2 (3)	C9—C8—C13—C12	0.4 (3)
C3—N4—C4—C1	-0.6 (3)	C7—C8—C13—C12	-179.82 (18)
C6—N4—C4—C1	179.82 (16)		

### Hydrogen-bond geometry ( $\text{\AA}$ , $^\circ$ )

$D-H\cdots A$	$D-H$	$H\cdots A$	$D\cdots A$	$D-H\cdots A$
O3—H3 $\cdots$ N2	0.84	1.86	2.611 (2)	148.
N1—H71 $\cdots$ O1 <sup>i</sup>	0.97 (2)	1.78 (2)	2.746 (2)	175.4 (19)
C9—H9 $\cdots$ O1 <sup>i</sup>	0.95	2.37	3.294 (2)	164.
C5—H5A $\cdots$ O3 <sup>ii</sup>	0.98	2.58	3.234 (2)	124.

Symmetry codes: (i)  $-x+1, -y+1, -z$ ; (ii)  $-x+2, -y-1, -z$ .



Fig. 1




Fig. 2




Fig. 3

