

(E)-2-[[1-Carboxy-2-(1*H*-indol-3-yl)ethyl-iminio]methyl]-6-hydroxyphenolateSalah Ahmed Ba-Salamah,^a Naser Eltaher Eltayeb,^{a,b} Siang Guan Teoh^{a*} and Kong Mun Lo^c^aSchool of Chemical Sciences, Universiti Sains Malaysia, Minden, Penang, Malaysia,^bDepartment of Chemistry, International University of Africa, Sudan, and ^cChemistry Department, Faculty of Science, University of Malaya, Malaysia

Correspondence e-mail: sgteoh@usm.my


Received 4 August 2011; accepted 5 August 2011

Key indicators: single-crystal X-ray study; $T = 100$ K; mean $\sigma(\text{C}-\text{C}) = 0.004$ Å; R factor = 0.038; wR factor = 0.093; data-to-parameter ratio = 8.4.

In the zwitterionic title compound, $\text{C}_{18}\text{H}_{16}\text{N}_2\text{O}_4$, the dihedral angle between the planes of the benzene and indole rings is 26.38 (10°). An intramolecular $\text{N}-\text{H}\cdots\text{O}$ hydrogen bond generates an $S(6)$ ring motif. In the crystal, molecules are linked through $\text{N}-\text{H}\cdots\text{O}$, $\text{O}-\text{H}\cdots\text{O}$ and $\text{C}-\text{H}\cdots\text{O}$ hydrogen bonds into infinite chains propagating in $[010]$.

Related literature

For a related structure and background to Schiff bases, see: Ba-Salamah *et al.* (2011). For other related structures, see: Eltayeb *et al.* (2010*a,b,c*). For reference bond lengths, see: Allen *et al.* (1987). For graph-set theory, see: Bernstein *et al.* (1995).

**Experimental***Crystal data*

$\text{C}_{18}\text{H}_{16}\text{N}_2\text{O}_4$
 $M_r = 324.33$
 Monoclinic, $P2_1$
 $a = 8.4351$ (2) Å

$b = 9.3038$ (3) Å
 $c = 9.5023$ (3) Å
 $\beta = 98.683$ (2)°
 $V = 737.18$ (4) Å³

$Z = 2$
 Mo $K\alpha$ radiation
 $\mu = 0.11$ mm⁻¹

$T = 100$ K
 $0.32 \times 0.22 \times 0.20$ mm

Data collection

Bruker SMART APEXII CCD diffractometer
 Absorption correction: multi-scan (SADABS; Bruker, 2009)
 $T_{\min} = 0.598$, $T_{\max} = 0.746$

7473 measured reflections
 1949 independent reflections
 1781 reflections with $I > 2\sigma(I)$
 $R_{\text{int}} = 0.046$

Refinement

$R[F^2 > 2\sigma(F^2)] = 0.038$
 $wR(F^2) = 0.093$
 $S = 1.14$
 1949 reflections
 233 parameters
 1 restraint

H atoms treated by a mixture of independent and constrained refinement
 $\Delta\rho_{\text{max}} = 0.27$ e Å⁻³
 $\Delta\rho_{\text{min}} = -0.23$ e Å⁻³

Table 1

Hydrogen-bond geometry (Å, °).

$D-\text{H}\cdots A$	$D-\text{H}$	$\text{H}\cdots A$	$D\cdots A$	$D-\text{H}\cdots A$
$\text{N1}-\text{H1N1}\cdots\text{O3}$	0.97 (3)	1.86 (3)	2.646 (2)	136 (2)
$\text{N2}-\text{H1N2}\cdots\text{O3}^i$	0.90 (3)	2.13 (3)	3.020 (2)	170 (2)
$\text{O2}-\text{H1O2}\cdots\text{O3}^{ii}$	0.89 (3)	1.64 (4)	2.526 (2)	174 (4)
$\text{O4}-\text{H1O4}\cdots\text{O3}$	0.89 (3)	2.45 (3)	2.823 (2)	106 (2)
$\text{O4}-\text{H1O4}\cdots\text{O1}^i$	0.89 (3)	1.83 (3)	2.665 (2)	156 (3)
$\text{C7}-\text{H7}\cdots\text{O4}^{iii}$	0.95	2.49	3.197 (3)	132

Symmetry codes: (i) $-x, y + \frac{1}{2}, -z$; (ii) $-x, y - \frac{1}{2}, -z$; (iii) $-x + 1, y - \frac{1}{2}, -z$.

Data collection: APEX2 (Bruker, 2009); cell refinement: SAINT (Bruker, 2009); data reduction: SAINT; program(s) used to solve structure: SHELXTL (Sheldrick, 2008); program(s) used to refine structure: SHELXTL; molecular graphics: SHELXTL; software used to prepare material for publication: SHELXTL and PLATON (Spek, 2009).

The authors thank the Malaysian Government and Universiti Sains Malaysia for the RU research grant (1001/PKIMIA/815067). NEE thanks Universiti Sains Malaysia for a post-doctoral fellowship and the International University of Africa (Sudan) for providing study leave. SAB thanks the Ministry of Higher Education and Scientific Research (Yemen) for a scholarship.

Supplementary data and figures for this paper are available from the IUCr electronic archives (Reference: HB6347).

References

- Allen, F. H., Kennard, O., Watson, D. G., Brammer, L., Orpen, A. G. & Taylor, R. (1987). *J. Chem. Soc. Perkin Trans. 2*, pp. S1–19.
- Ba-Salamah, S. A., Eltayeb, N. E., Teoh, S. G. & Lo, K. M. (2011). *Acta Cryst. E67*, o2113–o2114.
- Bernstein, J., Davis, R. E., Shimon, L. & Chang, N.-L. (1995). *Angew. Chem. Int. Ed. Engl.* **34**, 1555–1573.
- Bruker (2009). APEX2, SAINT and SADABS. Bruker AXS Inc., Madison, Wisconsin, USA.
- Eltayeb, N. E., Teoh, S. G., Chantrapomma, S. & Fun, H.-K. (2010a). *Acta Cryst. E66*, o934–o935.
- Eltayeb, N. E., Teoh, S. G., Fun, H.-K. & Chantrapomma, S. (2010b). *Acta Cryst. E66*, o1262–o1263.
- Eltayeb, N. E., Teoh, S. G., Fun, H.-K. & Chantrapomma, S. (2010c). *Acta Cryst. E66*, o1536–o1537.
- Sheldrick, G. M. (2008). *Acta Cryst. A64*, 112–122.
- Spek, A. L. (2009). *Acta Cryst. D65*, 148–155.

supplementary materials

Acta Cryst. (2011). E67, o2389 [doi:10.1107/S1600536811031709]

(*E*)-2-([1-Carboxy-2-(1*H*-indol-3-yl)ethyliminio]methyl)-6-hydroxyphenolate

S. A. Ba-Salamah, N. E. Eltayeb, S. G. Teoh and K. M. Lo

Comment

Recently, we reported the crystal structure of (*E*)-2-(2,4-dihydroxybenzylideneammonio)-3-(1*H*-indol-3-yl)propanoate (Ba-Salamah *et al.* 2011). In this paper, we report the crystal structure of the title compound, (I), (Fig. 1), obtained by the reaction of *L*-tryptophan and 2,3-dihydroxybenzaldehyde.

In the zwitterionic title compound, C₁₈H₁₆N₂O₄, the dihedral angle between the planes of the benzene and indole rings is 26.38 (10)°. Bond lengths have normal values (Allen *et al.*, 1987). The C1 in the benzene ring and C11 in the indole ring are connected together by a chain of four atoms, C7/N1/C8/C10 which has torsion angle 101.2 (2)°. The torsion angles of the chain C8/N1/C7/C1 and N1/C8/C10/C11 are -179.0 (2)° and 64.4 (2)°, respectively. The title molecule has a zwitterionic form with an intramolecular N1—H1N1···O3 hydrogen bond (Table 1, Fig. 1) which generates an S(6) ring motif (Bernstein *et al.*, 1995). The C7—N1 [1.295 (3) Å] and C2—O3 [1.318 (3) Å] bond distances are comparable to those [1.3129 (19) and 1.2915 (17) Å; 1.310 (2) and 1.304 (2) Å; 1.302 (2) and 1.316 (2) Å] observed in similar zwitterionic structures (Eltayeb *et al.*, 2010*a,b,c*).

In the crystal structure of (I) as shown in Fig. 2, the molecules are linked through N—H···O, O—H···O and C—H···O hydrogen bonds (Table 1). C—H···π interactions are also present; C8—H8···Cg3ⁱ = 2.70 Å, C16—H16···Cg2ⁱⁱ = 2.81 Å. Cg3 and Cg2 are centroids of C12—C17 and C1—C6 rings respectively, [symmetry codes: (i) = -*X*, -1/2+*Y*, 1-*Z*; (ii) = -*X*, 1/2+*Y*, -*Z*]. The molecules are linked into infinite one-dimensional chains along the *b* axis.

Experimental

To a 100-ml round-bottom flask were added 20 ml (3:1) methanol-water and 2 mmol of *L*-tryptophan (0.416 g), the mixture stirred to form a clear colorless solution, 2 mmol of 2,3-dihydroxybenzaldehyde (0.276 g) added to form a yellow clear solution. The mixture was refluxed under heating and stirring for seven hours after which the mixture was filtered and left to stand at room temperature for 12 h. Dark orange crystals which appeared, were filtered after two days and recrystallized from ethanol to yield orange blocks of (I), then left to dry. The IR showed broad peak due to O—H (carboxylic acid and phenol) stretch superimposed on the sharp band due to N—H stretch at 3500–3100 cm⁻¹. The C=O stretch (1708 cm⁻¹), C=N stretch (1644 cm⁻¹), C—O stretch (1230 cm⁻¹), O—H bends (1460 cm⁻¹).

Refinement

The N- and O-bound H atoms were located in a difference Fourier map and were refined freely. The remaining H atoms were positioned geometrically and refined using a riding model with C—H = 0.95 or 1.00 Å and $U_{\text{iso}}(\text{H}) = 1.2U_{\text{eq}}(\text{C})$. In the absence of significant anomalous scattering effects, 1577 Friedel pairs were merged.

Figures




Fig. 1. The molecular structure of the title compound, with 50% probability ellipsoids for non-H atoms.




Fig. 2. The crystal packing of title compound, viewed down *c* axis.

(*E*)-2-[[1-Carboxy-2-(1*H*-indol-3-yl)ethyliminio]methyl]-6- hydroxyphenolate

Crystal data

$C_{18}H_{16}N_2O_4$

$M_r = 324.33$

Monoclinic, $P2_1$

Hall symbol: P 2yb

$a = 8.4351$ (2) Å

$b = 9.3038$ (3) Å

$c = 9.5023$ (3) Å

$\beta = 98.683$ (2)°

$V = 737.18$ (4) Å³

$Z = 2$

$F(000) = 340$

$D_x = 1.461$ Mg m⁻³

Mo $K\alpha$ radiation, $\lambda = 0.71073$ Å

Cell parameters from 1920 reflections

$\theta = 3.0$ – 27.7 °

$\mu = 0.11$ mm⁻¹

$T = 100$ K

Block, orange

$0.32 \times 0.22 \times 0.20$ mm

Data collection

Bruker SMART APEXII CCD diffractometer

Radiation source: fine-focus sealed tube

graphite

φ and ω scans

Absorption correction: multi-scan (*SADABS*; Bruker, 2009)

$T_{\min} = 0.598$, $T_{\max} = 0.746$

7473 measured reflections

1949 independent reflections

1781 reflections with $I > 2\sigma(I)$

$R_{\text{int}} = 0.046$

$\theta_{\text{max}} = 28.4$ °, $\theta_{\text{min}} = 2.2$ °

$h = -11 \rightarrow 11$

$k = -12 \rightarrow 12$

$l = -12 \rightarrow 12$

Refinement

Refinement on F^2

Primary atom site location: structure-invariant direct methods

Least-squares matrix: full

$$R[F^2 > 2\sigma(F^2)] = 0.038$$

$$wR(F^2) = 0.093$$

$$S = 1.14$$

1949 reflections

233 parameters

1 restraint

Secondary atom site location: difference Fourier map

Hydrogen site location: inferred from neighbouring sites

H atoms treated by a mixture of independent and constrained refinement

$$w = 1/[\sigma^2(F_o^2) + (0.0454P)^2 + 0.0637P]$$

$$\text{where } P = (F_o^2 + 2F_c^2)/3$$

$$(\Delta/\sigma)_{\max} < 0.001$$

$$\Delta\rho_{\max} = 0.27 \text{ e } \text{\AA}^{-3}$$

$$\Delta\rho_{\min} = -0.23 \text{ e } \text{\AA}^{-3}$$

Special details

Geometry. All e.s.d.'s (except the e.s.d. in the dihedral angle between two l.s. planes) are estimated using the full covariance matrix. The cell e.s.d.'s are taken into account individually in the estimation of e.s.d.'s in distances, angles and torsion angles; correlations between e.s.d.'s in cell parameters are only used when they are defined by crystal symmetry. An approximate (isotropic) treatment of cell e.s.d.'s is used for estimating e.s.d.'s involving l.s. planes.

Refinement. Refinement of F^2 against ALL reflections. The weighted R -factor wR and goodness of fit S are based on F^2 , conventional R -factors R are based on F , with F set to zero for negative F^2 . The threshold expression of $F^2 > \sigma(F^2)$ is used only for calculating R -factors(gt) *etc.* and is not relevant to the choice of reflections for refinement. R -factors based on F^2 are statistically about twice as large as those based on F , and R -factors based on ALL data will be even larger.

Fractional atomic coordinates and isotropic or equivalent isotropic displacement parameters (\AA^2)

	<i>x</i>	<i>y</i>	<i>z</i>	$U_{\text{iso}}^*/U_{\text{eq}}$
O1	-0.07419 (19)	-0.1334 (2)	0.34014 (17)	0.0232 (4)
O2	-0.0062 (2)	-0.1221 (2)	0.12218 (17)	0.0211 (4)
O3	0.21206 (18)	0.18228 (18)	-0.05777 (17)	0.0182 (3)
O4	0.3782 (2)	0.2819 (2)	-0.27523 (17)	0.0208 (4)
N1	0.2349 (2)	0.0546 (2)	0.1933 (2)	0.0153 (4)
N2	-0.1227 (2)	0.4135 (2)	0.2298 (2)	0.0195 (4)
C1	0.4611 (3)	0.1047 (2)	0.0751 (2)	0.0155 (4)
C2	0.3695 (3)	0.1726 (2)	-0.0437 (2)	0.0170 (5)
C3	0.4555 (3)	0.2223 (3)	-0.1530 (2)	0.0171 (5)
C4	0.6201 (3)	0.2052 (3)	-0.1389 (3)	0.0197 (5)
H4	0.6748	0.2395	-0.2126	0.024*
C5	0.7084 (3)	0.1396 (3)	-0.0203 (3)	0.0201 (5)
H5	0.8216	0.1306	-0.0129	0.024*
C6	0.6296 (3)	0.0881 (3)	0.0856 (3)	0.0189 (5)
H6	0.6881	0.0413	0.1659	0.023*
C7	0.3873 (3)	0.0479 (3)	0.1886 (2)	0.0159 (4)
H7	0.4539	0.0024	0.2653	0.019*
C8	0.1609 (3)	-0.0021 (3)	0.3116 (2)	0.0155 (4)
H8	0.2412	-0.0673	0.3675	0.019*
C9	0.0139 (3)	-0.0931 (3)	0.2577 (2)	0.0163 (5)
C10	0.1237 (3)	0.1170 (3)	0.4127 (2)	0.0170 (4)
H10A	0.0851	0.0722	0.4957	0.020*

supplementary materials

H10B	0.2245	0.1686	0.4482	0.020*
C11	0.0013 (3)	0.2246 (3)	0.3481 (2)	0.0172 (5)
C12	-0.1594 (3)	0.2345 (3)	0.3819 (2)	0.0158 (4)
C13	-0.2460 (3)	0.1543 (3)	0.4697 (2)	0.0182 (5)
H13	-0.1991	0.0732	0.5208	0.022*
C14	-0.4016 (3)	0.1962 (3)	0.4803 (3)	0.0200 (5)
H14	-0.4619	0.1427	0.5389	0.024*
C15	-0.4712 (3)	0.3162 (3)	0.4059 (3)	0.0227 (5)
H15	-0.5772	0.3437	0.4167	0.027*
C16	-0.3889 (3)	0.3954 (3)	0.3172 (3)	0.0205 (5)
H16	-0.4373	0.4754	0.2654	0.025*
C17	-0.2323 (3)	0.3536 (3)	0.3065 (2)	0.0176 (5)
C18	0.0167 (3)	0.3350 (3)	0.2562 (2)	0.0189 (5)
H18	0.1109	0.3548	0.2160	0.023*
H1N1	0.171 (3)	0.093 (4)	0.108 (3)	0.034 (8)*
H1N2	-0.141 (3)	0.490 (4)	0.171 (3)	0.024 (7)*
H1O2	-0.084 (4)	-0.187 (4)	0.102 (3)	0.033 (8)*
H1O4	0.273 (4)	0.292 (4)	-0.278 (3)	0.046 (10)*

Atomic displacement parameters (\AA^2)

	U^{11}	U^{22}	U^{33}	U^{12}	U^{13}	U^{23}
O1	0.0195 (8)	0.0321 (10)	0.0189 (8)	-0.0065 (7)	0.0056 (6)	0.0020 (8)
O2	0.0196 (8)	0.0255 (10)	0.0191 (8)	-0.0075 (7)	0.0058 (6)	-0.0043 (7)
O3	0.0124 (7)	0.0216 (8)	0.0210 (8)	0.0019 (6)	0.0043 (6)	0.0020 (7)
O4	0.0162 (8)	0.0266 (9)	0.0208 (8)	0.0010 (7)	0.0069 (7)	0.0035 (7)
N1	0.0142 (9)	0.0166 (9)	0.0157 (9)	0.0013 (7)	0.0039 (7)	0.0001 (8)
N2	0.0194 (10)	0.0179 (10)	0.0228 (10)	0.0010 (8)	0.0087 (8)	0.0028 (9)
C1	0.0137 (10)	0.0168 (11)	0.0166 (10)	-0.0014 (8)	0.0039 (8)	-0.0031 (9)
C2	0.0143 (10)	0.0162 (11)	0.0208 (11)	0.0007 (8)	0.0041 (8)	-0.0019 (9)
C3	0.0169 (10)	0.0152 (11)	0.0197 (11)	-0.0008 (9)	0.0047 (8)	-0.0017 (9)
C4	0.0170 (11)	0.0221 (13)	0.0218 (11)	-0.0026 (9)	0.0086 (9)	-0.0004 (10)
C5	0.0121 (10)	0.0234 (12)	0.0250 (11)	-0.0021 (9)	0.0037 (9)	-0.0021 (10)
C6	0.0146 (10)	0.0225 (12)	0.0191 (11)	-0.0006 (9)	0.0012 (8)	0.0003 (10)
C7	0.0159 (10)	0.0144 (10)	0.0169 (10)	-0.0006 (8)	0.0006 (8)	-0.0024 (9)
C8	0.0136 (10)	0.0182 (11)	0.0149 (10)	0.0002 (9)	0.0028 (8)	0.0010 (9)
C9	0.0122 (10)	0.0162 (11)	0.0203 (11)	0.0004 (8)	0.0022 (8)	0.0021 (9)
C10	0.0150 (10)	0.0200 (12)	0.0165 (10)	0.0012 (9)	0.0040 (8)	-0.0001 (9)
C11	0.0166 (10)	0.0182 (11)	0.0170 (10)	-0.0010 (9)	0.0032 (8)	-0.0026 (9)
C12	0.0150 (10)	0.0177 (11)	0.0152 (10)	-0.0004 (9)	0.0043 (8)	-0.0036 (9)
C13	0.0195 (11)	0.0182 (12)	0.0171 (10)	0.0007 (9)	0.0038 (8)	-0.0011 (9)
C14	0.0181 (11)	0.0202 (12)	0.0230 (11)	-0.0031 (9)	0.0080 (9)	-0.0011 (10)
C15	0.0166 (11)	0.0228 (13)	0.0300 (13)	-0.0003 (9)	0.0074 (9)	-0.0049 (11)
C16	0.0188 (11)	0.0161 (11)	0.0272 (12)	0.0033 (8)	0.0054 (9)	-0.0002 (10)
C17	0.0182 (11)	0.0168 (11)	0.0189 (11)	-0.0017 (9)	0.0065 (8)	-0.0021 (9)
C18	0.0166 (11)	0.0207 (12)	0.0200 (11)	-0.0007 (9)	0.0048 (9)	-0.0036 (10)

Geometric parameters (Å, °)

O1—C9	1.218 (3)	C6—H6	0.9500
O2—C9	1.301 (3)	C7—H7	0.9500
O2—H1O2	0.89 (3)	C8—C9	1.525 (3)
O3—C2	1.318 (3)	C8—C10	1.530 (3)
O4—C3	1.361 (3)	C8—H8	1.0000
O4—H1O4	0.89 (3)	C10—C11	1.501 (3)
N1—C7	1.295 (3)	C10—H10A	0.9900
N1—C8	1.465 (3)	C10—H10B	0.9900
N1—H1N1	0.97 (3)	C11—C18	1.367 (4)
N2—C18	1.375 (3)	C11—C12	1.442 (3)
N2—C17	1.378 (3)	C12—C13	1.405 (3)
N2—H1N2	0.90 (3)	C12—C17	1.409 (3)
C1—C2	1.417 (3)	C13—C14	1.387 (3)
C1—C6	1.419 (3)	C13—H13	0.9500
C1—C7	1.426 (3)	C14—C15	1.403 (4)
C2—C3	1.429 (3)	C14—H14	0.9500
C3—C4	1.384 (3)	C15—C16	1.382 (4)
C4—C5	1.395 (3)	C15—H15	0.9500
C4—H4	0.9500	C16—C17	1.396 (3)
C5—C6	1.373 (3)	C16—H16	0.9500
C5—H5	0.9500	C18—H18	0.9500
C9—O2—H1O2	110 (2)	C10—C8—H8	107.0
C3—O4—H1O4	115 (2)	O1—C9—O2	124.9 (2)
C7—N1—C8	123.31 (19)	O1—C9—C8	119.9 (2)
C7—N1—H1N1	115.0 (18)	O2—C9—C8	115.22 (19)
C8—N1—H1N1	121.5 (18)	C11—C10—C8	114.84 (18)
C18—N2—C17	108.2 (2)	C11—C10—H10A	108.6
C18—N2—H1N2	126.3 (17)	C8—C10—H10A	108.6
C17—N2—H1N2	125.5 (17)	C11—C10—H10B	108.6
C2—C1—C6	121.6 (2)	C8—C10—H10B	108.6
C2—C1—C7	121.40 (19)	H10A—C10—H10B	107.5
C6—C1—C7	117.0 (2)	C18—C11—C12	106.1 (2)
O3—C2—C1	122.1 (2)	C18—C11—C10	129.5 (2)
O3—C2—C3	121.2 (2)	C12—C11—C10	124.3 (2)
C1—C2—C3	116.62 (19)	C13—C12—C17	119.6 (2)
O4—C3—C4	118.2 (2)	C13—C12—C11	133.6 (2)
O4—C3—C2	121.44 (19)	C17—C12—C11	106.9 (2)
C4—C3—C2	120.3 (2)	C14—C13—C12	118.5 (2)
C3—C4—C5	122.3 (2)	C14—C13—H13	120.8
C3—C4—H4	118.9	C12—C13—H13	120.8
C5—C4—H4	118.9	C13—C14—C15	121.1 (2)
C6—C5—C4	119.1 (2)	C13—C14—H14	119.5
C6—C5—H5	120.4	C15—C14—H14	119.5
C4—C5—H5	120.4	C16—C15—C14	121.4 (2)
C5—C6—C1	120.1 (2)	C16—C15—H15	119.3
C5—C6—H6	119.9	C14—C15—H15	119.3

supplementary materials

C1—C6—H6	119.9	C15—C16—C17	117.7 (2)
N1—C7—C1	123.8 (2)	C15—C16—H16	121.2
N1—C7—H7	118.1	C17—C16—H16	121.2
C1—C7—H7	118.1	N2—C17—C16	130.1 (2)
N1—C8—C9	111.15 (18)	N2—C17—C12	108.1 (2)
N1—C8—C10	111.87 (19)	C16—C17—C12	121.8 (2)
C9—C8—C10	112.33 (18)	C11—C18—N2	110.8 (2)
N1—C8—H8	107.0	C11—C18—H18	124.6
C9—C8—H8	107.0	N2—C18—H18	124.6
C6—C1—C2—O3	176.6 (2)	C9—C8—C10—C11	-61.5 (3)
C7—C1—C2—O3	-2.4 (3)	C8—C10—C11—C18	-75.8 (3)
C6—C1—C2—C3	0.5 (3)	C8—C10—C11—C12	108.5 (3)
C7—C1—C2—C3	-178.4 (2)	C18—C11—C12—C13	-179.4 (2)
O3—C2—C3—O4	0.3 (3)	C10—C11—C12—C13	-2.8 (4)
C1—C2—C3—O4	176.4 (2)	C18—C11—C12—C17	-0.2 (2)
O3—C2—C3—C4	-177.1 (2)	C10—C11—C12—C17	176.4 (2)
C1—C2—C3—C4	-1.0 (3)	C17—C12—C13—C14	-0.5 (3)
O4—C3—C4—C5	-177.2 (2)	C11—C12—C13—C14	178.6 (2)
C2—C3—C4—C5	0.3 (4)	C12—C13—C14—C15	-0.3 (3)
C3—C4—C5—C6	0.9 (4)	C13—C14—C15—C16	1.3 (4)
C4—C5—C6—C1	-1.3 (4)	C14—C15—C16—C17	-1.3 (4)
C2—C1—C6—C5	0.6 (4)	C18—N2—C17—C16	178.8 (2)
C7—C1—C6—C5	179.6 (2)	C18—N2—C17—C12	-0.3 (3)
C8—N1—C7—C1	-179.0 (2)	C15—C16—C17—N2	-178.5 (2)
C2—C1—C7—N1	-0.7 (4)	C15—C16—C17—C12	0.5 (4)
C6—C1—C7—N1	-179.7 (2)	C13—C12—C17—N2	179.6 (2)
C7—N1—C8—C9	-132.3 (2)	C11—C12—C17—N2	0.3 (2)
C7—N1—C8—C10	101.2 (2)	C13—C12—C17—C16	0.4 (3)
N1—C8—C9—O1	-170.5 (2)	C11—C12—C17—C16	-178.9 (2)
C10—C8—C9—O1	-44.3 (3)	C12—C11—C18—N2	0.0 (3)
N1—C8—C9—O2	9.7 (3)	C10—C11—C18—N2	-176.3 (2)
C10—C8—C9—O2	135.9 (2)	C17—N2—C18—C11	0.2 (3)
N1—C8—C10—C11	64.4 (2)		

Hydrogen-bond geometry (\AA , $^\circ$)

$D-H\cdots A$	$D-H$	$H\cdots A$	$D\cdots A$	$D-H\cdots A$
N1—H1N1 \cdots O3	0.97 (3)	1.86 (3)	2.646 (2)	136 (2)
N2—H1N2 \cdots O3 ⁱ	0.90 (3)	2.13 (3)	3.020 (2)	170 (2)
O2—H1O2 \cdots O3 ⁱⁱ	0.89 (3)	1.64 (4)	2.526 (2)	174 (4)
O4—H1O4 \cdots O3	0.89 (3)	2.45 (3)	2.823 (2)	106 (2)
O4—H1O4 \cdots O1 ⁱ	0.89 (3)	1.83 (3)	2.665 (2)	156 (3)
C7—H7 \cdots O4 ⁱⁱⁱ	0.95	2.49	3.197 (3)	132

Symmetry codes: (i) $-x, y+1/2, -z$; (ii) $-x, y-1/2, -z$; (iii) $-x+1, y-1/2, -z$.

Fig. 1




Fig. 2

