

Phenyl 3-methoxy-4-phenoxybenzoate

Jing Zhou, Shuai Zheng, Gang Chen, Wenjing Wang and
 Zhenxing Du*

College of Science, Northwest A&F University, Yangling, Shaanxi 712100, People's Republic of China

Correspondence e-mail: duz@nwsuaf.edu.cn


Received 8 July 2011; accepted 18 August 2011

Key indicators: single-crystal X-ray study; $T = 298\text{ K}$; mean $\sigma(\text{C}-\text{C}) = 0.003\text{ \AA}$; R factor = 0.041; wR factor = 0.118; data-to-parameter ratio = 13.1.

In the title molecule, $\text{C}_{20}\text{H}_{16}\text{O}_4$, the two outermost phenyl rings form dihedral angles of $79.80(7)$ and $69.35(7)^\circ$ with the central benzene ring. In the crystal structure, weak intermolecular $\text{C}-\text{H}\cdots\text{O}$ interactions link the molecules into ribbons propagating along $[1\bar{1}0]$.

Related literature

For the general synthesis of derivatives of diphenylethers, see: Paul & Gupta (2004). For related structures, see: Chen *et al.* (2006); Petek *et al.* (2005); Chantrapromma *et al.* (2001); Nakamura *et al.* (1983); Gopal *et al.* (1980). For applications of diphenylether derivatives, see: Dey & Desiraju (2005); Wang *et al.* (2005).



Experimental

Crystal data

$\text{C}_{20}\text{H}_{16}\text{O}_4$	$V = 3261.5(5)\text{ \AA}^3$
$M_r = 320.33$	$Z = 8$
Monoclinic, $C2/c$	Mo $K\alpha$ radiation
$a = 11.0261(10)\text{ \AA}$	$\mu = 0.09\text{ mm}^{-1}$
$b = 11.9624(11)\text{ \AA}$	$T = 298\text{ K}$
$c = 24.961(2)\text{ \AA}$	$0.49 \times 0.42 \times 0.40\text{ mm}$
$\beta = 97.842(1)^\circ$	

Data collection

Bruker SMART APEXII CCD area-detector diffractometer	7972 measured reflections
Absorption correction: multi-scan (<i>SADABS</i> ; Bruker, 2009)	2878 independent reflections
$T_{\min} = 0.957$, $T_{\max} = 0.965$	1785 reflections with $I > 2\sigma(I)$
	$R_{\text{int}} = 0.040$

Refinement

$R[F^2 > 2\sigma(F^2)] = 0.041$	219 parameters
$wR(F^2) = 0.118$	H-atom parameters constrained
$S = 1.05$	$\Delta\rho_{\max} = 0.16\text{ e \AA}^{-3}$
2878 reflections	$\Delta\rho_{\min} = -0.17\text{ e \AA}^{-3}$

Table 1
 Hydrogen-bond geometry (\AA , $^\circ$).

$D-\text{H}\cdots A$	$D-\text{H}$	$\text{H}\cdots A$	$D\cdots A$	$D-\text{H}\cdots A$
C8—H8C···O2 ⁱ	0.96	2.52	3.408 (4)	153
C14—H14···O2 ⁱⁱ	0.93	2.53	3.446 (3)	167
C20—H20···O3 ⁱⁱ	0.93	2.60	3.468 (3)	155

Symmetry codes: (i) $-x + \frac{1}{2}, -y + \frac{3}{2}, -z + 1$; (ii) $-x + 1, -y + 1, -z + 1$.

Data collection: *APEX2* (Bruker, 2009); cell refinement: *SAINT* (Bruker, 2009); data reduction: *SAINT*; program(s) used to solve structure: *SHELXS97* (Sheldrick, 2008); program(s) used to refine structure: *SHELXL97* (Sheldrick, 2008); molecular graphics: *SHELXTL* (Sheldrick, 2008); software used to prepare material for publication: *SHELXTL*.

Financial support from the Fundamental Research Funds for the Central Universities in NWSUAF (grant No. QN2009048) and from the National Natural Science Foundation of China (grant No. 20802058) is greatly appreciated.

Supplementary data and figures for this paper are available from the IUCr electronic archives (Reference: CV5133).

References

- Bruker (2009). *APEX2, SAINT* and *SADABS*. Bruker AXS Inc., Madison, Wisconsin, USA.
- Chantrapromma, S., Fun, H.-K., Razak, I. A., Saewon, N., Karalai, C., Ponglimanont, C. & Chantrapromma, K. (2001). *Acta Cryst. E57*, o1047–o1049.
- Chen, Z.-B., Wu, J., Zhang, P.-Z. & Zhang, P.-M. (2006). *Acta Cryst. E62*, o1336–o1337.
- Dey, A. & Desiraju, G. R. (2005). *Chem. Commun.*, pp. 2486–2488.
- Gopal, R., Chandler, W. D. & Robertson, B. E. (1980). *Can. J. Chem. 58*, 658–663.
- Nakamura, M., Fukuyama, K., Tsukihara, T., Katube, Y. & Hamasaki, T. (1983). *Acta Cryst. C39*, 268–270.
- Paul, S. & Gupta, M. (2004). *Tetrahedron Lett. 45*, 8825–8829.
- Petek, H., Akdemir, N., Kantar, C., Ağar, E. & Şenel, İ. (2005). *Acta Cryst. E61*, o1233–o1234.
- Sheldrick, G. M. (2008). *Acta Cryst. A64*, 112–122.
- Wang, X. L., Qin, C., Wang, E. B., Li, Y. G., Su, Z. M., Xu, L. & Carlucci, L. (2005). *Angew. Chem. Int. Ed. Engl. 44*, 5824–5827.

supplementary materials

Acta Cryst. (2011). E67, o2393 [doi:10.1107/S1600536811033678]

Phenyl 3-methoxy-4-phenoxybenzoate

J. Zhou, S. Zheng, G. Chen, W. Wang and Z. Du

Comment

4,4'-Dicarboxy diphenyl ether was widely used in polymer frameworks which show potential applications in microelectronics, nonlinear optics, porous materials and catalysis (Dey & Desiraju, 2005; Wang *et al.*, 2005). The title compound (**I**) was obtained as unexpected product in the cyclization of diazonium salt to dibenzofuran. Accidentally, in our pursuing new methodology of synthesis of dibenzofuran the diazonium salt was heated without any catalyst. In order to determine the structure of a new compound, it was characterized by single-crystal X-ray analysis.

In (**I**) (Fig. 1) all bond lengths and angles are normal and correspond to those observed in the related compounds (Chen *et al.*, 2006; Petek *et al.*, 2005; Chantrapromma *et al.*, 2001; Nakamura *et al.*, 1983; Gopal *et al.*, 1980). Two utmost phenyl rings form the dihedral angles of 79.80 (7) and 69.35 (7) $^{\circ}$, respectively, with the central benzene ring. In the crystal structure, weak intermolecular C—H \cdots O interactions (Table 1) link molecules into ribbons propagated in [1 $\bar{1}$ 0].

Experimental

A solution of 2-(2-methoxy-4-(phenoxy carbonyl)phenoxy)benzenediazonium tetrafluoroborate (0.868 g, 2 mmol) in ethanol (30 ml) was refluxed for 1 h. Then the solvent was evaporated and the residue was purified through column chromatography. The compound was dissolved in hexane, and white square crystals were obtained by slow evaporation of the solvent over one week.

Figures




Fig. 1. The molecular structure of (**I**). Displacement ellipsoids are drawn at the 30% probability level.

Phenyl 3-methoxy-4-phenoxybenzoate

Crystal data

C ₂₀ H ₁₆ O ₄	D _x = 1.305 Mg m ⁻³
M _r = 320.33	Melting point: 383 K
Monoclinic, C2/c	Mo K α radiation, λ = 0.71073 Å
<i>a</i> = 11.0261 (10) Å	Cell parameters from 2372 reflections
<i>b</i> = 11.9624 (11) Å	θ = 2.5–27.3 $^{\circ}$
<i>c</i> = 24.961 (2) Å	μ = 0.09 mm ⁻¹
β = 97.842 (1) $^{\circ}$	T = 298 K
<i>V</i> = 3261.5 (5) Å ³	Block, white

supplementary materials

$Z = 8$ $0.49 \times 0.42 \times 0.40$ mm
 $F(000) = 1344$

Data collection

Bruker SMART APEXII CCD area-detector diffractometer 2878 independent reflections
Radiation source: fine-focus sealed tube 1785 reflections with $I > 2\sigma(I)$
graphite $R_{\text{int}} = 0.040$
 φ and ω scans $\theta_{\text{max}} = 25.0^\circ$, $\theta_{\text{min}} = 2.5^\circ$
Absorption correction: multi-scan (*SADABS*; Bruker, 2009) $h = -12 \rightarrow 13$
 $T_{\text{min}} = 0.957$, $T_{\text{max}} = 0.965$ $k = -14 \rightarrow 8$
7972 measured reflections $l = -29 \rightarrow 29$

Refinement

Refinement on F^2 Secondary atom site location: difference Fourier map
Least-squares matrix: full Hydrogen site location: inferred from neighbouring sites
 $R[F^2 > 2\sigma(F^2)] = 0.041$ H-atom parameters constrained
 $wR(F^2) = 0.118$ $w = 1/[\sigma^2(F_o^2) + (0.0413P)^2 + 1.7982P]$
 $S = 1.05$ where $P = (F_o^2 + 2F_c^2)/3$
2878 reflections $(\Delta/\sigma)_{\text{max}} < 0.001$
219 parameters $\Delta\rho_{\text{max}} = 0.16 \text{ e } \text{\AA}^{-3}$
0 restraints $\Delta\rho_{\text{min}} = -0.17 \text{ e } \text{\AA}^{-3}$
Extinction correction: *SHELXL97* (Sheldrick, 2008),
 $F_c^* = kFc[1 + 0.001xFc^2\lambda^3/\sin(2\theta)]^{-1/4}$
Primary atom site location: structure-invariant direct Extinction coefficient: 0.0053 (4)
methods

Special details

Geometry. All e.s.d.'s (except the e.s.d. in the dihedral angle between two l.s. planes) are estimated using the full covariance matrix. The cell e.s.d.'s are taken into account individually in the estimation of e.s.d.'s in distances, angles and torsion angles; correlations between e.s.d.'s in cell parameters are only used when they are defined by crystal symmetry. An approximate (isotropic) treatment of cell e.s.d.'s is used for estimating e.s.d.'s involving l.s. planes.

Refinement. Refinement of F^2 against ALL reflections. The weighted R -factor wR and goodness of fit S are based on F^2 , conventional R -factors R are based on F , with F set to zero for negative F^2 . The threshold expression of $F^2 > \sigma(F^2)$ is used only for calculating R -factors(gt) etc. and is not relevant to the choice of reflections for refinement. R -factors based on F^2 are statistically about twice as large as those based on F , and R -factors based on ALL data will be even larger.

Fractional atomic coordinates and isotropic or equivalent isotropic displacement parameters (\AA^2)

	x	y	z	$U_{\text{iso}}^*/U_{\text{eq}}$
O1	0.20235 (14)	0.40188 (12)	0.47155 (6)	0.0542 (5)
O2	0.22859 (15)	0.58572 (13)	0.46169 (6)	0.0617 (5)

O3	0.55526 (13)	0.68666 (12)	0.61835 (6)	0.0510 (4)
O4	0.59538 (14)	0.50787 (12)	0.67467 (5)	0.0562 (5)
C1	0.25750 (19)	0.50110 (18)	0.48580 (8)	0.0416 (5)
C2	0.35069 (17)	0.49359 (16)	0.53388 (7)	0.0365 (5)
C3	0.40974 (17)	0.59252 (17)	0.55134 (7)	0.0383 (5)
H3	0.3923	0.6579	0.5316	0.046*
C4	0.49363 (17)	0.59433 (16)	0.59748 (8)	0.0362 (5)
C5	0.51873 (18)	0.49544 (17)	0.62679 (7)	0.0393 (5)
C6	0.4624 (2)	0.39727 (18)	0.60920 (8)	0.0484 (6)
H6	0.4809	0.3316	0.6285	0.058*
C7	0.37782 (19)	0.39605 (18)	0.56250 (8)	0.0454 (6)
H7	0.3396	0.3296	0.5506	0.054*
C8	0.5401 (3)	0.7870 (2)	0.58760 (11)	0.0768 (9)
H8A	0.5629	0.7742	0.5524	0.115*
H8B	0.5911	0.8444	0.6056	0.115*
H8C	0.4560	0.8101	0.5841	0.115*
C9	0.6653 (2)	0.41886 (18)	0.69837 (8)	0.0459 (6)
C10	0.6668 (2)	0.4066 (2)	0.75311 (9)	0.0601 (7)
H10	0.6174	0.4510	0.7718	0.072*
C11	0.7429 (3)	0.3271 (3)	0.77982 (10)	0.0760 (9)
H11	0.7440	0.3173	0.8169	0.091*
C12	0.8166 (2)	0.2624 (2)	0.75287 (12)	0.0739 (9)
H12	0.8679	0.2094	0.7714	0.089*
C13	0.8145 (2)	0.2764 (2)	0.69818 (11)	0.0663 (7)
H13	0.8648	0.2329	0.6796	0.080*
C14	0.7378 (2)	0.3548 (2)	0.67055 (9)	0.0566 (7)
H14	0.7357	0.3638	0.6334	0.068*
C15	0.1108 (2)	0.40450 (17)	0.42591 (8)	0.0436 (6)
C16	-0.0096 (2)	0.40588 (19)	0.43430 (9)	0.0548 (6)
H16	-0.0307	0.4064	0.4691	0.066*
C17	-0.0987 (2)	0.4065 (2)	0.38964 (10)	0.0623 (7)
H17	-0.1809	0.4077	0.3944	0.075*
C18	-0.0671 (2)	0.4055 (2)	0.33860 (10)	0.0606 (7)
H18	-0.1276	0.4068	0.3088	0.073*
C19	0.0534 (2)	0.4024 (2)	0.33141 (9)	0.0620 (7)
H19	0.0747	0.4013	0.2966	0.074*
C20	0.1434 (2)	0.40105 (19)	0.37542 (9)	0.0539 (6)
H20	0.2255	0.3978	0.3706	0.065*

Atomic displacement parameters (\AA^2)

	U^{11}	U^{22}	U^{33}	U^{12}	U^{13}	U^{23}
O1	0.0618 (10)	0.0448 (9)	0.0481 (9)	-0.0063 (8)	-0.0211 (8)	0.0062 (7)
O2	0.0733 (11)	0.0466 (10)	0.0555 (10)	-0.0032 (8)	-0.0256 (8)	0.0097 (8)
O3	0.0550 (9)	0.0386 (9)	0.0528 (9)	-0.0041 (7)	-0.0163 (7)	0.0042 (7)
O4	0.0706 (11)	0.0496 (10)	0.0405 (8)	0.0025 (8)	-0.0208 (8)	0.0034 (7)
C1	0.0434 (12)	0.0420 (13)	0.0374 (12)	-0.0009 (11)	-0.0016 (10)	0.0005 (10)
C2	0.0368 (11)	0.0383 (12)	0.0328 (10)	0.0004 (9)	-0.0008 (9)	0.0020 (9)

supplementary materials

C3	0.0383 (11)	0.0392 (12)	0.0361 (11)	0.0030 (10)	0.0000 (9)	0.0093 (9)
C4	0.0348 (11)	0.0366 (12)	0.0359 (11)	0.0007 (9)	-0.0001 (9)	-0.0001 (9)
C5	0.0412 (12)	0.0430 (13)	0.0309 (10)	0.0019 (10)	-0.0053 (9)	0.0030 (9)
C6	0.0592 (14)	0.0394 (13)	0.0423 (12)	0.0007 (11)	-0.0086 (11)	0.0114 (10)
C7	0.0496 (13)	0.0397 (13)	0.0434 (12)	-0.0039 (10)	-0.0061 (10)	0.0018 (10)
C8	0.0853 (19)	0.0437 (15)	0.089 (2)	-0.0156 (14)	-0.0321 (16)	0.0169 (14)
C9	0.0454 (13)	0.0466 (13)	0.0403 (12)	-0.0045 (11)	-0.0131 (10)	0.0126 (11)
C10	0.0530 (14)	0.0817 (18)	0.0419 (13)	-0.0012 (13)	-0.0065 (11)	0.0126 (13)
C11	0.0707 (19)	0.104 (2)	0.0473 (15)	-0.0006 (18)	-0.0142 (14)	0.0313 (16)
C12	0.0597 (17)	0.078 (2)	0.0744 (19)	0.0020 (15)	-0.0244 (15)	0.0286 (16)
C13	0.0544 (15)	0.0668 (17)	0.0740 (18)	0.0055 (13)	-0.0047 (13)	0.0088 (14)
C14	0.0622 (16)	0.0592 (15)	0.0452 (13)	-0.0011 (13)	-0.0039 (12)	0.0090 (12)
C15	0.0481 (14)	0.0376 (12)	0.0405 (12)	-0.0027 (10)	-0.0108 (10)	0.0020 (10)
C16	0.0587 (16)	0.0583 (15)	0.0466 (13)	-0.0017 (12)	0.0044 (11)	-0.0078 (11)
C17	0.0422 (13)	0.0724 (18)	0.0694 (17)	0.0012 (12)	-0.0025 (12)	-0.0132 (14)
C18	0.0569 (16)	0.0643 (17)	0.0534 (15)	-0.0035 (13)	-0.0183 (12)	-0.0025 (12)
C19	0.0620 (17)	0.0809 (19)	0.0408 (13)	-0.0123 (14)	-0.0015 (11)	0.0036 (12)
C20	0.0456 (13)	0.0633 (16)	0.0514 (14)	-0.0077 (12)	0.0010 (11)	0.0023 (12)

Geometric parameters (\AA , $^\circ$)

O1—C1	1.359 (2)	C9—C10	1.372 (3)
O1—C15	1.415 (2)	C10—C11	1.379 (3)
O2—C1	1.198 (2)	C10—H10	0.9300
O3—C4	1.362 (2)	C11—C12	1.364 (4)
O3—C8	1.422 (3)	C11—H11	0.9300
O4—C5	1.375 (2)	C12—C13	1.372 (4)
O4—C9	1.398 (2)	C12—H12	0.9300
C1—C2	1.472 (3)	C13—C14	1.383 (3)
C2—C7	1.379 (3)	C13—H13	0.9300
C2—C3	1.392 (3)	C14—H14	0.9300
C3—C4	1.375 (3)	C15—C20	1.358 (3)
C3—H3	0.9300	C15—C16	1.372 (3)
C4—C5	1.399 (3)	C16—C17	1.382 (3)
C5—C6	1.372 (3)	C16—H16	0.9300
C6—C7	1.390 (3)	C17—C18	1.366 (3)
C6—H6	0.9300	C17—H17	0.9300
C7—H7	0.9300	C18—C19	1.365 (3)
C8—H8A	0.9600	C18—H18	0.9300
C8—H8B	0.9600	C19—C20	1.377 (3)
C8—H8C	0.9600	C19—H19	0.9300
C9—C14	1.363 (3)	C20—H20	0.9300
C1—O1—C15	115.84 (15)	C9—C10—C11	118.6 (3)
C4—O3—C8	117.53 (15)	C9—C10—H10	120.7
C5—O4—C9	121.60 (16)	C11—C10—H10	120.7
O2—C1—O1	121.84 (18)	C12—C11—C10	121.1 (2)
O2—C1—C2	124.68 (19)	C12—C11—H11	119.4
O1—C1—C2	113.44 (18)	C10—C11—H11	119.4
C7—C2—C3	119.86 (17)	C11—C12—C13	119.4 (2)

C7—C2—C1	123.48 (19)	C11—C12—H12	120.3
C3—C2—C1	116.62 (17)	C13—C12—H12	120.3
C4—C3—C2	120.53 (18)	C12—C13—C14	120.4 (3)
C4—C3—H3	119.7	C12—C13—H13	119.8
C2—C3—H3	119.7	C14—C13—H13	119.8
O3—C4—C3	125.18 (17)	C9—C14—C13	119.2 (2)
O3—C4—C5	115.66 (16)	C9—C14—H14	120.4
C3—C4—C5	119.15 (18)	C13—C14—H14	120.4
C6—C5—O4	124.64 (18)	C20—C15—C16	121.75 (19)
C6—C5—C4	120.53 (17)	C20—C15—O1	119.8 (2)
O4—C5—C4	114.69 (17)	C16—C15—O1	118.42 (19)
C5—C6—C7	119.99 (19)	C15—C16—C17	118.3 (2)
C5—C6—H6	120.0	C15—C16—H16	120.9
C7—C6—H6	120.0	C17—C16—H16	120.9
C2—C7—C6	119.91 (19)	C18—C17—C16	120.6 (2)
C2—C7—H7	120.0	C18—C17—H17	119.7
C6—C7—H7	120.0	C16—C17—H17	119.7
O3—C8—H8A	109.5	C19—C18—C17	120.0 (2)
O3—C8—H8B	109.5	C19—C18—H18	120.0
H8A—C8—H8B	109.5	C17—C18—H18	120.0
O3—C8—H8C	109.5	C18—C19—C20	120.3 (2)
H8A—C8—H8C	109.5	C18—C19—H19	119.9
H8B—C8—H8C	109.5	C20—C19—H19	119.9
C14—C9—C10	121.3 (2)	C15—C20—C19	119.1 (2)
C14—C9—O4	122.70 (19)	C15—C20—H20	120.4
C10—C9—O4	115.7 (2)	C19—C20—H20	120.4
C15—O1—C1—O2	1.3 (3)	C5—C6—C7—C2	0.1 (3)
C15—O1—C1—C2	179.29 (18)	C5—O4—C9—C14	50.8 (3)
O2—C1—C2—C7	176.6 (2)	C5—O4—C9—C10	-135.7 (2)
O1—C1—C2—C7	-1.3 (3)	C14—C9—C10—C11	-0.4 (3)
O2—C1—C2—C3	-1.1 (3)	O4—C9—C10—C11	-174.1 (2)
O1—C1—C2—C3	-179.05 (17)	C9—C10—C11—C12	0.7 (4)
C7—C2—C3—C4	-1.1 (3)	C10—C11—C12—C13	-0.4 (4)
C1—C2—C3—C4	176.72 (19)	C11—C12—C13—C14	-0.3 (4)
C8—O3—C4—C3	-5.9 (3)	C10—C9—C14—C13	-0.3 (3)
C8—O3—C4—C5	175.0 (2)	O4—C9—C14—C13	172.91 (19)
C2—C3—C4—O3	-179.20 (19)	C12—C13—C14—C9	0.6 (4)
C2—C3—C4—C5	-0.2 (3)	C1—O1—C15—C20	81.7 (2)
C9—O4—C5—C6	28.7 (3)	C1—O1—C15—C16	-100.9 (2)
C9—O4—C5—C4	-155.67 (19)	C20—C15—C16—C17	-1.6 (3)
O3—C4—C5—C6	-179.50 (19)	O1—C15—C16—C17	-178.9 (2)
C3—C4—C5—C6	1.4 (3)	C15—C16—C17—C18	0.2 (4)
O3—C4—C5—O4	4.6 (3)	C16—C17—C18—C19	0.8 (4)
C3—C4—C5—O4	-174.49 (18)	C17—C18—C19—C20	-0.4 (4)
O4—C5—C6—C7	174.1 (2)	C16—C15—C20—C19	2.0 (3)
C4—C5—C6—C7	-1.3 (3)	O1—C15—C20—C19	179.3 (2)
C3—C2—C7—C6	1.1 (3)	C18—C19—C20—C15	-1.0 (4)
C1—C2—C7—C6	-176.5 (2)		

supplementary materials

Hydrogen-bond geometry (Å, °)

<i>D—H···A</i>	<i>D—H</i>	<i>H···A</i>	<i>D···A</i>	<i>D—H···A</i>
C8—H8C···O2 ⁱ	0.96	2.52	3.408 (4)	153
C14—H14···O2 ⁱⁱ	0.93	2.53	3.446 (3)	167
C20—H20···O3 ⁱⁱ	0.93	2.60	3.468 (3)	155

Symmetry codes: (i) $-x+1/2, -y+3/2, -z+1$; (ii) $-x+1, -y+1, -z+1$.

Fig. 1

