

Monoclinic, $P2_1/c$
 $a = 13.6794 (4)$ Å
 $b = 13.4304 (4)$ Å
 $c = 7.3678 (2)$ Å
 $\beta = 91.055 (3)^\circ$
 $V = 1353.38 (7)$ Å³

$Z = 4$
Cu $K\alpha$ radiation
 $\mu = 2.29$ mm⁻¹
 $T = 100$ K
 $0.30 \times 0.05 \times 0.05$ mm

4-(5-Oxo-3-phenyl-4,5-dihydro-1*H*-pyrazol-1-yl)benzenesulfonamide

Abdulrahman O. Al-Youbi,^a Abdullah M. Asiri,^{a,b}
Hassan M. Faidallah,^a Khalid A. Alamry^a and Seik Weng
Ng^{c,a*}

^aChemistry Department, Faculty of Science, King Abdulaziz University, PO Box 80203 Jeddah, Saudi Arabia, ^bCenter of Excellence for Advanced Materials Research, King Abdulaziz University, PO Box 80203 Jeddah, Saudi Arabia, and ^cDepartment of Chemistry, University of Malaya, 50603 Kuala Lumpur, Malaysia
Correspondence e-mail: seikweng@um.edu.my


Received 11 August 2011; accepted 13 August 2011

Key indicators: single-crystal X-ray study; $T = 100$ K; mean $\sigma(C-C) = 0.003$ Å;
 R factor = 0.041; wR factor = 0.121; data-to-parameter ratio = 13.2.

With respect to the aliphatic planar five-membered ring (r.m.s. deviation = 0.011 Å) of the title compound, C₁₅H₁₃N₃O₃S, the phenyl ring is aligned at 6.9 (1)° and the phenylene ring at 2.4 (1)°, so that the three rings are nearly coplanar. The amino group has the N atom in a pyramidal geometry; the group is a hydrogen-bond donor to the sulfonyl O atom of one molecule and to the ketonic O atom of another molecule, resulting in the formation of a layer parallel to the *bc* plane.

Related literature

For the synthesis, see: Casoni (1956); Itano (1955).



Experimental

Crystal data

C₁₅H₁₃N₃O₃S

$M_r = 315.34$

Data collection

Agilent SuperNova Dual diffractometer with an Atlas detector
Absorption correction: multi-scan (*CrysAlis PRO*; Agilent, 2010)
 $T_{\min} = 0.546$, $T_{\max} = 0.894$

10404 measured reflections
2731 independent reflections
2444 reflections with $I > 2\sigma(I)$
 $R_{\text{int}} = 0.041$

Refinement

$R[F^2 > 2\sigma(F^2)] = 0.041$
 $wR(F^2) = 0.121$
 $S = 1.06$
2731 reflections
207 parameters
2 restraints

H atoms treated by a mixture of independent and constrained refinement
 $\Delta\rho_{\max} = 0.73$ e Å⁻³
 $\Delta\rho_{\min} = -0.51$ e Å⁻³

Table 1
Hydrogen-bond geometry (Å, °).

D–H···A	D–H	H···A	D···A	D–H···A
N3–H1···O3 ⁱ	0.88 (1)	2.12 (1)	2.975 (2)	164 (2)
N3–H2···O2 ⁱⁱ	0.87 (1)	2.12 (1)	2.978 (2)	168 (2)

Symmetry codes: (i) $-x + 2$, $-y + 1$, $-z + 1$; (ii) x , $-y + \frac{3}{2}$, $z + \frac{1}{2}$.

Data collection: *CrysAlis PRO* (Agilent, 2010); cell refinement: *CrysAlis PRO*; data reduction: *CrysAlis PRO*; program(s) used to solve structure: *SHELXS97* (Sheldrick, 2008); program(s) used to refine structure: *SHELXL97* (Sheldrick, 2008); molecular graphics: *X-SEED* (Barbour, 2001); software used to prepare material for publication: *publCIF* (Westrip, 2010).

We thank King Abdulaziz University and the University of Malaya for supporting this study.

Supplementary data and figures for this paper are available from the IUCr electronic archives (Reference: BT5611).

References

- Agilent (2010). *CrysAlis PRO*. Agilent Technologies, Yarnton, Oxfordshire, England.
Barbour, L. J. (2001). *J. Supramol. Chem.* **1**, 189–191.
Casoni, D. (1956). *Boll. Sci. Fac. Chim. Ind. Bologna*, **14**, 22–30.
Itano, K. (1955). *Yakugaku Zasshi*, **5**, 441–444.
Sheldrick, G. M. (2008). *Acta Cryst. A* **64**, 112–122.
Westrip, S. P. (2010). *J. Appl. Cryst.* **43**, 920–925.

supplementary materials

Acta Cryst. (2011). E67, o2428 [doi:10.1107/S1600536811032879]

4-(5-Oxo-3-phenyl-4,5-dihydro-1*H*-pyrazol-1-yl)benzenesulfonamide

A. O. Al-Youbi, A. M. Asiri, H. M. Faidallah, K. A. Alamry and S. W. Ng

Comment

We are examining the medicinal properties of phenylpyrazolones of which the 4-benzenesulfonamide derivative (Scheme I) is expected to show enhanced activity. The synthesis of the compound was reported a long time ago (Casoni, 1956; Itano, 1955). With respect to the aliphatic planar five-membered ring, the phenyl ring is aligned at 6.9 (1) $^{\circ}$ and the phenylene ring at 2.4 (1) $^{\circ}$. The amino group is hydrogen bond donor to the sulfonyl O atom of one molecule and to the ketonic O atom of another molecule to result in the formation of a layer parallel to the bc plane.

Experimental

Ethyl benzoylacetate (10 mmol) and 4-hydrazinobenzenesulfonamide hydrochloride (10 mmol) were heated in ethanol (50 ml) for 4 h; water was added to precipitate the product, which was collected and recrystallized from ethanol as brownish-orange crystals; m.p. 510–511 K.

Refinement

Carbon bound H-atoms were placed in calculated positions [C–H 0.95 to 0.98 Å, $U_{\text{iso}}(\text{H})$ 1.2–1.5 $U_{\text{eq}}(\text{C})$] and were included in the refinement in the riding model approximation.

The amino H-atoms were located in a difference Fouier map and were refined with a distance restraint of N–H 0.88±0.01 Å; their isotropic displacement parameter were freely refined.

Figures




Fig. 1. Anisotropic displacement ellipsoid plot (Barbour, 2001) of $\text{C}_{15}\text{H}_{13}\text{N}_3\text{O}_2\text{S}$ at the 70% probability level; hydrogen atoms are drawn as spheres of arbitrary radius.

4-(5-Oxo-3-phenyl-4,5-dihydro-1*H*-pyrazol-1-yl)benzenesulfonamide

Crystal data

$\text{C}_{15}\text{H}_{13}\text{N}_3\text{O}_3\text{S}$	$F(000) = 656$
$M_r = 315.34$	$D_x = 1.548 \text{ Mg m}^{-3}$
Monoclinic, $P2_1/c$	$\text{Cu } K\alpha$ radiation, $\lambda = 1.54184 \text{ \AA}$
Hall symbol: -P 2ybc	Cell parameters from 4563 reflections
$a = 13.6794 (4) \text{ \AA}$	$\theta = 3.2\text{--}74.4^\circ$

supplementary materials

$b = 13.4304(4)$ Å	$\mu = 2.29$ mm $^{-1}$
$c = 7.3678(2)$ Å	$T = 100$ K
$\beta = 91.055(3)^\circ$	Prism, brown orange
$V = 1353.38(7)$ Å 3	$0.30 \times 0.05 \times 0.05$ mm
$Z = 4$	

Data collection

Agilent SuperNova Dual diffractometer with an Atlas detector	2731 independent reflections
Radiation source: SuperNova (Cu) X-ray Source	2444 reflections with $I > 2\sigma(I)$
Mirror	$R_{\text{int}} = 0.041$
Detector resolution: 10.4041 pixels mm $^{-1}$	$\theta_{\text{max}} = 74.5^\circ, \theta_{\text{min}} = 3.2^\circ$
ω scan	$h = -17 \rightarrow 16$
Absorption correction: multi-scan (<i>CrysAlis PRO</i> ; Agilent, 2010)	$k = -16 \rightarrow 16$
$T_{\text{min}} = 0.546, T_{\text{max}} = 0.894$	$l = -5 \rightarrow 9$
10404 measured reflections	

Refinement

Refinement on F^2	Primary atom site location: structure-invariant direct methods
Least-squares matrix: full	Secondary atom site location: difference Fourier map
$R[F^2 > 2\sigma(F^2)] = 0.041$	Hydrogen site location: inferred from neighbouring sites
$wR(F^2) = 0.121$	H atoms treated by a mixture of independent and constrained refinement
$S = 1.06$	$w = 1/[\sigma^2(F_o^2) + (0.0759P)^2 + 0.5321P]$ where $P = (F_o^2 + 2F_c^2)/3$
2731 reflections	$(\Delta/\sigma)_{\text{max}} = 0.001$
207 parameters	$\Delta\rho_{\text{max}} = 0.73$ e Å $^{-3}$
2 restraints	$\Delta\rho_{\text{min}} = -0.51$ e Å $^{-3}$

Special details

Geometry. All e.s.d.'s (except the e.s.d. in the dihedral angle between two l.s. planes) are estimated using the full covariance matrix. The cell e.s.d.'s are taken into account individually in the estimation of e.s.d.'s in distances, angles and torsion angles; correlations between e.s.d.'s in cell parameters are only used when they are defined by crystal symmetry. An approximate (isotropic) treatment of cell e.s.d.'s is used for estimating e.s.d.'s involving l.s. planes.

Fractional atomic coordinates and isotropic or equivalent isotropic displacement parameters (Å 2)

	x	y	z	$U_{\text{iso}}^*/U_{\text{eq}}$
S1	0.91654(3)	0.64477(3)	0.49527(6)	0.01653(16)
O1	0.52785(10)	0.32661(11)	0.6467(2)	0.0319(4)
O2	0.90632(9)	0.74972(10)	0.46298(18)	0.0220(3)
O3	0.95114(9)	0.58212(10)	0.35168(17)	0.0209(3)

N2	0.46514 (11)	0.56244 (12)	0.8066 (2)	0.0203 (3)
N3	0.99249 (12)	0.63163 (12)	0.6642 (2)	0.0197 (3)
H1	1.0122 (17)	0.5704 (9)	0.682 (3)	0.028 (6)*
H2	0.9740 (16)	0.6620 (16)	0.763 (2)	0.023 (6)*
N1	0.52929 (11)	0.49471 (11)	0.7246 (2)	0.0197 (3)
C1	0.49011 (14)	0.39912 (14)	0.7147 (3)	0.0227 (4)
C2	0.39264 (14)	0.40640 (14)	0.8033 (3)	0.0224 (4)
H2A	0.3390	0.3864	0.7189	0.027*
H2B	0.3903	0.3645	0.9138	0.027*
C3	0.38673 (13)	0.51403 (14)	0.8491 (2)	0.0198 (4)
C4	0.30272 (13)	0.56293 (14)	0.9324 (2)	0.0205 (4)
C5	0.30792 (15)	0.66235 (15)	0.9877 (3)	0.0261 (4)
H5	0.3663	0.6993	0.9713	0.031*
C6	0.22764 (16)	0.70694 (16)	1.0664 (3)	0.0312 (5)
H6	0.2311	0.7745	1.1040	0.037*
C7	0.14192 (15)	0.65297 (16)	1.0905 (3)	0.0294 (5)
H7	0.0870	0.6837	1.1441	0.035*
C8	0.13695 (14)	0.55477 (17)	1.0363 (3)	0.0281 (5)
H8	0.0784	0.5181	1.0528	0.034*
C9	0.21695 (13)	0.50911 (15)	0.9577 (3)	0.0235 (4)
H9	0.2132	0.4413	0.9213	0.028*
C10	0.62051 (12)	0.52987 (13)	0.6675 (2)	0.0178 (4)
C11	0.68801 (13)	0.46447 (14)	0.5912 (2)	0.0197 (4)
H11	0.6720	0.3961	0.5759	0.024*
C12	0.77831 (13)	0.49990 (13)	0.5381 (2)	0.0184 (4)
H12	0.8246	0.4557	0.4875	0.022*
C13	0.80101 (13)	0.60030 (13)	0.5590 (2)	0.0175 (4)
C14	0.73339 (14)	0.66554 (14)	0.6338 (3)	0.0201 (4)
H14	0.7495	0.7339	0.6481	0.024*
C15	0.64299 (14)	0.63125 (14)	0.6873 (3)	0.0200 (4)
H15	0.5967	0.6759	0.7369	0.024*

Atomic displacement parameters (\AA^2)

	U^{11}	U^{22}	U^{33}	U^{12}	U^{13}	U^{23}
S1	0.0140 (2)	0.0175 (3)	0.0182 (3)	-0.00041 (14)	0.00381 (17)	0.00075 (14)
O1	0.0230 (7)	0.0225 (7)	0.0506 (9)	-0.0015 (6)	0.0103 (7)	-0.0049 (6)
O2	0.0202 (6)	0.0184 (7)	0.0276 (7)	-0.0010 (5)	0.0052 (5)	0.0040 (5)
O3	0.0194 (6)	0.0241 (7)	0.0194 (6)	-0.0005 (5)	0.0061 (5)	-0.0021 (5)
N2	0.0148 (7)	0.0241 (8)	0.0221 (8)	0.0014 (6)	0.0047 (6)	-0.0003 (6)
N3	0.0180 (8)	0.0205 (8)	0.0208 (8)	0.0015 (6)	0.0012 (6)	-0.0009 (6)
N1	0.0161 (7)	0.0206 (8)	0.0226 (8)	-0.0004 (6)	0.0050 (6)	-0.0023 (6)
C1	0.0194 (9)	0.0227 (10)	0.0259 (9)	-0.0017 (7)	0.0013 (7)	0.0021 (7)
C2	0.0179 (9)	0.0251 (10)	0.0242 (9)	-0.0018 (7)	0.0024 (7)	0.0015 (7)
C3	0.0168 (9)	0.0248 (9)	0.0178 (8)	-0.0014 (7)	0.0008 (7)	0.0022 (7)
C4	0.0162 (9)	0.0274 (9)	0.0179 (8)	0.0009 (7)	0.0022 (7)	0.0049 (7)
C5	0.0236 (10)	0.0259 (10)	0.0290 (10)	0.0004 (8)	0.0083 (8)	0.0056 (8)
C6	0.0326 (11)	0.0261 (10)	0.0353 (11)	0.0068 (8)	0.0120 (9)	0.0078 (8)

supplementary materials

C7	0.0220 (10)	0.0400 (12)	0.0266 (10)	0.0123 (8)	0.0080 (8)	0.0114 (8)
C8	0.0153 (9)	0.0450 (12)	0.0241 (10)	-0.0017 (8)	0.0019 (7)	0.0068 (8)
C9	0.0180 (9)	0.0329 (10)	0.0195 (9)	-0.0034 (8)	0.0007 (7)	0.0007 (7)
C10	0.0146 (8)	0.0220 (9)	0.0170 (8)	-0.0006 (7)	0.0012 (6)	0.0019 (7)
C11	0.0190 (9)	0.0203 (9)	0.0200 (8)	-0.0009 (7)	0.0019 (7)	-0.0004 (7)
C12	0.0161 (8)	0.0193 (8)	0.0199 (9)	0.0024 (7)	0.0029 (7)	0.0004 (6)
C13	0.0141 (8)	0.0208 (9)	0.0178 (8)	-0.0012 (7)	0.0015 (6)	0.0019 (6)
C14	0.0195 (9)	0.0171 (8)	0.0240 (9)	-0.0002 (7)	0.0033 (7)	0.0000 (7)
C15	0.0177 (9)	0.0211 (9)	0.0213 (9)	0.0017 (7)	0.0029 (7)	-0.0007 (7)

Geometric parameters (\AA , $^\circ$)

S1—O2	1.4358 (13)	C5—H5	0.9500
S1—O3	1.4386 (13)	C6—C7	1.393 (3)
S1—N3	1.6160 (17)	C6—H6	0.9500
S1—C13	1.7611 (18)	C7—C8	1.379 (3)
O1—C1	1.215 (2)	C7—H7	0.9500
N2—C3	1.298 (2)	C8—C9	1.390 (3)
N2—N1	1.408 (2)	C8—H8	0.9500
N3—H1	0.875 (10)	C9—H9	0.9500
N3—H2	0.873 (10)	C10—C11	1.400 (2)
N1—C1	1.393 (2)	C10—C15	1.403 (3)
N1—C10	1.406 (2)	C11—C12	1.387 (3)
C1—C2	1.499 (3)	C11—H11	0.9500
C2—C3	1.487 (3)	C12—C13	1.392 (2)
C2—H2A	0.9900	C12—H12	0.9500
C2—H2B	0.9900	C13—C14	1.395 (2)
C3—C4	1.468 (2)	C14—C15	1.384 (3)
C4—C9	1.393 (3)	C14—H14	0.9500
C4—C5	1.397 (3)	C15—H15	0.9500
C5—C6	1.387 (3)		
O2—S1—O3	119.00 (8)	C5—C6—C7	120.2 (2)
O2—S1—N3	107.10 (8)	C5—C6—H6	119.9
O3—S1—N3	106.67 (8)	C7—C6—H6	119.9
O2—S1—C13	106.97 (8)	C8—C7—C6	119.86 (19)
O3—S1—C13	107.85 (8)	C8—C7—H7	120.1
N3—S1—C13	108.97 (8)	C6—C7—H7	120.1
C3—N2—N1	107.72 (15)	C7—C8—C9	120.49 (19)
S1—N3—H1	114.1 (16)	C7—C8—H8	119.8
S1—N3—H2	113.3 (16)	C9—C8—H8	119.8
H1—N3—H2	114 (2)	C8—C9—C4	119.88 (19)
C1—N1—C10	129.54 (15)	C8—C9—H9	120.1
C1—N1—N2	112.08 (14)	C4—C9—H9	120.1
C10—N1—N2	118.38 (14)	C11—C10—N1	120.35 (16)
O1—C1—N1	126.49 (17)	C11—C10—C15	120.39 (16)
O1—C1—C2	128.35 (18)	N1—C10—C15	119.26 (16)
N1—C1—C2	105.17 (16)	C12—C11—C10	119.71 (17)
C3—C2—C1	102.40 (15)	C12—C11—H11	120.1
C3—C2—H2A	111.3	C10—C11—H11	120.1

C1—C2—H2A	111.3	C11—C12—C13	119.97 (16)
C3—C2—H2B	111.3	C11—C12—H12	120.0
C1—C2—H2B	111.3	C13—C12—H12	120.0
H2A—C2—H2B	109.2	C12—C13—C14	120.26 (16)
N2—C3—C4	122.26 (17)	C12—C13—S1	119.91 (13)
N2—C3—C2	112.57 (16)	C14—C13—S1	119.81 (14)
C4—C3—C2	125.17 (16)	C15—C14—C13	120.41 (17)
C9—C4—C5	119.68 (17)	C15—C14—H14	119.8
C9—C4—C3	119.44 (17)	C13—C14—H14	119.8
C5—C4—C3	120.88 (17)	C14—C15—C10	119.25 (17)
C6—C5—C4	119.85 (19)	C14—C15—H15	120.4
C6—C5—H5	120.1	C10—C15—H15	120.4
C4—C5—H5	120.1		
C3—N2—N1—C1	-0.1 (2)	C5—C4—C9—C8	0.5 (3)
C3—N2—N1—C10	-179.52 (16)	C3—C4—C9—C8	-179.92 (17)
C10—N1—C1—O1	-2.7 (3)	C1—N1—C10—C11	-1.6 (3)
N2—N1—C1—O1	177.9 (2)	N2—N1—C10—C11	177.72 (15)
C10—N1—C1—C2	177.83 (17)	C1—N1—C10—C15	178.21 (18)
N2—N1—C1—C2	-1.6 (2)	N2—N1—C10—C15	-2.4 (2)
O1—C1—C2—C3	-177.1 (2)	N1—C10—C11—C12	-178.97 (16)
N1—C1—C2—C3	2.34 (19)	C15—C10—C11—C12	1.2 (3)
N1—N2—C3—C4	-178.03 (16)	C10—C11—C12—C13	-0.7 (3)
N1—N2—C3—C2	1.7 (2)	C11—C12—C13—C14	0.2 (3)
C1—C2—C3—N2	-2.6 (2)	C11—C12—C13—S1	178.85 (14)
C1—C2—C3—C4	177.17 (17)	O2—S1—C13—C12	157.60 (15)
N2—C3—C4—C9	173.44 (17)	O3—S1—C13—C12	28.50 (17)
C2—C3—C4—C9	-6.3 (3)	N3—S1—C13—C12	-86.93 (16)
N2—C3—C4—C5	-7.0 (3)	O2—S1—C13—C14	-23.78 (17)
C2—C3—C4—C5	173.27 (18)	O3—S1—C13—C14	-152.88 (15)
C9—C4—C5—C6	-0.3 (3)	N3—S1—C13—C14	91.69 (16)
C3—C4—C5—C6	-179.88 (18)	C12—C13—C14—C15	-0.3 (3)
C4—C5—C6—C7	0.0 (3)	S1—C13—C14—C15	-178.89 (14)
C5—C6—C7—C8	0.2 (3)	C13—C14—C15—C10	0.8 (3)
C6—C7—C8—C9	0.0 (3)	C11—C10—C15—C14	-1.2 (3)
C7—C8—C9—C4	-0.4 (3)	N1—C10—C15—C14	178.94 (16)

Hydrogen-bond geometry (Å, °)

D—H···A	D—H	H···A	D···A	D—H···A
N3—H1···O3 ⁱ	0.88 (1)	2.12 (1)	2.975 (2)	164 (2)
N3—H2···O2 ⁱⁱ	0.87 (1)	2.12 (1)	2.978 (2)	168 (2)

Symmetry codes: (i) -x+2, -y+1, -z+1; (ii) x, -y+3/2, z+1/2.

supplementary materials

Fig. 1

