

Poly[[aquatris(μ_4 -benzene-1,2-dicarboxylato)dilanthanum(III)]hemihydrate]

 Shie Fu Lush^a and Fwu Ming Shen^{b*}
^aDepartment of General Education Center, Yuanpei University, HsinChu, 30015 Taiwan, and ^bDepartment of Biotechnology, Yuanpei University, No. 306, Yuanpei St, HsinChu, 30015 Taiwan


Correspondence e-mail: fmshe@mail.ypu.edu.tw

Received 29 August 2011; accepted 6 September 2011

 Key indicators: single-crystal X-ray study; $T = 150$ K; mean $\sigma(\text{C}-\text{C}) = 0.012$ Å; disorder in solvent or counterion; R factor = 0.037; wR factor = 0.087; data-to-parameter ratio = 11.6.

The asymmetric unit of the title coordination polymer, $\{[\text{La}_2(\text{C}_8\text{H}_4\text{O}_4)_3(\text{H}_2\text{O})]\cdot 0.5\text{H}_2\text{O}\}_n$, contains two independent La^{III} atoms, one of which is surrounded by eight carboxylate-O atoms from six benzene-1,2-dicarboxylate (BDC) anions in a bicapped trigonal-prismatic geometry. The other La^{III} atom is nine-coordinated in a tricapped trigonal-prismatic geometry, formed by eight carboxylate-O atoms from six BDC anions and a coordinated water molecule. The BDC anions bridge the La^{III} atoms, forming a two-dimensional polymeric complex parallel to (001). The crystal structure contains weak $\text{O}-\text{H}\cdots\text{O}$ and non-classical $\text{C}-\text{H}\cdots\text{O}$ hydrogen bonds. A $\text{C}-\text{H}\cdots\pi$ interaction is also present in the crystal structure. The uncoordinated water molecule shows half-occupation.

Related literature

 For a related structure, see: Wang *et al.* (2009).


Experimental

Crystal data

$[\text{La}_2(\text{C}_8\text{H}_4\text{O}_4)_3(\text{H}_2\text{O})]\cdot 0.5\text{H}_2\text{O}$
 $M_r = 1594.36$
 Triclinic, $P\bar{1}$
 $a = 8.6269$ (19) Å
 $b = 10.5832$ (17) Å
 $c = 14.323$ (2) Å
 $\alpha = 97.271$ (18)°
 $\beta = 102.199$ (6)°

$\gamma = 104.489$ (8)°
 $V = 1215.1$ (4) Å³
 $Z = 1$
 Mo $K\alpha$ radiation
 $\mu = 3.54$ mm⁻¹
 $T = 150$ K
 $0.47 \times 0.24 \times 0.04$ mm

Data collection

Nonius KappaCCD diffractometer
 Absorption correction: multi-scan
 (*SCALEPACK*; Otwinowski & Minor, 1997)
 $T_{\text{min}} = 0.287$, $T_{\text{max}} = 0.871$

8765 measured reflections
 4145 independent reflections
 3430 reflections with $I > 2\sigma(I)$
 $R_{\text{int}} = 0.032$

Refinement

$R[F^2 > 2\sigma(F^2)] = 0.037$
 $wR(F^2) = 0.087$
 $S = 1.11$
 4145 reflections
 356 parameters

12 restraints
 H-atom parameters constrained
 $\Delta\rho_{\text{max}} = 1.25$ e Å⁻³
 $\Delta\rho_{\text{min}} = -0.91$ e Å⁻³

Table 1

Selected bond lengths (Å).

La1—O1	2.599 (4)	La2—O1	2.535 (4)
La1—O2	2.645 (5)	La2—O4 ^{iv}	2.482 (4)
La1—O3	2.695 (6)	La2—O5 ^{iv}	2.625 (4)
La1—O4	2.613 (4)	La2—O6 ⁱⁱⁱ	2.466 (4)
La1—O5 ⁱ	2.617 (4)	La2—O8 ^v	2.549 (5)
La1—O7 ⁱⁱ	2.439 (5)	La2—O17	2.608 (4)
La1—O17 ⁱⁱⁱ	2.543 (4)	La2—O18 ⁱⁱⁱ	2.495 (4)
La1—O20	2.478 (5)	La2—O19	2.417 (5)
La1—O22	2.611 (4)		

Symmetry codes: (i) $-x+1, -y+1, -z+1$; (ii) $-x, -y, -z+1$; (iii) $-x, -y+1, -z+1$; (iv) $x-1, y, z$; (v) $-x-1, -y, -z+1$.

Table 2

Hydrogen-bond geometry (Å, °).

 C_g is the centroid of the C2–C7 ring.

$D-H\cdots A$	$D-H$	$H\cdots A$	$D\cdots A$	$D-H\cdots A$
O22—H22A ⁱ ···O1 ⁱⁱ	0.90	2.20	3.001 (6)	148
O22—H22B ⁱ ···O20 ⁱⁱ	0.88	2.15	3.009 (8)	164
O27—H27A ⁱ ···O2 ^{vi}	0.91	2.38	3.17 (2)	146
O27—H27B ⁱ ···O3 ^z	0.88	1.86	2.69 (2)	158
C3—H3 ⁱ ···O27 ^{vii}	0.93	2.15	2.96 (2)	145
C16—H16 ⁱ ···O2 ^z	0.93	2.58	3.331 (10)	138
C19—H19 ⁱ ···C _g ^v	0.93	2.98	3.902 (9)	169

Symmetry codes: (i) $-x+1, -y+1, -z+1$; (ii) $-x, -y, -z+1$; (v) $-x-1, -y, -z+1$; (vi) $x+1, y+1, z$; (vii) $x-1, y-1, z$.

Data collection: *COLLECT* (Nonius, 2000); cell refinement: *SCALEPACK* (Otwinowski & Minor, 1997); data reduction: *DENZO* (Otwinowski & Minor, 1997) and *SCALEPACK*; program(s) used to solve structure: *SHELXS97* (Sheldrick, 2008); program(s) used to refine structure: *SHELXL97* (Sheldrick, 2008); molecular graphics: *PLATON* (Spek, 2009); software used to prepare material for publication: *PLATON*.

This work was supported financially by Yuanpei University, Taiwan.

Supplementary data and figures for this paper are available from the IUCr electronic archives (Reference: XU5316).

References

- Nonius (2000). *COLLECT*. Nonius BV, Delft, The Netherlands.
- Otwinowski, Z. & Minor, W. (1997). *Methods in Enzymology*, Vol. 276, *Macromolecular Crystallography*, Part A, edited by C. W. Carter Jr & R. M. Sweet, pp. 307–326. New York: Academic Press.
- Sheldrick, G. M. (2008). *Acta Cryst.* **A64**, 112–122.
- Spek, A. L. (2009). *Acta Cryst.* **D65**, 148–155.
- Wang, G.-M., Xue, S.-Y., Li, H. & Liu, H.-L. (2009). *Acta Cryst.* **C65**, m469–m471.

supplementary materials

Acta Cryst. (2011). E67, m1370-m1371 [doi:10.1107/S1600536811036282]

Poly[[aquatris(μ_4 -benzene-1,2-dicarboxylato)dilanthanum(III)] hemihydrate]

S. F. Lush and F. M. Shen

Comment

Benzene-1,2-dicarboxylic acid (H_2BDC) are widely used in the construction of coordination polymers due to their capability of acting as bridging ligands in various coordination modes. But to the best of our knowledge, H_2BDC is seldom involved in lanthanide complexes (Wang *et al.*, 2009). In this paper, we describe the hydrothermal synthesis and structure properties of a lanthanide phthalate coordination complex $\{[La_2(C_8H_4O_4)_3(H_2O)] \cdot 0.5 H_2O\}_n$.

The molecular structure of the title compound is shown in Fig. 1. There are two independent lanthanum ions in the asymmetric unit. The La(1) ion is nine-coordinated with O_9 donors sets to form tricapped trigonal prismatic geometries by eight carboxylate O atoms and one water molecule, where La(2) ion is eight-coordinated with O_8 donors sets to form distorted bicapped trigonal-prismatic geometries by eight carboxylate O atoms, from six benzene-1,2-dicarboxylate anions. The selected bond lengths (\AA) of title compound are listed in Table 1. The two La^{III} cations are separated by a non-bonding distance of 4.453 (9) and 4.419 (10) \AA . The benzene-1,2-dicarboxylate anions bridge the La^{III} cations, forming a two-dimensional polymeric complex.

There are extensive intermolecular $O-H\cdots O$ and weak $C-H\cdots O$ hydrogen bonds, which cause the stability of the crystal structure (Fig. 2, Table 2). There are no $\pi-\pi$ stacking interactions in the title compound. Furthermore, there is $C-H\cdots\pi$ interaction between $C-H$ group of the BDC ligand, with an $C-H\cdots$ centroid distance of 3.902 (9) \AA [$C19-H19\cdots Cg1^V(C2-C7)$] (Symmetry code: $-1-X, -Y, 1-Z$).

Experimental

$LaCl_3 \cdot 6H_2O$ (0.0868 g, 0.20 mmol), benzene-1,2-dicarboxylic acid (0.0348 g, 0.20 mmol) and 1,2-bis(4-pyridyl)ethane were mixed in 10 ml of deionized water. After stirring for 30 min, the mixture was placed in a 23 ml Teflon-lined reactor, heated at 453 K for 48 h, then cooled slowly to room temperature. The colorless transparent single crystals of the title compound were obtained in 36.76% yield (based on La).

Refinement

The site occupancy factor of the lattice water O27 was refined to 0.46 (3), and was set as 0.5 at the final cycles of refinement. Water H atoms were placed in calculated positions and refined in riding mode with $U_{iso}(H) = 1.5U_{eq}(O)$. Other H atoms were positioned geometrically with $C-H = 0.93 \text{\AA}$ and refined using a riding model with $U_{iso}(H) = 1.2U_{eq}(C)$.

Figures




Fig. 1. View of the title compound with the atom numbering scheme. Displacement ellipsoids for non-H atoms are drawn at the 50% probability level. H atoms have been omitted for clarity. Symmetry codes: (i) $-x, -y, -z + 1$; (ii) $-x, -y + 1, -z + 1$; (iii) $-x + 1, -y + 1, -z + 1$; (iv) $x - 1, y, z$; (v) $-x - 1, -y, -z + 1$.




Fig. 2. The molecular packing for the title compound, viewed along the c axis. Hydrogen bonds are shown as dashed lines.

Poly[[aquatris(μ_4 -benzene-1,2-dicarboxylato)dilanthanum(III)] hemihydrate]

Crystal data

$[\text{La}_2(\text{C}_8\text{H}_4\text{O}_4)_3(\text{H}_2\text{O})] \cdot 0.5\text{H}_2\text{O}$

$M_r = 1594.36$

Triclinic, $P\bar{1}$

Hall symbol: $-P\ 1$

$a = 8.6269\ (19)\ \text{\AA}$

$b = 10.5832\ (17)\ \text{\AA}$

$c = 14.323\ (2)\ \text{\AA}$

$\alpha = 97.271\ (18)^\circ$

$\beta = 102.199\ (6)^\circ$

$\gamma = 104.489\ (8)^\circ$

$V = 1215.1\ (4)\ \text{\AA}^3$

$Z = 1$

$F(000) = 762$

$D_x = 2.179\ \text{Mg m}^{-3}$

Mo $K\alpha$ radiation, $\lambda = 0.71073\ \text{\AA}$

Cell parameters from 4145 reflections

$\theta = 2.0\text{--}25.0^\circ$

$\mu = 3.54\ \text{mm}^{-1}$

$T = 150\ \text{K}$

Prism, colorless

$0.47 \times 0.24 \times 0.04\ \text{mm}$

Data collection

Nonius KappaCCD
diffractometer

Radiation source: fine-focus sealed tube
graphite

Detector resolution: 9 pixels mm^{-1}

$\omega/2\theta$ scans

Absorption correction: multi-scan
(*SCALEPACK*; Otwinowski & Minor, 1997)

$T_{\min} = 0.287$, $T_{\max} = 0.871$

8765 measured reflections

4145 independent reflections

3430 reflections with $I > 2\sigma(I)$

$R_{\text{int}} = 0.032$

$\theta_{\max} = 25.0^\circ$, $\theta_{\min} = 2.0^\circ$

$h = -10 \rightarrow 10$

$k = -12 \rightarrow 12$

$l = -17 \rightarrow 16$

Refinement

Refinement on F^2	Primary atom site location: structure-invariant direct methods
Least-squares matrix: full	Secondary atom site location: difference Fourier map
$R[F^2 > 2\sigma(F^2)] = 0.037$	Hydrogen site location: inferred from neighbouring sites
$wR(F^2) = 0.087$	H-atom parameters constrained
$S = 1.11$	$w = 1/[\sigma^2(F_o^2) + (0.0369P)^2 + 2.5018P]$
4145 reflections	where $P = (F_o^2 + 2F_c^2)/3$
356 parameters	$(\Delta/\sigma)_{\max} = 0.002$
12 restraints	$\Delta\rho_{\max} = 1.25 \text{ e } \text{\AA}^{-3}$
	$\Delta\rho_{\min} = -0.91 \text{ e } \text{\AA}^{-3}$

Special details

Geometry. Bond distances, angles *etc.* have been calculated using the rounded fractional coordinates. All su's are estimated from the variances of the (full) variance-covariance matrix. The cell e.s.d.'s are taken into account in the estimation of distances, angles and torsion angles

Refinement. Refinement on F^2 for ALL reflections except those flagged by the user for potential systematic errors. Weighted R -factors wR and all goodnesses of fit S are based on F^2 , conventional R -factors R are based on F , with F set to zero for negative F^2 . The observed criterion of $F^2 > \sigma(F^2)$ is used only for calculating $-R$ -factor-obs *etc.* and is not relevant to the choice of reflections for refinement. R -factors based on F^2 are statistically about twice as large as those based on F , and R -factors based on ALL data will be even larger.

Fractional atomic coordinates and isotropic or equivalent isotropic displacement parameters (\AA^2)

	x	y	z	$U_{\text{iso}}^*/U_{\text{eq}}$	Occ. (<1)
La1	0.16724 (4)	0.22260 (3)	0.50902 (3)	0.0255 (1)	
La2	-0.30351 (4)	0.34775 (3)	0.49673 (3)	0.0181 (1)	
O1	-0.0850 (4)	0.2911 (4)	0.4193 (3)	0.0186 (13)	
O2	0.0101 (5)	0.1690 (4)	0.3225 (4)	0.0377 (16)	
O3	0.3563 (6)	0.2580 (6)	0.6913 (4)	0.048 (2)	
O4	0.4799 (5)	0.3465 (4)	0.5867 (3)	0.0239 (14)	
O5	0.7181 (5)	0.5887 (4)	0.5813 (3)	0.0214 (14)	
O6	0.5239 (5)	0.6448 (4)	0.6419 (3)	0.0233 (14)	
O7	-0.3449 (6)	-0.1149 (4)	0.5627 (4)	0.0351 (18)	
O8	-0.5707 (5)	-0.1219 (4)	0.6165 (4)	0.0312 (16)	
O17	-0.1888 (5)	0.5423 (4)	0.4122 (3)	0.0208 (14)	
O18	0.0471 (5)	0.5575 (4)	0.3684 (3)	0.0217 (14)	
O19	-0.2597 (5)	0.1804 (4)	0.5913 (4)	0.0313 (16)	
O20	0.0031 (5)	0.1774 (4)	0.6299 (4)	0.0369 (16)	
O22	0.1603 (6)	-0.0025 (4)	0.5669 (4)	0.0436 (18)	
C1	-0.0786 (7)	0.2446 (6)	0.3327 (6)	0.029 (2)	
C2	-0.1747 (7)	0.2850 (7)	0.2501 (5)	0.032 (2)	
C3	-0.2494 (9)	0.1966 (9)	0.1623 (6)	0.050 (3)	

supplementary materials

C4	-0.3483 (10)	0.2326 (13)	0.0877 (6)	0.071 (4)	
C5	-0.3728 (10)	0.3569 (13)	0.1003 (6)	0.067 (4)	
C6	-0.2982 (8)	0.4457 (9)	0.1859 (5)	0.040 (3)	
C7	-0.1968 (7)	0.4135 (7)	0.2618 (5)	0.028 (2)	
C8	-0.1036 (7)	0.5116 (6)	0.3539 (5)	0.0192 (19)	
C9	0.4775 (8)	0.3355 (7)	0.6748 (6)	0.033 (2)	
C10	0.6412 (7)	0.5955 (6)	0.6477 (5)	0.0205 (19)	
C11	0.6971 (8)	0.5475 (8)	0.7397 (5)	0.032 (2)	
C12	0.6165 (8)	0.4227 (8)	0.7530 (5)	0.037 (3)	
C13	0.6653 (10)	0.3832 (12)	0.8412 (7)	0.069 (4)	
C14	0.7926 (11)	0.4676 (16)	0.9153 (7)	0.091 (6)	
C15	0.8710 (11)	0.5890 (14)	0.9012 (7)	0.079 (5)	
C16	0.8232 (9)	0.6330 (9)	0.8153 (6)	0.046 (3)	
C17	-0.4155 (8)	-0.0836 (6)	0.6270 (5)	0.0254 (19)	
C18	-0.3075 (7)	-0.0081 (6)	0.7229 (5)	0.027 (2)	
C19	-0.3358 (9)	-0.0561 (7)	0.8052 (6)	0.040 (3)	
C20	-0.2256 (10)	-0.0039 (9)	0.8933 (7)	0.059 (3)	
C21	-0.0847 (11)	0.1003 (10)	0.9026 (7)	0.070 (4)	
C22	-0.0567 (9)	0.1496 (8)	0.8216 (7)	0.055 (3)	
C23	-0.1669 (7)	0.0972 (6)	0.7317 (6)	0.033 (2)	
C24	-0.1373 (7)	0.1562 (6)	0.6441 (6)	0.028 (2)	
O27	0.717 (3)	0.922 (2)	0.1959 (16)	0.163 (9)*	0.500
H3	-0.23260	0.11290	0.15400	0.0600*	
H4	-0.39840	0.17330	0.02890	0.0860*	
H5	-0.44080	0.38060	0.05010	0.0810*	
H6	-0.31620	0.52910	0.19300	0.0480*	
H13	0.61180	0.29940	0.85010	0.0830*	
H14	0.82430	0.44130	0.97420	0.1090*	
H15	0.95910	0.64440	0.95050	0.0940*	
H16	0.87490	0.71840	0.80830	0.0550*	
H19	-0.43070	-0.12450	0.80030	0.0480*	
H20	-0.24470	-0.03830	0.94780	0.0710*	
H21	-0.01040	0.13620	0.96300	0.0840*	
H22	0.03760	0.21900	0.82730	0.0660*	
H22A	0.17560	-0.07490	0.58980	0.0650*	
H22B	0.10320	-0.04070	0.50710	0.0650*	
H27A	0.82020	0.96320	0.23430	0.2440*	0.500
H27B	0.67000	0.87010	0.23130	0.2440*	0.500

Atomic displacement parameters (\AA^2)

	U^{11}	U^{22}	U^{33}	U^{12}	U^{13}	U^{23}
La1	0.0128 (2)	0.0071 (2)	0.0613 (3)	0.0054 (1)	0.0149 (2)	0.0089 (2)
La2	0.0126 (2)	0.0076 (2)	0.0386 (3)	0.0058 (1)	0.0113 (2)	0.0063 (2)
O1	0.0111 (19)	0.0089 (19)	0.036 (3)	0.0011 (16)	0.0098 (18)	0.0016 (19)
O2	0.021 (2)	0.024 (2)	0.062 (4)	0.009 (2)	0.008 (2)	-0.015 (2)
O3	0.022 (3)	0.069 (4)	0.069 (4)	0.013 (3)	0.020 (3)	0.053 (3)
O4	0.017 (2)	0.022 (2)	0.042 (3)	0.0137 (18)	0.012 (2)	0.015 (2)

O5	0.017 (2)	0.018 (2)	0.029 (3)	0.0066 (17)	0.0060 (19)	0.0004 (19)
O6	0.017 (2)	0.014 (2)	0.042 (3)	0.0082 (18)	0.010 (2)	0.005 (2)
O7	0.034 (3)	0.017 (2)	0.069 (4)	0.016 (2)	0.030 (3)	0.015 (2)
O8	0.017 (2)	0.010 (2)	0.070 (4)	0.0035 (18)	0.020 (2)	0.005 (2)
O17	0.017 (2)	0.015 (2)	0.037 (3)	0.0098 (17)	0.0123 (19)	0.0093 (19)
O18	0.017 (2)	0.017 (2)	0.036 (3)	0.0084 (18)	0.0109 (19)	0.0081 (19)
O19	0.014 (2)	0.016 (2)	0.066 (4)	0.0068 (18)	0.006 (2)	0.017 (2)
O20	0.019 (2)	0.022 (2)	0.080 (4)	0.0089 (19)	0.020 (2)	0.028 (3)
O22	0.039 (3)	0.017 (2)	0.092 (4)	0.018 (2)	0.033 (3)	0.024 (3)
C1	0.015 (3)	0.016 (3)	0.051 (5)	0.000 (3)	0.012 (3)	-0.010 (3)
C2	0.010 (3)	0.041 (4)	0.037 (5)	0.001 (3)	0.007 (3)	-0.012 (3)
C3	0.023 (4)	0.071 (6)	0.044 (5)	0.003 (4)	0.015 (4)	-0.023 (4)
C4	0.028 (5)	0.129 (10)	0.031 (6)	-0.007 (6)	0.013 (4)	-0.029 (6)
C5	0.026 (4)	0.151 (11)	0.020 (5)	0.013 (6)	0.010 (4)	0.015 (6)
C6	0.022 (4)	0.076 (6)	0.028 (5)	0.017 (4)	0.011 (3)	0.020 (4)
C7	0.015 (3)	0.046 (4)	0.025 (4)	0.012 (3)	0.009 (3)	0.002 (3)
C8	0.019 (3)	0.016 (3)	0.031 (4)	0.009 (3)	0.012 (3)	0.017 (3)
C9	0.018 (3)	0.040 (4)	0.056 (5)	0.020 (3)	0.015 (3)	0.031 (4)
C10	0.008 (3)	0.017 (3)	0.030 (4)	-0.003 (2)	0.003 (3)	-0.003 (3)
C11	0.017 (3)	0.063 (5)	0.026 (4)	0.023 (3)	0.011 (3)	0.008 (4)
C12	0.017 (3)	0.068 (5)	0.037 (5)	0.017 (4)	0.011 (3)	0.029 (4)
C13	0.031 (5)	0.133 (9)	0.067 (7)	0.033 (6)	0.025 (5)	0.060 (7)
C14	0.027 (5)	0.216 (15)	0.044 (6)	0.039 (7)	0.014 (5)	0.059 (8)
C15	0.024 (5)	0.178 (13)	0.033 (6)	0.041 (7)	-0.002 (4)	0.003 (7)
C16	0.023 (4)	0.083 (6)	0.036 (5)	0.026 (4)	0.007 (3)	0.002 (4)
C17	0.024 (3)	0.011 (3)	0.048 (4)	0.008 (2)	0.014 (3)	0.016 (3)
C18	0.017 (3)	0.019 (3)	0.052 (5)	0.008 (3)	0.015 (3)	0.014 (3)
C19	0.027 (4)	0.033 (4)	0.060 (6)	0.002 (3)	0.015 (4)	0.016 (4)
C20	0.047 (5)	0.061 (6)	0.069 (7)	0.005 (4)	0.012 (5)	0.036 (5)
C21	0.045 (5)	0.080 (7)	0.061 (7)	-0.016 (5)	-0.008 (5)	0.031 (6)
C22	0.025 (4)	0.053 (5)	0.076 (7)	-0.006 (4)	0.002 (4)	0.030 (5)
C23	0.016 (3)	0.027 (3)	0.060 (5)	0.010 (3)	0.008 (3)	0.022 (4)
C24	0.016 (3)	0.013 (3)	0.063 (5)	0.008 (3)	0.016 (3)	0.015 (3)

Geometric parameters (Å, °)

La1—O1	2.599 (4)	C2—C7	1.414 (10)
La1—O2	2.645 (5)	C3—C4	1.377 (13)
La1—O3	2.695 (6)	C4—C5	1.380 (18)
La1—O4	2.613 (4)	C5—C6	1.369 (12)
La1—O5 ⁱ	2.617 (4)	C6—C7	1.380 (10)
La1—O7 ⁱⁱ	2.439 (5)	C7—C8	1.501 (10)
La1—O17 ⁱⁱⁱ	2.543 (4)	C9—C12	1.470 (11)
La1—O20	2.478 (5)	C10—C11	1.500 (10)
La1—O22	2.611 (4)	C11—C16	1.392 (11)
La2—O1	2.535 (4)	C11—C12	1.387 (12)
La2—O4 ^{iv}	2.482 (4)	C12—C13	1.395 (13)
La2—O5 ^{iv}	2.625 (4)	C13—C14	1.381 (16)

supplementary materials

La2—O6 ⁱⁱⁱ	2.466 (4)	C14—C15	1.36 (2)
La2—O8 ^v	2.549 (5)	C15—C16	1.388 (14)
La2—O17	2.608 (4)	C17—C18	1.484 (10)
La2—O18 ⁱⁱⁱ	2.495 (4)	C18—C19	1.385 (10)
La2—O19	2.417 (5)	C18—C23	1.397 (9)
O1—C1	1.293 (9)	C19—C20	1.362 (13)
O2—C1	1.253 (8)	C20—C21	1.391 (14)
O3—C9	1.244 (9)	C21—C22	1.373 (14)
O4—C9	1.286 (9)	C22—C23	1.381 (12)
O5—C10	1.274 (8)	C23—C24	1.511 (11)
O6—C10	1.242 (8)	C3—H3	0.9300
O7—C17	1.261 (9)	C4—H4	0.9300
O8—C17	1.268 (9)	C5—H5	0.9300
O17—C8	1.286 (8)	C6—H6	0.9300
O18—C8	1.230 (8)	C13—H13	0.9300
O19—C24	1.265 (9)	C14—H14	0.9300
O20—C24	1.240 (8)	C15—H15	0.9300
O22—H22B	0.8800	C16—H16	0.9300
O22—H22A	0.9000	C19—H19	0.9300
O27—H27B	0.8800	C20—H20	0.9300
O27—H27A	0.9100	C21—H21	0.9300
C1—C2	1.470 (10)	C22—H22	0.9300
C2—C3	1.389 (11)		
O1—La1—O2	49.50 (14)	La1 ⁱⁱⁱ —O17—C8	125.3 (4)
O1—La1—O3	138.66 (15)	La2 ⁱⁱⁱ —O18—C8	141.3 (4)
O1—La1—O4	134.61 (13)	La2—O19—C24	136.6 (4)
O1—La1—O20	83.93 (14)	La1—O20—C24	145.0 (5)
O1—La1—O22	126.61 (15)	La1—O22—H22B	86.00
O1—La1—O7 ⁱⁱ	127.83 (16)	H22A—O22—H22B	99.00
O1—La1—O17 ⁱⁱⁱ	71.11 (14)	La1—O22—H22A	171.00
O1—La1—O5 ⁱ	73.05 (13)	H27A—O27—H27B	104.00
O2—La1—O3	171.83 (16)	O1—C1—O2	119.2 (7)
O2—La1—O4	127.22 (14)	O1—C1—C2	118.0 (6)
O2—La1—O20	118.61 (16)	O2—C1—C2	122.8 (7)
O2—La1—O22	108.06 (15)	C3—C2—C7	119.8 (7)
O2—La1—O7 ⁱⁱ	78.77 (16)	C1—C2—C3	120.7 (7)
O2—La1—O17 ⁱⁱⁱ	111.42 (13)	C1—C2—C7	119.4 (6)
O2—La1—O5 ⁱ	68.20 (13)	C2—C3—C4	120.2 (9)
O3—La1—O4	48.67 (15)	C3—C4—C5	119.8 (9)
O3—La1—O20	66.90 (17)	C4—C5—C6	120.7 (9)
O3—La1—O22	67.90 (18)	C5—C6—C7	121.0 (9)
O3—La1—O7 ⁱⁱ	93.09 (18)	C2—C7—C6	118.4 (7)
O3—La1—O17 ⁱⁱⁱ	74.93 (16)	C2—C7—C8	119.1 (6)
O3—La1—O5 ⁱ	111.83 (16)	C6—C7—C8	122.4 (7)
O4—La1—O20	113.75 (15)	O17—C8—C7	116.9 (6)
O4—La1—O22	98.40 (15)	O18—C8—C7	118.5 (6)

O4—La1—O7 ⁱⁱ	68.65 (15)	O17—C8—O18	124.6 (6)
O4—La1—O17 ⁱⁱⁱ	71.98 (14)	O4—C9—C12	117.6 (6)
O4—La1—O5 ⁱ	67.52 (13)	O3—C9—O4	119.8 (7)
O20—La1—O22	65.81 (16)	O3—C9—C12	122.5 (7)
O7 ⁱⁱ —La1—O20	135.44 (15)	O6—C10—C11	117.4 (6)
O17 ⁱⁱⁱ —La1—O20	79.05 (14)	O5—C10—C11	119.5 (6)
O5 ⁱ —La1—O20	140.95 (14)	O5—C10—O6	123.1 (6)
O7 ⁱⁱ —La1—O22	69.81 (16)	C10—C11—C16	119.5 (7)
O17 ⁱⁱⁱ —La1—O22	136.23 (15)	C12—C11—C16	119.6 (7)
O5 ⁱ —La1—O22	152.50 (16)	C10—C11—C12	120.8 (6)
O7 ⁱⁱ —La1—O17 ⁱⁱⁱ	135.85 (15)	C9—C12—C11	119.4 (7)
O5 ⁱ —La1—O7 ⁱⁱ	82.87 (15)	C9—C12—C13	120.9 (8)
O5 ⁱ —La1—O17 ⁱⁱⁱ	63.96 (13)	C11—C12—C13	119.6 (8)
O1—La2—O17	71.82 (14)	C12—C13—C14	120.5 (11)
O1—La2—O19	84.69 (15)	C13—C14—C15	119.3 (10)
O1—La2—O4 ^{iv}	164.86 (14)	C14—C15—C16	121.8 (10)
O1—La2—O5 ^{iv}	123.63 (14)	C11—C16—C15	119.1 (9)
O1—La2—O8 ^v	71.19 (14)	O7—C17—O8	123.7 (6)
O1—La2—O6 ⁱⁱⁱ	104.32 (14)	O8—C17—C18	119.0 (6)
O1—La2—O18 ⁱⁱⁱ	77.94 (14)	O7—C17—C18	117.1 (6)
O17—La2—O19	150.39 (15)	C17—C18—C23	122.3 (6)
O4 ^{iv} —La2—O17	123.32 (14)	C17—C18—C19	117.9 (6)
O5 ^{iv} —La2—O17	62.98 (13)	C19—C18—C23	119.2 (7)
O8 ^v —La2—O17	114.56 (15)	C18—C19—C20	120.5 (7)
O6 ⁱⁱⁱ —La2—O17	70.92 (14)	C19—C20—C21	120.7 (9)
O17—La2—O18 ⁱⁱⁱ	87.01 (14)	C20—C21—C22	119.3 (9)
O4 ^{iv} —La2—O19	80.80 (15)	C21—C22—C23	120.7 (8)
O5 ^{iv} —La2—O19	121.13 (15)	C18—C23—C22	119.6 (7)
O8 ^v —La2—O19	72.45 (17)	C18—C23—C24	120.4 (7)
O6 ⁱⁱⁱ —La2—O19	134.18 (15)	C22—C23—C24	119.9 (6)
O18 ⁱⁱⁱ —La2—O19	70.27 (15)	O19—C24—C23	116.4 (6)
O4 ^{iv} —La2—O5 ^{iv}	68.53 (14)	O20—C24—C23	118.2 (6)
O4 ^{iv} —La2—O8 ^v	100.10 (15)	O19—C24—O20	125.5 (7)
O4 ^{iv} —La2—O6 ⁱⁱⁱ	83.05 (14)	C2—C3—H3	120.00
O4 ^{iv} —La2—O18 ⁱⁱⁱ	100.99 (14)	C4—C3—H3	120.00
O5 ^{iv} —La2—O8 ^v	158.53 (14)	C5—C4—H4	120.00
O5 ^{iv} —La2—O6 ⁱⁱⁱ	91.38 (14)	C3—C4—H4	120.00
O5 ^{iv} —La2—O18 ⁱⁱⁱ	68.20 (14)	C4—C5—H5	120.00
O6 ⁱⁱⁱ —La2—O8 ^v	68.66 (15)	C6—C5—H5	120.00
O8 ^v —La2—O18 ⁱⁱⁱ	133.08 (15)	C7—C6—H6	120.00
O6 ⁱⁱⁱ —La2—O18 ⁱⁱⁱ	155.34 (14)	C5—C6—H6	119.00

supplementary materials

La1—O1—La2	124.95 (16)	C12—C13—H13	120.00
La1—O1—C1	95.4 (4)	C14—C13—H13	120.00
La2—O1—C1	136.0 (4)	C15—C14—H14	120.00
La1—O2—C1	94.3 (5)	C13—C14—H14	120.00
La1—O3—C9	93.6 (5)	C14—C15—H15	119.00
La1—O4—C9	96.4 (4)	C16—C15—H15	119.00
La1—O4—La2 ^{vi}	122.76 (16)	C15—C16—H16	120.00
La2 ^{vi} —O4—C9	132.0 (4)	C11—C16—H16	120.00
La2 ^{vi} —O5—C10	115.6 (4)	C18—C19—H19	120.00
La1 ⁱ —O5—C10	127.7 (4)	C20—C19—H19	120.00
La1 ⁱ —O5—La2 ^{vi}	114.93 (16)	C21—C20—H20	120.00
La2 ⁱⁱⁱ —O6—C10	132.6 (4)	C19—C20—H20	120.00
La1 ⁱⁱ —O7—C17	156.2 (5)	C20—C21—H21	120.00
La2 ^v —O8—C17	110.5 (4)	C22—C21—H21	120.00
La2—O17—C8	112.8 (4)	C21—C22—H22	120.00
La1 ⁱⁱⁱ —O17—La2	118.14 (16)	C23—C22—H22	120.00
O2—La1—O1—La2	154.4 (3)	O5 ^{iv} —La2—O19—C24	-74.0 (7)
O2—La1—O1—C1	-7.1 (3)	O8 ^v —La2—O19—C24	124.4 (7)
O3—La1—O1—La2	-26.0 (3)	O6 ⁱⁱⁱ —La2—O19—C24	157.3 (6)
O3—La1—O1—C1	172.4 (4)	O18 ⁱⁱⁱ —La2—O19—C24	-26.6 (7)
O4—La1—O1—La2	-99.6 (2)	O17—La2—O4 ^{iv} —La1 ^{iv}	97.8 (2)
O4—La1—O1—C1	98.9 (4)	O17—La2—O4 ^{iv} —C9 ^{iv}	-122.9 (6)
O20—La1—O1—La2	18.1 (2)	O19—La2—O4 ^{iv} —La1 ^{iv}	-100.8 (2)
O20—La1—O1—C1	-143.5 (4)	O19—La2—O4 ^{iv} —C9 ^{iv}	38.4 (6)
O22—La1—O1—La2	71.7 (3)	O1—La2—O5 ^{iv} —C10 ^{iv}	-153.0 (4)
O22—La1—O1—C1	-89.8 (4)	O1—La2—O5 ^{iv} —La1 ⁱⁱⁱ	40.9 (2)
O7 ⁱⁱ —La1—O1—La2	163.59 (18)	O17—La2—O5 ^{iv} —C10 ^{iv}	166.4 (5)
O7 ⁱⁱ —La1—O1—C1	2.1 (4)	O17—La2—O5 ^{iv} —La1 ⁱⁱⁱ	0.34 (15)
O17 ⁱⁱⁱ —La1—O1—La2	-62.4 (2)	O19—La2—O5 ^{iv} —C10 ^{iv}	-47.3 (5)
O17 ⁱⁱⁱ —La1—O1—C1	136.0 (4)	O19—La2—O5 ^{iv} —La1 ⁱⁱⁱ	146.65 (17)
O5 ⁱ —La1—O1—La2	-130.1 (2)	O1—La2—O8 ^v —C17 ^v	-170.5 (5)
O5 ⁱ —La1—O1—C1	68.4 (4)	O17—La2—O8 ^v —C17 ^v	-111.7 (4)
O1—La1—O2—C1	7.3 (3)	O19—La2—O8 ^v —C17 ^v	99.3 (5)
O4—La1—O2—C1	-113.4 (4)	O1—La2—O6 ⁱⁱⁱ —C10 ⁱⁱⁱ	-162.0 (5)
O20—La1—O2—C1	58.7 (4)	O17—La2—O6 ⁱⁱⁱ —C10 ⁱⁱⁱ	-97.4 (5)
O22—La1—O2—C1	130.4 (4)	O19—La2—O6 ⁱⁱⁱ —C10 ⁱⁱⁱ	101.4 (6)
O7 ⁱⁱ —La1—O2—C1	-165.4 (4)	O1—La2—O18 ⁱⁱⁱ —C8 ⁱⁱⁱ	27.4 (6)
O17 ⁱⁱⁱ —La1—O2—C1	-30.3 (4)	O17—La2—O18 ⁱⁱⁱ —C8 ⁱⁱⁱ	-44.7 (6)
O5 ⁱ —La1—O2—C1	-78.6 (4)	O19—La2—O18 ⁱⁱⁱ —C8 ⁱⁱⁱ	116.0 (7)
O1—La1—O3—C9	-107.5 (5)	La1—O1—C1—O2	13.1 (6)
O4—La1—O3—C9	7.1 (4)	La1—O1—C1—C2	-166.0 (5)
O20—La1—O3—C9	-156.2 (5)	La2—O1—C1—O2	-144.9 (5)

O22—La1—O3—C9	131.7 (5)	La2—O1—C1—C2	36.0 (9)
O7 ⁱⁱ —La1—O3—C9	64.9 (5)	La1—O2—C1—O1	-12.9 (6)
O17 ⁱⁱⁱ —La1—O3—C9	-71.9 (5)	La1—O2—C1—C2	166.2 (6)
O5 ⁱ —La1—O3—C9	-18.6 (5)	La1—O3—C9—O4	-12.6 (7)
O1—La1—O4—C9	115.6 (4)	La1—O3—C9—C12	163.6 (6)
O1—La1—O4—La2 ^{vi}	-93.6 (2)	La1—O4—C9—O3	13.1 (7)
O2—La1—O4—C9	-177.8 (4)	La1—O4—C9—C12	-163.3 (6)
O2—La1—O4—La2 ^{vi}	-27.0 (3)	La2 ^{vi} —O4—C9—O3	-133.4 (6)
O3—La1—O4—C9	-6.9 (4)	La2 ^{vi} —O4—C9—C12	50.2 (9)
O3—La1—O4—La2 ^{vi}	143.9 (3)	La2 ^{vi} —O5—C10—O6	-116.6 (6)
O20—La1—O4—C9	9.8 (4)	La2 ^{vi} —O5—C10—C11	66.1 (7)
O20—La1—O4—La2 ^{vi}	160.59 (18)	La1 ⁱ —O5—C10—O6	47.4 (8)
O22—La1—O4—C9	-57.4 (4)	La1 ⁱ —O5—C10—C11	-129.9 (5)
O22—La1—O4—La2 ^{vi}	93.4 (2)	La2 ⁱⁱⁱ —O6—C10—O5	33.2 (9)
O7 ⁱⁱ —La1—O4—C9	-121.8 (4)	La2 ⁱⁱⁱ —O6—C10—C11	-149.4 (5)
O7 ⁱⁱ —La1—O4—La2 ^{vi}	29.00 (19)	La1 ⁱⁱ —O7—C17—O8	-113.2 (11)
O17 ⁱⁱⁱ —La1—O4—C9	78.6 (4)	La1 ⁱⁱ —O7—C17—C18	61.0 (13)
O17 ⁱⁱⁱ —La1—O4—La2 ^{vi}	-130.6 (2)	La2 ^v —O8—C17—O7	29.7 (8)
O5 ⁱ —La1—O4—C9	147.2 (4)	La2 ^v —O8—C17—C18	-144.4 (5)
O5 ⁱ —La1—O4—La2 ^{vi}	-62.00 (19)	La2—O17—C8—O18	-110.4 (6)
O1—La1—O20—C24	19.1 (7)	La2—O17—C8—C7	70.1 (6)
O2—La1—O20—C24	-17.6 (8)	La1 ⁱⁱⁱ —O17—C8—O18	47.3 (8)
O3—La1—O20—C24	169.2 (8)	La1 ⁱⁱⁱ —O17—C8—C7	-132.3 (5)
O4—La1—O20—C24	155.6 (7)	La2 ⁱⁱⁱ —O18—C8—O17	27.6 (11)
O22—La1—O20—C24	-115.7 (8)	La2 ⁱⁱⁱ —O18—C8—C7	-152.9 (5)
O7 ⁱⁱ —La1—O20—C24	-121.3 (7)	La2—O19—C24—O20	-34.9 (11)
O17 ⁱⁱⁱ —La1—O20—C24	91.0 (7)	La2—O19—C24—C23	144.8 (5)
O5 ⁱ —La1—O20—C24	72.5 (8)	La1—O20—C24—O19	-17.0 (12)
O1—La1—O7 ⁱⁱ —C17 ⁱⁱ	26.2 (12)	La1—O20—C24—C23	163.3 (5)
O2—La1—O7 ⁱⁱ —C17 ⁱⁱ	33.3 (11)	O1—C1—C2—C3	-144.4 (7)
O3—La1—O7 ⁱⁱ —C17 ⁱⁱ	-147.4 (11)	O1—C1—C2—C7	33.0 (9)
O4—La1—O7 ⁱⁱ —C17 ⁱⁱ	-104.4 (11)	O2—C1—C2—C3	36.6 (10)
O20—La1—O7 ⁱⁱ —C17 ⁱⁱ	152.9 (10)	O2—C1—C2—C7	-146.1 (7)
O22—La1—O7 ⁱⁱ —C17 ⁱⁱ	147.5 (11)	C1—C2—C3—C4	175.6 (8)
O1—La1—O17 ⁱⁱⁱ —La2 ⁱⁱⁱ	-80.23 (19)	C7—C2—C3—C4	-1.7 (12)
O1—La1—O17 ⁱⁱⁱ —C8 ⁱⁱⁱ	76.4 (5)	C1—C2—C7—C6	-174.9 (6)
O2—La1—O17 ⁱⁱⁱ —La2 ⁱⁱⁱ	-50.9 (2)	C1—C2—C7—C8	7.9 (9)
O2—La1—O17 ⁱⁱⁱ —C8 ⁱⁱⁱ	105.7 (5)	C3—C2—C7—C6	2.4 (10)
O3—La1—O17 ⁱⁱⁱ —La2 ⁱⁱⁱ	123.7 (2)	C3—C2—C7—C8	-174.8 (7)
O3—La1—O17 ⁱⁱⁱ —C8 ⁱⁱⁱ	-79.7 (5)	C2—C3—C4—C5	0.1 (14)
O4—La1—O17 ⁱⁱⁱ —La2 ⁱⁱⁱ	72.88 (19)	C3—C4—C5—C6	0.8 (15)

supplementary materials

O4—La1—O17 ⁱⁱⁱ —C8 ⁱⁱⁱ	-130.5 (5)	C4—C5—C6—C7	-0.1 (14)
O20—La1—O17 ⁱⁱⁱ —La2 ⁱⁱⁱ	-167.5 (2)	C5—C6—C7—C2	-1.5 (11)
O20—La1—O17 ⁱⁱⁱ —C8 ⁱⁱⁱ	-10.9 (5)	C5—C6—C7—C8	175.6 (8)
O22—La1—O17 ⁱⁱⁱ —La2 ⁱⁱⁱ	156.13 (19)	C2—C7—C8—O17	-110.5 (7)
O22—La1—O17 ⁱⁱⁱ —C8 ⁱⁱⁱ	-47.3 (6)	C2—C7—C8—O18	69.9 (8)
O1—La1—O5 ⁱ —La2 ⁱⁱⁱ	77.17 (18)	C6—C7—C8—O17	72.4 (9)
O1—La1—O5 ⁱ —C10 ⁱ	-118.7 (5)	C6—C7—C8—O18	-107.2 (8)
O2—La1—O5 ⁱ —La2 ⁱⁱⁱ	129.6 (2)	O3—C9—C12—C11	-145.8 (8)
O2—La1—O5 ⁱ —C10 ⁱ	-66.3 (5)	O3—C9—C12—C13	32.4 (12)
O3—La1—O5 ⁱ —La2 ⁱⁱⁱ	-59.2 (2)	O4—C9—C12—C11	30.5 (10)
O3—La1—O5 ⁱ —C10 ⁱ	104.9 (5)	O4—C9—C12—C13	-151.3 (8)
O4—La1—O5 ⁱ —La2 ⁱⁱⁱ	-79.85 (18)	O5—C10—C11—C12	-99.6 (8)
O4—La1—O5 ⁱ —C10 ⁱ	84.3 (5)	O5—C10—C11—C16	85.3 (9)
O20—La1—O5 ⁱ —La2 ⁱⁱⁱ	20.6 (3)	O6—C10—C11—C12	82.9 (9)
O20—La1—O5 ⁱ —C10 ⁱ	-175.3 (5)	O6—C10—C11—C16	-92.3 (8)
O22—La1—O5 ⁱ —La2 ⁱⁱⁱ	-143.0 (3)	C10—C11—C12—C9	1.6 (11)
O22—La1—O5 ⁱ —C10 ⁱ	21.1 (7)	C10—C11—C12—C13	-176.6 (7)
O17—La2—O1—La1	127.4 (2)	C16—C11—C12—C9	176.8 (7)
O17—La2—O1—C1	-79.6 (6)	C16—C11—C12—C13	-1.5 (12)
O19—La2—O1—La1	-34.3 (2)	C10—C11—C16—C15	178.2 (8)
O19—La2—O1—C1	118.7 (6)	C12—C11—C16—C15	3.0 (12)
O5 ^{iv} —La2—O1—La1	89.9 (2)	C9—C12—C13—C14	-177.9 (9)
O5 ^{iv} —La2—O1—C1	-117.2 (5)	C11—C12—C13—C14	0.3 (13)
O8 ^v —La2—O1—La1	-107.6 (2)	C12—C13—C14—C15	-0.6 (16)
O8 ^v —La2—O1—C1	45.4 (5)	C13—C14—C15—C16	2.3 (16)
O6 ⁱⁱⁱ —La2—O1—La1	-168.64 (18)	C14—C15—C16—C11	-3.4 (15)
O6 ⁱⁱⁱ —La2—O1—C1	-15.7 (6)	O7—C17—C18—C19	-128.0 (7)
O18 ⁱⁱⁱ —La2—O1—La1	36.59 (19)	O7—C17—C18—C23	43.1 (9)
O18 ⁱⁱⁱ —La2—O1—C1	-170.4 (6)	O8—C17—C18—C19	46.5 (9)
O1—La2—O17—C8	13.8 (4)	O8—C17—C18—C23	-142.4 (6)
O1—La2—O17—La1 ⁱⁱⁱ	-145.6 (2)	C17—C18—C19—C20	169.6 (7)
O19—La2—O17—C8	53.0 (5)	C23—C18—C19—C20	-1.8 (11)
O19—La2—O17—La1 ⁱⁱⁱ	-106.4 (3)	C17—C18—C23—C22	-169.5 (7)
O4 ^{iv} —La2—O17—C8	-166.7 (4)	C17—C18—C23—C24	12.1 (10)
O4 ^{iv} —La2—O17—La1 ⁱⁱⁱ	33.9 (2)	C19—C18—C23—C22	1.5 (10)
O5 ^{iv} —La2—O17—C8	159.1 (5)	C19—C18—C23—C24	-176.9 (6)
O5 ^{iv} —La2—O17—La1 ⁱⁱⁱ	-0.36 (16)	C18—C19—C20—C21	1.4 (13)
O8 ^v —La2—O17—C8	-44.7 (4)	C19—C20—C21—C22	-0.6 (14)
O8 ^v —La2—O17—La1 ⁱⁱⁱ	155.92 (17)	C20—C21—C22—C23	0.2 (14)
O6 ⁱⁱⁱ —La2—O17—C8	-99.1 (4)	C21—C22—C23—C18	-0.7 (12)
O6 ⁱⁱⁱ —La2—O17—La1 ⁱⁱⁱ	101.5 (2)	C21—C22—C23—C24	177.7 (8)
O18 ⁱⁱⁱ —La2—O17—C8	92.1 (4)	C18—C23—C24—O19	46.1 (9)

O18 ⁱⁱⁱ —La2—O17—La1 ⁱⁱⁱ	-67.35 (19)	C18—C23—C24—O20	-134.2 (7)
O1—La2—O19—C24	52.5 (7)	C22—C23—C24—O19	-132.3 (7)
O17—La2—O19—C24	15.4 (9)	C22—C23—C24—O20	47.4 (10)
O4 ^{iv} —La2—O19—C24	-131.9 (7)		

Symmetry codes: (i) $-x+1, -y+1, -z+1$; (ii) $-x, -y, -z+1$; (iii) $-x, -y+1, -z+1$; (iv) $x-1, y, z$; (v) $-x-1, -y, -z+1$; (vi) $x+1, y, z$.

Hydrogen-bond geometry (\AA , $^\circ$)

Cg is the centroid of the C2–C7 ring.

$D-H\cdots A$	$D-H$	$H\cdots A$	$D\cdots A$	$D-H\cdots A$
O22—H22A \cdots O1 ⁱⁱ	0.90	2.20	3.001 (6)	148.
O22—H22B \cdots O20 ⁱⁱ	0.88	2.15	3.009 (8)	164.
O27—H27A \cdots O2 ^{vii}	0.91	2.38	3.17 (2)	146.00
O27—H27B \cdots O3 ⁱ	0.88	1.86	2.69 (2)	158.00
C3—H3 \cdots O27 ^{viii}	0.93	2.15	2.96 (2)	145.
C16—H16 \cdots O2 ⁱ	0.93	2.58	3.331 (10)	138.
C19—H19 \cdots Cg ^v	0.93	2.98	3.902 (9)	169.

Symmetry codes: (ii) $-x, -y, -z+1$; (vii) $x+1, y+1, z$; (i) $-x+1, -y+1, -z+1$; (viii) $x-1, y-1, z$; (v) $-x-1, -y, -z+1$.

Fig. 1




Fig. 2

