

{5,5'-Dimethoxy-2,2'-[1,1'-(2,2-dimethylpropane-1,3-diylidinitrilo)diethylidyne]-diphenolato- $\kappa^4 O, N, N', O'$ }copper(II) monohydrate

Akbar Ghaemi,^a ‡ Saeed Rayati,^b Ehsan Elahi,^a Seik Weng Ng^{c,d} and Edward R. T. Tiekink^{c*}

^aDepartment of Chemistry, Saveh Branch, Islamic Azad University, Saveh, Iran,

^bDepartment of Chemistry, K. N. Toosi University of Technology, PO Box, 16315-1618, Tehran, Iran, ^cDepartment of Chemistry, University of Malaya, 50603 Kuala Lumpur, Malaysia, and ^dChemistry Department, Faculty of Science, King

Abdulaziz University, PO Box 80203 Jeddah, Saudi Arabia

Correspondence e-mail: Edward.Tiekink@gmail.com


Received 21 September 2011; accepted 22 September 2011

Key indicators: single-crystal X-ray study; $T = 294$ K; mean $\sigma(\text{C}-\text{C}) = 0.004$ Å; R factor = 0.037; wR factor = 0.105; data-to-parameter ratio = 17.7.

The tetradentate dianion in the title complex hydrate, $[\text{Cu}(\text{C}_{23}\text{H}_{28}\text{N}_2\text{O}_4)] \cdot \text{H}_2\text{O}$, provides the Cu^{II} atom with a *cis*- N_2O_2 donor set. There is a significant twist from a regular square-planar geometry with the dihedral angle formed between the two six-membered CuOC_3N chelate rings being $32.14(8)^\circ$. The water molecule forms hydrogen bonds to each of the coordinating O atoms of a given complex molecule. Supramolecular layers in the *bc* plane are formed in the crystal packing through $\text{C}-\text{H} \cdots \text{O}$ and $\text{C}-\text{H} \cdots \pi$ interactions.

Related literature

For the catalytic potential of Schiff base complexes of Cu^{II} , see: Gupta & Sutar (2008); Rayati *et al.* (2010). For the structure of the ligand, see: Ghaemi *et al.* (2011). For crystallization conditions, see: Harrowfield *et al.* (1996).



‡ Additional correspondence author, e-mail: akbarghaemi@yahoo.com.

Experimental

Crystal data

$[\text{Cu}(\text{C}_{23}\text{H}_{28}\text{N}_2\text{O}_4)] \cdot \text{H}_2\text{O}$

$M_r = 478.03$

Triclinic, $P\bar{1}$

$a = 10.4721(7)$ Å

$b = 10.8023(9)$ Å

$c = 10.8487(7)$ Å

$\alpha = 106.699(7)^\circ$

$\beta = 99.823(5)^\circ$

$\gamma = 100.035(6)^\circ$

$V = 1125.37(14)$ Å³

$Z = 2$

Mo $K\alpha$ radiation

$\mu = 1.01$ mm⁻¹

$T = 294$ K

$0.40 \times 0.40 \times 0.20$ mm

Data collection

Agilent SuperNova Dual diffractometer with Atlas detector

Absorption correction: multi-scan (*CrysAlis PRO*; Agilent, 2010)

$T_{\text{min}} = 0.643$, $T_{\text{max}} = 1.000$

11143 measured reflections

5034 independent reflections

4332 reflections with $I > 2\sigma(I)$

$R_{\text{int}} = 0.024$

Refinement

$R[F^2 > 2\sigma(F^2)] = 0.037$

$wR(F^2) = 0.105$

$S = 0.99$

5034 reflections

285 parameters

6 restraints

H-atom parameters constrained

$\Delta\rho_{\text{max}} = 0.26$ e Å⁻³

$\Delta\rho_{\text{min}} = -0.47$ e Å⁻³

Table 1

Selected bond lengths (Å).

Cu—O2	1.8825 (16)	Cu—N1	1.9597 (17)
Cu—O3	1.8776 (15)	Cu—N2	1.9524 (18)

Table 2

Hydrogen-bond geometry (Å, °).

$D-H \cdots A$	$D-H$	$H \cdots A$	$D \cdots A$	$D-H \cdots A$
O1w—H1w ⁱ ···O2	0.84	2.12	2.832 (3)	142
O1w—H2w ⁱ ···O3	0.84	2.32	3.035 (3)	143
C7—H7c···O1w ⁱ	0.96	2.55	3.476 (5)	163
C16—H16c···O2 ⁱⁱ	0.96	2.52	3.409 (3)	153
C14—H14b···Cg1 ⁱⁱ	0.97	2.62	3.426 (2)	141

Symmetry codes: (i) $-x + 1, -y, -z$; (ii) $-x + 1, -y + 1, -z + 1$.

Data collection: *CrysAlis PRO* (Agilent, 2010); cell refinement: *CrysAlis PRO*; data reduction: *CrysAlis PRO*; program(s) used to solve structure: *SHELXS97* (Sheldrick, 2008); program(s) used to refine structure: *SHELXL97* (Sheldrick, 2008); molecular graphics: *ORTEP-3* (Farrugia, 1997) and *DIAMOND* (Brandenburg, 2006); software used to prepare material for publication: *publCIF* (Westrip, 2010).

We gratefully acknowledge practical support of this study by K. N. Toosi University of Technology, Islamic Azad University (Saveh Branch), and thank the University of Malaya for supporting the crystallographic facility.

Supplementary data and figures for this paper are available from the IUCr electronic archives (Reference: HG5100).

References

- Agilent (2010). *CrysAlis PRO*. Agilent Technologies, Yarnton, England.
- Brandenburg, K. (2006). *DIAMOND*. Crystal Impact GbR, Bonn, Germany.
- Farrugia, L. J. (1997). *J. Appl. Cryst.* **30**, 565.
- Ghaemi, A., Rayati, S., Elahi, E., Ng, S. W. & Tiekink, E. R. T. (2011). *Acta Cryst.* **E67**, o2760.
- Gupta, K. C. & Sutar, A. K. (2008). *Coord. Chem. Rev.* **252**, 1420–1450.
- Harrowfield, J. M., Miyamae, H., Skelton, B. W., Soudi, A. A. & White, A. H. (1996). *Aust. J. Chem.* **49**, 1165–1169.
- Rayati, S., Zakavi, S., Koliaei, M., Wojtczak, A. & Kozakiewicz, A. (2010). *Inorg. Chem. Commun.* **13**, 203–207.
- Sheldrick, G. M. (2008). *Acta Cryst.* **A64**, 112–122.
- Westrip, S. P. (2010). *J. Appl. Cryst.* **43**, 920–925.

supplementary materials

Acta Cryst. (2011). E67, m1445-m1446 [doi:10.1107/S160053681103889X]

{5,5'-Dimethoxy-2,2'-[1,1'-(2,2-dimethylpropane-1,3-diyl)dinitrilo]diethyldiynediethyldiphenolato- κ^4 O,N,N',O'}copper(II) monohydrate

A. Ghaemi, S. Rayati, E. Elahi, S. W. Ng and E. R. T. Tiekink

Comment

Synthetic copper(II) Schiff base complexes have long been of great interest because of their potential as catalysts in the oxidation of various organic compounds (Gupta & Sutar, 2008). In continuation of research in this field (Rayati *et al.*, 2010), the title complex, (I), was investigated.

The tetradentate dianion in the title monohydrate, (I), Fig. 1, provides a *cis*-N₂O₂ donor set, Table 1. Three six-membered chelate rings are formed as a result of coordination of the dianion. The CuNC₃N ring adopts a half-chair conformation. While the CuOC₃N chelate ring containing the O3 atom approaches planarity with a r.m.s. deviation of 0.031 Å, the other ring displays significant distortions. Thus, the r.m.s. deviation for the O2-containing CuOC₃N chelate ring is 0.163 Å with maximum deviations of 0.162 (2) Å for atom O2 and -0.159 (1) Å for the Cu atom. The dihedral angle formed between the two CuOC₃N chelate rings is 32.14 (8)° indicating a significant distortion from a regular square planar geometry. Each of the methoxy groups is co-planar with the benzene ring to which it is attached as seen in the values of the C7—O1—C3—C2 and C23—O4—C20—C19 of -0.8 (4) and -179.3 (3)°, respectively. The water molecule of solvation is associated with the complex, forming a bridge *via* its hydrogen atoms between the two coordinated oxygen atoms, Table 2.

The crystal packing features C—H...O and C—H... π interactions, Table 2, that assemble molecules into layers in the *bc* plane, Fig. 2, which stack along the *a* axis, Fig. 3.

Experimental

The title complex was obtained by the template method in a branch tube (Harrowfield *et al.*, 1996). The recently described (Ghaemi *et al.*, 2011) *N,N*-bis(2-hydroxy-4-methoxyacetophenone)-2,2-dimethylpropane-1,3-diamine (0.40 g, 1 mmol) and copper(II) acetate monohydrate (0.199 g, 1 mmol) were placed in the main arm of a branched tube. Ethanol was added to fill both arms. The tube was sealed and the main arm immersed in an oil bath at 333 K while the other was held at ambient temperature. After one week, crystals deposited in the cooler arm. These were filtered off and air dried. Yield: 75%. FT—IR data: $\nu(\text{C}=\text{N})$ 1595 cm⁻¹.

Refinement

The H-atoms were placed in calculated positions (C—H 0.93 to 0.97 Å) and were included in the refinement in the riding model approximation, with $U_{\text{iso}}(\text{H})$ set to 1.2 to 1.5 $U_{\text{equiv}}(\text{C})$. The water-H atoms were placed in calculated positions (O—H = 0.84 Å; 1.5 $U_{\text{equiv}}(\text{O})$) on the basis of hydrogen bonding.

Figures




Fig. 1. The molecular structure of (I) showing the atom-labelling scheme and displacement ellipsoids at the 50% probability level.




Fig. 2. Supramolecular layer in the *bc* plane in (I) sustained by C—H...O and C—H... π interactions shown as blue and black dashed lines, respectively. The O—H...O hydrogen bonds are shown as orange dashed lines.




Fig. 3. A view in projection down the *c* axis of the unit-cell contents of (I), highlighting the stacking of layers along the *a* axis. The C—H...O and C—H... π interactions shown as blue and black dashed lines, respectively, and the O—H...O hydrogen bonds are shown as orange dashed lines.

{5,5'-Dimethoxy-2,2'-[1,1'-(2,2-dimethylpropane-1,3-diyl)dinitrilo] diethyldiyne]diphenolato- κ^4 O,N,N',\backslash

O'}copper(II) monohydrate

Crystal data

[Cu(C₂₃H₂₈N₂O₄)]·H₂O

M_r = 478.03

Triclinic, *P* $\bar{1}$

Hall symbol: -P 1

a = 10.4721 (7) Å

b = 10.8023 (9) Å

c = 10.8487 (7) Å

α = 106.699 (7)°

β = 99.823 (5)°

γ = 100.035 (6)°

V = 1125.37 (14) Å³

Z = 2

F(000) = 502

D_x = 1.411 Mg m⁻³

Mo *K* α radiation, λ = 0.71073 Å

Cell parameters from 5694 reflections

θ = 2.3–29.3°

μ = 1.01 mm⁻¹

T = 294 K

Block, dark-brown

0.40 × 0.40 × 0.20 mm

Data collection

Agilent SuperNova Dual
diffractometer with Atlas detector

Radiation source: SuperNova (Mo) X-ray Source

5034 independent reflections

4332 reflections with *I* > 2 σ (*I*)

Mirror	$R_{\text{int}} = 0.024$
Detector resolution: 10.4041 pixels mm^{-1}	$\theta_{\text{max}} = 27.5^\circ$, $\theta_{\text{min}} = 2.5^\circ$
ω scan	$h = -12 \rightarrow 13$
Absorption correction: multi-scan (<i>CrysAlis PRO</i> ; Agilent, 2010)	$k = -14 \rightarrow 13$
$T_{\text{min}} = 0.643$, $T_{\text{max}} = 1.000$	$l = -14 \rightarrow 11$
11143 measured reflections	

Refinement

Refinement on F^2	Primary atom site location: structure-invariant direct methods
Least-squares matrix: full	Secondary atom site location: difference Fourier map
$R[F^2 > 2\sigma(F^2)] = 0.037$	Hydrogen site location: inferred from neighbouring sites
$wR(F^2) = 0.105$	H-atom parameters constrained
$S = 0.99$	$w = 1/[\sigma^2(F_o^2) + (0.0531P)^2 + 0.3861P]$
5034 reflections	where $P = (F_o^2 + 2F_c^2)/3$
285 parameters	$(\Delta/\sigma)_{\text{max}} = 0.002$
6 restraints	$\Delta\rho_{\text{max}} = 0.26 \text{ e } \text{\AA}^{-3}$
	$\Delta\rho_{\text{min}} = -0.47 \text{ e } \text{\AA}^{-3}$

Special details

Geometry. All e.s.d.'s (except the e.s.d. in the dihedral angle between two l.s. planes) are estimated using the full covariance matrix. The cell e.s.d.'s are taken into account individually in the estimation of e.s.d.'s in distances, angles and torsion angles; correlations between e.s.d.'s in cell parameters are only used when they are defined by crystal symmetry. An approximate (isotropic) treatment of cell e.s.d.'s is used for estimating e.s.d.'s involving l.s. planes.

Refinement. Refinement of F^2 against ALL reflections. The weighted R -factor wR and goodness of fit S are based on F^2 , conventional R -factors R are based on F , with F set to zero for negative F^2 . The threshold expression of $F^2 > \sigma(F^2)$ is used only for calculating R -factors(gt) *etc.* and is not relevant to the choice of reflections for refinement. R -factors based on F^2 are statistically about twice as large as those based on F , and R -factors based on ALL data will be even larger.

Fractional atomic coordinates and isotropic or equivalent isotropic displacement parameters (\AA^2)

	x	y	z	$U_{\text{iso}}^*/U_{\text{eq}}$
Cu	0.48963 (3)	0.47337 (3)	0.24182 (2)	0.04175 (11)
O1	0.04084 (19)	-0.0426 (2)	-0.1635 (2)	0.0746 (6)
O2	0.40205 (16)	0.30214 (16)	0.12419 (15)	0.0519 (4)
O3	0.62185 (18)	0.39699 (17)	0.30882 (16)	0.0580 (5)
O4	0.96607 (19)	0.3235 (2)	0.60237 (19)	0.0702 (5)
O1w	0.5646 (3)	0.1173 (3)	0.1144 (3)	0.1143 (10)
H1w	0.4956	0.1403	0.0867	0.171*
H2w	0.6121	0.1815	0.1794	0.171*
N1	0.40191 (19)	0.55577 (19)	0.12292 (18)	0.0440 (4)
N2	0.54285 (18)	0.63414 (17)	0.39652 (18)	0.0404 (4)
C1	0.2881 (2)	0.2697 (2)	0.0363 (2)	0.0429 (5)

supplementary materials

C2	0.2249 (2)	0.1342 (2)	-0.0152 (2)	0.0472 (5)
H2	0.2638	0.0743	0.0155	0.057*
C3	0.1068 (2)	0.0873 (3)	-0.1098 (2)	0.0553 (6)
C4	0.0476 (3)	0.1766 (3)	-0.1551 (3)	0.0685 (8)
H4	-0.0329	0.1462	-0.2182	0.082*
C5	0.1076 (3)	0.3079 (3)	-0.1070 (3)	0.0607 (7)
H5	0.0664	0.3657	-0.1392	0.073*
C6	0.2299 (2)	0.3623 (2)	-0.0100 (2)	0.0461 (5)
C7	0.0998 (3)	-0.1364 (3)	-0.1189 (3)	0.0780 (9)
H7A	0.0449	-0.2245	-0.1647	0.117*
H7B	0.1071	-0.1161	-0.0255	0.117*
H7C	0.1869	-0.1317	-0.1367	0.117*
C8	0.2965 (2)	0.5025 (3)	0.0267 (2)	0.0481 (6)
C9	0.2384 (3)	0.5809 (3)	-0.0542 (3)	0.0696 (8)
H9A	0.2902	0.6711	-0.0212	0.104*
H9B	0.1480	0.5795	-0.0475	0.104*
H9C	0.2401	0.5420	-0.1452	0.104*
C10	0.4828 (3)	0.6911 (2)	0.1535 (2)	0.0520 (6)
H10A	0.4573	0.7230	0.0801	0.062*
H10B	0.5759	0.6882	0.1623	0.062*
C11	0.4675 (3)	0.7898 (2)	0.2811 (3)	0.0511 (6)
C12	0.3407 (3)	0.8406 (3)	0.2542 (3)	0.0714 (8)
H12A	0.3482	0.8891	0.1932	0.107*
H12B	0.3301	0.8980	0.3358	0.107*
H12C	0.2646	0.7665	0.2170	0.107*
C13	0.5910 (3)	0.9054 (3)	0.3299 (3)	0.0732 (8)
H13A	0.5994	0.9441	0.2615	0.110*
H13B	0.6688	0.8732	0.3519	0.110*
H13C	0.5822	0.9713	0.4070	0.110*
C14	0.4513 (2)	0.7208 (2)	0.3848 (2)	0.0456 (5)
H14A	0.4652	0.7883	0.4703	0.055*
H14B	0.3604	0.6681	0.3623	0.055*
C15	0.6363 (2)	0.6611 (2)	0.5039 (2)	0.0434 (5)
C16	0.6651 (3)	0.7919 (2)	0.6149 (3)	0.0632 (7)
H16A	0.6322	0.8562	0.5812	0.095*
H16B	0.7596	0.8230	0.6508	0.095*
H16C	0.6218	0.7794	0.6831	0.095*
C17	0.7159 (2)	0.5667 (2)	0.5215 (2)	0.0423 (5)
C18	0.8061 (3)	0.5949 (3)	0.6445 (2)	0.0564 (6)
H18	0.8115	0.6730	0.7120	0.068*
C19	0.8854 (3)	0.5134 (3)	0.6686 (3)	0.0635 (7)
H19	0.9418	0.5351	0.7516	0.076*
C20	0.8819 (2)	0.3976 (3)	0.5691 (2)	0.0508 (6)
C21	0.7941 (2)	0.3632 (2)	0.4486 (2)	0.0459 (5)
H21	0.7917	0.2854	0.3822	0.055*
C22	0.7074 (2)	0.4443 (2)	0.4242 (2)	0.0425 (5)
C23	0.9685 (3)	0.2051 (4)	0.5051 (3)	0.0800 (9)
H23A	1.0308	0.1625	0.5422	0.120*
H23B	0.9951	0.2259	0.4315	0.120*

H23C 0.8812 0.1465 0.4754 0.120*

Atomic displacement parameters (\AA^2)

	U^{11}	U^{22}	U^{33}	U^{12}	U^{13}	U^{23}
Cu	0.04730 (18)	0.04062 (18)	0.03637 (16)	0.01857 (13)	0.00166 (12)	0.01100 (12)
O1	0.0582 (11)	0.0691 (13)	0.0700 (12)	0.0033 (10)	-0.0159 (10)	0.0074 (10)
O2	0.0550 (9)	0.0437 (9)	0.0456 (9)	0.0209 (8)	-0.0136 (7)	0.0063 (7)
O3	0.0669 (11)	0.0539 (10)	0.0417 (8)	0.0319 (9)	-0.0107 (8)	0.0014 (7)
O4	0.0647 (12)	0.0861 (14)	0.0622 (11)	0.0343 (11)	-0.0056 (9)	0.0297 (11)
O1w	0.0964 (18)	0.0788 (16)	0.153 (2)	0.0434 (14)	-0.0017 (17)	0.0207 (17)
N1	0.0504 (10)	0.0491 (11)	0.0414 (9)	0.0236 (9)	0.0130 (8)	0.0198 (8)
N2	0.0443 (10)	0.0368 (9)	0.0434 (9)	0.0130 (8)	0.0108 (8)	0.0154 (8)
C1	0.0449 (12)	0.0550 (13)	0.0294 (9)	0.0224 (10)	0.0044 (9)	0.0110 (9)
C2	0.0471 (12)	0.0536 (14)	0.0371 (11)	0.0195 (11)	0.0007 (9)	0.0099 (10)
C3	0.0486 (13)	0.0643 (16)	0.0437 (12)	0.0144 (12)	0.0010 (10)	0.0087 (12)
C4	0.0521 (15)	0.087 (2)	0.0535 (15)	0.0173 (15)	-0.0137 (12)	0.0185 (15)
C5	0.0559 (15)	0.0807 (19)	0.0504 (14)	0.0302 (14)	-0.0010 (12)	0.0283 (14)
C6	0.0480 (12)	0.0607 (15)	0.0348 (10)	0.0256 (11)	0.0063 (9)	0.0178 (10)
C7	0.0662 (18)	0.0586 (17)	0.086 (2)	0.0091 (15)	-0.0071 (16)	0.0059 (16)
C8	0.0545 (13)	0.0627 (15)	0.0400 (11)	0.0308 (12)	0.0133 (10)	0.0251 (11)
C9	0.0762 (19)	0.079 (2)	0.0666 (17)	0.0324 (16)	0.0040 (14)	0.0419 (16)
C10	0.0565 (14)	0.0581 (15)	0.0553 (14)	0.0217 (12)	0.0194 (11)	0.0313 (12)
C11	0.0610 (14)	0.0421 (12)	0.0596 (14)	0.0211 (11)	0.0165 (12)	0.0242 (11)
C12	0.087 (2)	0.0629 (17)	0.0770 (19)	0.0444 (16)	0.0179 (16)	0.0274 (15)
C13	0.084 (2)	0.0560 (16)	0.081 (2)	0.0073 (15)	0.0165 (17)	0.0320 (15)
C14	0.0522 (13)	0.0409 (12)	0.0482 (12)	0.0187 (10)	0.0169 (10)	0.0136 (10)
C15	0.0487 (12)	0.0373 (11)	0.0406 (11)	0.0041 (10)	0.0098 (10)	0.0113 (9)
C16	0.086 (2)	0.0412 (13)	0.0514 (14)	0.0139 (13)	0.0024 (13)	0.0066 (11)
C17	0.0414 (11)	0.0394 (11)	0.0411 (11)	0.0037 (9)	0.0021 (9)	0.0136 (9)
C18	0.0568 (14)	0.0482 (14)	0.0470 (13)	0.0036 (12)	-0.0092 (11)	0.0069 (11)
C19	0.0554 (15)	0.0667 (17)	0.0531 (14)	0.0070 (13)	-0.0161 (12)	0.0173 (13)
C20	0.0406 (12)	0.0599 (15)	0.0524 (13)	0.0119 (11)	0.0003 (10)	0.0252 (12)
C21	0.0434 (12)	0.0536 (14)	0.0422 (11)	0.0174 (10)	0.0060 (9)	0.0169 (10)
C22	0.0400 (11)	0.0478 (12)	0.0378 (11)	0.0098 (10)	0.0018 (9)	0.0157 (9)
C23	0.082 (2)	0.099 (2)	0.077 (2)	0.056 (2)	0.0146 (17)	0.0381 (19)

Geometric parameters (\AA , $^\circ$)

Cu—O2	1.8825 (16)	C9—H9C	0.9600
Cu—O3	1.8776 (15)	C10—C11	1.538 (3)
Cu—N1	1.9597 (17)	C10—H10A	0.9700
Cu—N2	1.9524 (18)	C10—H10B	0.9700
O1—C3	1.358 (3)	C11—C13	1.528 (4)
O1—C7	1.428 (3)	C11—C14	1.535 (3)
O2—C1	1.316 (3)	C11—C12	1.538 (3)
O3—C22	1.312 (3)	C12—H12A	0.9600
O4—C20	1.361 (3)	C12—H12B	0.9600
O4—C23	1.412 (4)	C12—H12C	0.9600

supplementary materials

O1w—H1w	0.8400	C13—H13A	0.9600
O1w—H2w	0.8400	C13—H13B	0.9600
N1—C8	1.296 (3)	C13—H13C	0.9600
N1—C10	1.469 (3)	C14—H14A	0.9700
N2—C15	1.310 (3)	C14—H14B	0.9700
N2—C14	1.466 (3)	C15—C17	1.458 (3)
C1—C2	1.400 (3)	C15—C16	1.512 (3)
C1—C6	1.421 (3)	C16—H16A	0.9600
C2—C3	1.374 (3)	C16—H16B	0.9600
C2—H2	0.9300	C16—H16C	0.9600
C3—C4	1.390 (4)	C17—C22	1.412 (3)
C4—C5	1.353 (4)	C17—C18	1.414 (3)
C4—H4	0.9300	C18—C19	1.359 (4)
C5—C6	1.420 (3)	C18—H18	0.9300
C5—H5	0.9300	C19—C20	1.387 (4)
C6—C8	1.459 (4)	C19—H19	0.9300
C7—H7A	0.9600	C20—C21	1.373 (3)
C7—H7B	0.9600	C21—C22	1.411 (3)
C7—H7C	0.9600	C21—H21	0.9300
C8—C9	1.511 (3)	C23—H23A	0.9600
C9—H9A	0.9600	C23—H23B	0.9600
C9—H9B	0.9600	C23—H23C	0.9600
O3—Cu—O2	87.70 (7)	C13—C11—C10	107.4 (2)
O3—Cu—N2	93.30 (7)	C14—C11—C10	110.41 (18)
O2—Cu—N2	161.99 (8)	C13—C11—C12	110.3 (2)
O3—Cu—N1	156.09 (8)	C14—C11—C12	106.3 (2)
O2—Cu—N1	91.13 (7)	C10—C11—C12	110.7 (2)
N2—Cu—N1	95.08 (8)	C11—C12—H12A	109.5
C3—O1—C7	117.2 (2)	C11—C12—H12B	109.5
C1—O2—Cu	126.53 (14)	H12A—C12—H12B	109.5
C22—O3—Cu	128.03 (15)	C11—C12—H12C	109.5
C20—O4—C23	118.3 (2)	H12A—C12—H12C	109.5
H1w—O1w—H2w	107.4	H12B—C12—H12C	109.5
C8—N1—C10	123.47 (19)	C11—C13—H13A	109.5
C8—N1—Cu	128.32 (17)	C11—C13—H13B	109.5
C10—N1—Cu	108.06 (14)	H13A—C13—H13B	109.5
C15—N2—C14	121.92 (19)	C11—C13—H13C	109.5
C15—N2—Cu	127.82 (15)	H13A—C13—H13C	109.5
C14—N2—Cu	109.93 (14)	H13B—C13—H13C	109.5
O2—C1—C2	116.13 (19)	N2—C14—C11	114.24 (18)
O2—C1—C6	124.1 (2)	N2—C14—H14A	108.7
C2—C1—C6	119.7 (2)	C11—C14—H14A	108.7
C3—C2—C1	121.7 (2)	N2—C14—H14B	108.7
C3—C2—H2	119.1	C11—C14—H14B	108.7
C1—C2—H2	119.1	H14A—C14—H14B	107.6
O1—C3—C2	124.5 (2)	N2—C15—C17	121.6 (2)
O1—C3—C4	116.1 (2)	N2—C15—C16	120.9 (2)
C2—C3—C4	119.4 (3)	C17—C15—C16	117.5 (2)
C5—C4—C3	119.7 (2)	C15—C16—H16A	109.5

C5—C4—H4	120.1	C15—C16—H16B	109.5
C3—C4—H4	120.1	H16A—C16—H16B	109.5
C4—C5—C6	123.6 (2)	C15—C16—H16C	109.5
C4—C5—H5	118.2	H16A—C16—H16C	109.5
C6—C5—H5	118.2	H16B—C16—H16C	109.5
C5—C6—C1	115.8 (2)	C22—C17—C18	116.3 (2)
C5—C6—C8	120.7 (2)	C22—C17—C15	124.43 (19)
C1—C6—C8	123.2 (2)	C18—C17—C15	119.3 (2)
O1—C7—H7A	109.5	C19—C18—C17	123.1 (2)
O1—C7—H7B	109.5	C19—C18—H18	118.5
H7A—C7—H7B	109.5	C17—C18—H18	118.5
O1—C7—H7C	109.5	C18—C19—C20	119.9 (2)
H7A—C7—H7C	109.5	C18—C19—H19	120.1
H7B—C7—H7C	109.5	C20—C19—H19	120.1
N1—C8—C6	121.1 (2)	O4—C20—C21	124.7 (2)
N1—C8—C9	122.0 (2)	O4—C20—C19	115.5 (2)
C6—C8—C9	116.9 (2)	C21—C20—C19	119.7 (2)
C8—C9—H9A	109.4	C20—C21—C22	120.9 (2)
C8—C9—H9B	109.4	C20—C21—H21	119.6
H9A—C9—H9B	109.5	C22—C21—H21	119.6
C8—C9—H9C	109.6	O3—C22—C21	115.4 (2)
H9A—C9—H9C	109.5	O3—C22—C17	124.6 (2)
H9B—C9—H9C	109.5	C21—C22—C17	119.95 (19)
N1—C10—C11	113.23 (19)	O4—C23—H23A	109.5
N1—C10—H10A	108.9	O4—C23—H23B	109.5
C11—C10—H10A	108.9	H23A—C23—H23B	109.5
N1—C10—H10B	108.9	O4—C23—H23C	109.5
C11—C10—H10B	108.9	H23A—C23—H23C	109.5
H10A—C10—H10B	107.7	H23B—C23—H23C	109.5
C13—C11—C14	111.8 (2)		
O3—Cu—O2—C1	178.8 (2)	C5—C6—C8—N1	174.0 (2)
N2—Cu—O2—C1	85.3 (3)	C1—C6—C8—N1	-13.1 (3)
N1—Cu—O2—C1	-25.1 (2)	C5—C6—C8—C9	-7.1 (3)
O2—Cu—O3—C22	-158.8 (2)	C1—C6—C8—C9	165.8 (2)
N2—Cu—O3—C22	3.2 (2)	C8—N1—C10—C11	108.3 (3)
N1—Cu—O3—C22	113.6 (2)	Cu—N1—C10—C11	-76.0 (2)
O3—Cu—N1—C8	104.7 (3)	N1—C10—C11—C13	158.1 (2)
O2—Cu—N1—C8	17.8 (2)	N1—C10—C11—C14	36.0 (3)
N2—Cu—N1—C8	-145.2 (2)	N1—C10—C11—C12	-81.4 (2)
O3—Cu—N1—C10	-70.8 (2)	C15—N2—C14—C11	113.8 (2)
O2—Cu—N1—C10	-157.67 (15)	Cu—N2—C14—C11	-72.3 (2)
N2—Cu—N1—C10	39.26 (15)	C13—C11—C14—N2	-75.8 (3)
O3—Cu—N2—C15	-2.6 (2)	C10—C11—C14—N2	43.7 (3)
O2—Cu—N2—C15	90.1 (3)	C12—C11—C14—N2	163.8 (2)
N1—Cu—N2—C15	-160.20 (19)	C14—N2—C15—C17	172.1 (2)
O3—Cu—N2—C14	-176.04 (14)	Cu—N2—C15—C17	-0.6 (3)
O2—Cu—N2—C14	-83.4 (2)	C14—N2—C15—C16	-7.6 (3)
N1—Cu—N2—C14	26.38 (15)	Cu—N2—C15—C16	179.70 (18)
Cu—O2—C1—C2	-163.66 (16)	N2—C15—C17—C22	4.5 (4)

supplementary materials

Cu—O2—C1—C6	18.1 (3)	C16—C15—C17—C22	-175.7 (2)
O2—C1—C2—C3	-178.5 (2)	N2—C15—C17—C18	-173.7 (2)
C6—C1—C2—C3	-0.2 (3)	C16—C15—C17—C18	6.0 (3)
C7—O1—C3—C2	-0.8 (4)	C22—C17—C18—C19	2.6 (4)
C7—O1—C3—C4	179.8 (3)	C15—C17—C18—C19	-179.0 (2)
C1—C2—C3—O1	-180.0 (2)	C17—C18—C19—C20	1.4 (4)
C1—C2—C3—C4	-0.6 (4)	C23—O4—C20—C21	2.8 (4)
O1—C3—C4—C5	-179.7 (3)	C23—O4—C20—C19	-179.3 (3)
C2—C3—C4—C5	0.9 (4)	C18—C19—C20—O4	179.2 (2)
C3—C4—C5—C6	-0.4 (5)	C18—C19—C20—C21	-2.9 (4)
C4—C5—C6—C1	-0.3 (4)	O4—C20—C21—C22	178.0 (2)
C4—C5—C6—C8	173.1 (3)	C19—C20—C21—C22	0.3 (4)
O2—C1—C6—C5	178.8 (2)	Cu—O3—C22—C21	178.78 (16)
C2—C1—C6—C5	0.7 (3)	Cu—O3—C22—C17	-0.6 (4)
O2—C1—C6—C8	5.6 (3)	C20—C21—C22—O3	-175.6 (2)
C2—C1—C6—C8	-172.6 (2)	C20—C21—C22—C17	3.9 (4)
C10—N1—C8—C6	172.0 (2)	C18—C17—C22—O3	174.3 (2)
Cu—N1—C8—C6	-2.9 (3)	C15—C17—C22—O3	-4.0 (4)
C10—N1—C8—C9	-6.9 (4)	C18—C17—C22—C21	-5.1 (3)
Cu—N1—C8—C9	178.26 (18)	C15—C17—C22—C21	176.6 (2)

Hydrogen-bond geometry (\AA , $^\circ$)

$D-H\cdots A$	$D-H$	$H\cdots A$	$D\cdots A$	$D-H\cdots A$
O1w—H1w \cdots O2	0.84	2.12	2.832 (3)	142
O1w—H2w \cdots O3	0.84	2.32	3.035 (3)	143
C7—H7c \cdots O1w ⁱ	0.96	2.55	3.476 (5)	163
C16—H16c \cdots O2 ⁱⁱ	0.96	2.52	3.409 (3)	153
C14—H14b \cdots Cg1 ⁱⁱ	0.97	2.62	3.426 (2)	141

Symmetry codes: (i) $-x+1, -y, -z$; (ii) $-x+1, -y+1, -z+1$.

Fig. 1




Fig. 2




Fig. 3

