

Poly[tris(μ_3 -2-aminoethanesulfonato)-cobalt(II)potassium]

 Xiao-lin Li,^{a*} Jing Yu^a and Yang-Miao Ou^b

^aPhysics and Chemistry Department, Jiangxi College of Traditional Chinese Medicine, Fuzhou, Jiangxi 344000, People's Republic of China, and ^bDepartment of Chemistry and Life Science, Hechi University, Yizhou, Guangxi 546300, People's Republic of China

Correspondence e-mail: caizhou2006@126.com


Received 18 September 2011; accepted 26 September 2011

Key indicators: single-crystal X-ray study; $T = 296$ K; mean $\sigma(\text{C}-\text{C}) = 0.004$ Å; R factor = 0.024; wR factor = 0.058; data-to-parameter ratio = 16.3.

The title compound, $[\text{CoK}(\text{C}_2\text{H}_6\text{NO}_3\text{S})_3]_n$, is isotypic with its Ni^{II} analogue. The Co^{II} atom is chelated by the three taurinate ligands in a distorted octahedral geometry and in a facial manner. Each taurinate ligand bridges two K^+ ions *via* its sulfonate group, forming a three-dimensional framework. Weak $\text{N}-\text{H}\cdots\text{O}$ hydrogen bonding is observed in the crystal structure.

Related literature

For the isotypic Ni^{II} structure, see: Jiang *et al.* (2005). For the applications of taurine in medicine and biochemistry, see: Bottari & Festa (1998); Jiang *et al.* (2003). For general background to taurine complexes and their derivatives, see: Zhang & Jiang (2002); Zhong *et al.* (2003); Cai *et al.* (2004, 2006, 2011); Yang *et al.* (2010a,b). For S–O(–Co) bond lengths in bridging sulfonate groups, see: Zeng *et al.* (2009); Yang *et al.* (2010b).



Experimental

Crystal data

$[\text{CoK}(\text{C}_2\text{H}_6\text{NO}_3\text{S})_3]$	$V = 1609.6$ (5) Å ³
$M_r = 470.44$	$Z = 4$
Orthorhombic, $Pna2_1$	Mo $K\alpha$ radiation
$a = 10.6901$ (19) Å	$\mu = 1.76$ mm ⁻¹
$b = 15.669$ (3) Å	$T = 296$ K
$c = 9.6094$ (17) Å	$0.26 \times 0.22 \times 0.14$ mm

Data collection

Bruker SMART CCD area-detector diffractometer	10792 measured reflections
Absorption correction: multi-scan (SADABS; Bruker, 1999)	3386 independent reflections
$T_{\text{min}} = 0.657$, $T_{\text{max}} = 0.791$	3146 reflections with $I > 2\sigma(I)$
	$R_{\text{int}} = 0.027$

Refinement

$R[F^2 > 2\sigma(F^2)] = 0.024$	H-atom parameters constrained
$wR(F^2) = 0.058$	$\Delta\rho_{\text{max}} = 0.22$ e Å ⁻³
$S = 1.02$	$\Delta\rho_{\text{min}} = -0.33$ e Å ⁻³
3386 reflections	Absolute structure: Flack (1983),
208 parameters	1528 Friedel pairs
1 restraint	Flack parameter: 0.020 (13)

Table 1

Selected bond lengths (Å).

Co1–O1	2.1316 (18)	K1–O6 ⁱ	2.687 (2)
Co1–O7	2.1411 (18)	K1–O3 ⁱⁱ	2.7140 (19)
Co1–O4	2.142 (2)	K1–O5 ⁱⁱⁱ	2.8361 (19)
Co1–N2	2.143 (2)	K1–O2 ^{iv}	2.8522 (19)
Co1–N3	2.146 (2)	K1–O8 ^v	2.8893 (19)
Co1–N1	2.149 (2)	K1–O9	2.816 (2)

Symmetry codes: (i) $x, y, z - 1$; (ii) $-x - \frac{1}{2}, y - \frac{1}{2}, z - \frac{1}{2}$; (iii) $x - \frac{1}{2}, -y + \frac{3}{2}, z - 1$; (iv) $x - \frac{1}{2}, -y + \frac{3}{2}, z$; (v) $-x, -y + 1, z - \frac{1}{2}$.

Table 2

Hydrogen-bond geometry (Å, °).

$D-\text{H}\cdots A$	$D-\text{H}$	$\text{H}\cdots A$	$D\cdots A$	$D-\text{H}\cdots A$
N3–H3B \cdots O7 ^{vi}	0.90	2.22	3.097 (3)	163
N2–H2B \cdots O4 ^{vi}	0.90	2.40	3.166 (3)	143
N2–H2B \cdots O1 ^{vi}	0.90	2.47	3.255 (3)	146
N1–H1D \cdots O4 ^{vi}	0.90	2.55	3.423 (3)	165
N3–H3A \cdots O2	0.90	2.28	3.113 (3)	153
N2–H2A \cdots O8	0.90	2.39	3.219 (3)	152
N1–H1C \cdots O5	0.90	2.49	3.229 (3)	140

Symmetry code: (vi) $x + \frac{1}{2}, -y + \frac{3}{2}, z$.

Data collection: SMART (Bruker, 1999); cell refinement: SAINT (Bruker, 1999); data reduction: SAINT; program(s) used to solve structure: SHELXS97 (Sheldrick, 2008); program(s) used to refine structure: SHELXL97 (Sheldrick, 2008); molecular graphics: SHELXTL (Sheldrick, 2008); software used to prepare material for publication: SHELXTL.

We are grateful to the Education Department Foundation of the Guangxi Zhuang Autonomous Region of the People's Republic of China (201012MS203) and the Start-up Foundation for Advanced Talents of Hechi University (No. 2008QS-N019).

Supplementary data and figures for this paper are available from the IUCr electronic archives (Reference: BX2374).

References

- Bottari, E. & Festa, M. R. (1998). *Talanta*, **46**, 91–99.
- Bruker (1999). *SMART*, *SAINT* and *SADABS*. Bruker AXS Inc., Madison, Wisconsin, USA.
- Cai, J.-H., Jiang, Y.-M. & Ng, S. W. (2006). *Acta Cryst.* **E62**, m3059–m3061.
- Cai, J.-H., Jiang, Y.-M., Wang, X.-J. & Liu, Z.-M. (2004). *Acta Cryst.* **E60**, m1659–m1661.
- Cai, J. H., Zhong, F. & Jiang, Y. M. (2011). *Chin. J. Struct. Chem.* **30**, 743–747.
- Flack, H. D. (1983). *Acta Cryst.* **A39**, 876–881.
- Jiang, Y.-M., Cai, J.-H., Liu, Z.-M. & Liu, X.-H. (2005). *Acta Cryst.* **E61**, m878–m880.
- Jiang, Y. M., Zhang, S. H., Xu, Q. & Xiao, Y. (2003). *Acta Chim. Sin.* **61**, 573–577.
- Sheldrick, G. M. (2008). *Acta Cryst.* **A64**, 112–122.
- Yang, F., Liu, X.-H. & Zhao, C.-Q. (2010*b*). *Acta Cryst.* **E66**, m1343–m1344.
- Yang, F., Wu, Z.-H. & Cai, J.-H. (2010*a*). *Acta Cryst.* **E66**, m748.
- Zeng, J.-L., Jiang, Y.-M., Sun, L.-X., Cao, Z. & Yang, D.-W. (2009). *Acta Cryst.* **E65**, m1067–m1068.
- Zhang, S. H. & Jiang, Y. M. (2002). *Chin. J. Inorg. Chem.* **18**, 497–500.
- Zhong, F., Jiang, Y. M. & Zhang, S. H. (2003). *Chin. J. Inorg. Chem.* **6**, 559–602.

supplementary materials

Acta Cryst. (2011). E67, m1466-m1467 [doi:10.1107/S1600536811039390]

Poly[tris(μ_3 -2-aminoethanesulfonato)cobalt(II)potassium]

X. Li, J. Yu and Y.-M. Ou

Comment

Taurine, an amino acid containing sulfur, is indispensable to human beings because of its applications in medicine and biochemistry (Bottari & Festa, 1998; Jiang *et al.*, 2003). Several taurine complexes and their derivatives have recently been prepared (Zhong *et al.* (2003); Cai *et al.* (2004, 2006, 2011); Yang *et al.* (2010a,b)). We found that the taurine has manifold coordination modes. For the much less well studied of the coordination modes of the sulfonate group, the title polymeric Co^{II} complex, (I), has been prepared and its structure determined.

The coordinated modes of the title compound are similar to the previously reported Ni(II) structure (Jiang *et al.* (2005)). The molecular structure of (I) is shown in Fig. 1 and the important bond lengths are listed in Table 1. The asymmetric unit of (I) consists of one Co^{II} atom, three taurinate ligands and one K⁺ ion. The cobalt is six-coordinate with three nitrogen atoms [N(1), N(2), N(3)] and three oxygen atoms [O(1), O(4), O(7)], thus giving an octahedral configuration. The Co atom forms six-membered chelate rings (NiNC₂SO) with each taurinate ligand. This is a facial isomer. Each sulfonate group of the taurinate ligand takes part in the formation of a hydrogen bond (Table 2) with the amino group of a neighbouring ligand in the complex. The most common coordination modes of the sulfonate group are monodentate and μ_2 -bridging, while μ_3 -bridging is very rare. The coordination mode of the sulfonate group in (I) is μ_3 -bridging, which makes the S–O(–Co) bonds [1.475 (2)–1.479 (2) Å] much longer than those previously reported [S–O(–Co) 1.464 (2) and 1.456 (3) Å; Yang *et al.*, 2010b; Zeng *et al.*, 2009]. The S=O(⋯K) bonds [1.443 (2)–1.453 (2) Å] are slightly longer than the uncoordinated S=O bond in taurine [1.446 (12)–1.457 (13) Å; Zhang & Jiang, 2002]. The K atom is surrounded by six O atoms from different taurinate ligands. The title complex forms a three-dimensional structure through the K⋯O linkage. The K⋯O distances are in the range 2.678 (2)–2.889 (2) Å, suggesting weak electrostatic interactions.

Experimental

A mixture of Co(CH₃COO)₂·7H₂O (0.5 mmol, 152 mg), taurine (1.5 mmol, 187 mg), KOH (1.5 mmol, 84 mg) and anhydrous methanol (15.0 ml) was placed in a Teflon-lined stainless steel vessel, and heated directly to 120 °C. After keeping at 120 °C for 4 days, it was cooled to room temperature at a rate for 10 °C/h. Block red crystals of the complex were obtained.

Refinement

H atoms were positioned geometrically (C–H = 0.97 Å and N–H = 0.90 Å) and included in the refinement in the riding model approximation, with $U_{\text{iso}}(\text{H}) = 1.2U_{\text{eq}}(\text{carrier atom})$.

Figures




Fig. 1. Molecular structure of the title compound. H atoms have been omitted for clarity. [Symmetry code: (i) $x, y, z - 1$; (ii) $-x - 1/2, y - 1/2, z - 1/2$; (iii) $x - 1/2, -y + 3/2, z - 1$; (iv) $x - 1/2, -y + 3/2, z$; (v) $-x, -y + 1, z - 1/2$.]




Fig. 2. View of a three-dimensional supramolecular structure.

Poly[tris(μ_3 -2-aminoethanesulfonato)cobalt(II)potassium]

Crystal data

[CoK(C₂H₆NO₃S)₃]

$M_r = 470.44$

Orthorhombic, $Pna2_1$

Hall symbol: P 2c -2n

$a = 10.6901 (19) \text{ \AA}$

$b = 15.669 (3) \text{ \AA}$

$c = 9.6094 (17) \text{ \AA}$

$V = 1609.6 (5) \text{ \AA}^3$

$Z = 4$

$F(000) = 964$

$D_x = 1.941 \text{ Mg m}^{-3}$

Mo $K\alpha$ radiation, $\lambda = 0.71073 \text{ \AA}$

Cell parameters from 7233 reflections

$\theta = 2.5\text{--}28.1^\circ$

$\mu = 1.76 \text{ mm}^{-1}$

$T = 296 \text{ K}$

Block, red

$0.26 \times 0.22 \times 0.14 \text{ mm}$

Data collection

Bruker SMART CCD area-detector diffractometer

Radiation source: fine-focus sealed tube graphite

φ and ω scans

Absorption correction: multi-scan (SADABS; Bruker, 1999)

$T_{\min} = 0.657, T_{\max} = 0.791$

10792 measured reflections

3386 independent reflections

3146 reflections with $I > 2\sigma(I)$

$R_{\text{int}} = 0.027$

$\theta_{\max} = 27.0^\circ, \theta_{\min} = 2.3^\circ$

$h = -13 \rightarrow 13$

$k = -20 \rightarrow 20$

$l = -11 \rightarrow 12$

Refinement

Refinement on F^2

Least-squares matrix: full

Secondary atom site location: difference Fourier map

Hydrogen site location: inferred from neighbouring sites

$$R[F^2 > 2\sigma(F^2)] = 0.024$$

$$wR(F^2) = 0.058$$

$$S = 1.02$$

3386 reflections

208 parameters

1 restraint

Primary atom site location: structure-invariant direct methods

H-atom parameters constrained

$$w = 1/[\sigma^2(F_o^2) + (0.0303P)^2]$$

$$\text{where } P = (F_o^2 + 2F_c^2)/3$$

$$(\Delta/\sigma)_{\max} = 0.001$$

$$\Delta\rho_{\max} = 0.22 \text{ e } \text{\AA}^{-3}$$

$$\Delta\rho_{\min} = -0.33 \text{ e } \text{\AA}^{-3}$$

Absolute structure: Flack (1983), 1528 Friedel pairs

Flack parameter: 0.020 (13)

Special details

Geometry. All e.s.d.'s (except the e.s.d. in the dihedral angle between two l.s. planes) are estimated using the full covariance matrix. The cell e.s.d.'s are taken into account individually in the estimation of e.s.d.'s in distances, angles and torsion angles; correlations between e.s.d.'s in cell parameters are only used when they are defined by crystal symmetry. An approximate (isotropic) treatment of cell e.s.d.'s is used for estimating e.s.d.'s involving l.s. planes.

Refinement. Refinement of F^2 against ALL reflections. The weighted R -factor wR and goodness of fit S are based on F^2 , conventional R -factors R are based on F , with F set to zero for negative F^2 . The threshold expression of $F^2 > \sigma(F^2)$ is used only for calculating R -factors(gt) *etc.* and is not relevant to the choice of reflections for refinement. R -factors based on F^2 are statistically about twice as large as those based on F , and R -factors based on ALL data will be even larger.

Fractional atomic coordinates and isotropic or equivalent isotropic displacement parameters (\AA^2)

	<i>x</i>	<i>y</i>	<i>z</i>	$U_{\text{iso}}^*/U_{\text{eq}}$
Co1	0.11671 (3)	0.758726 (19)	0.15941 (5)	0.02649 (9)
S1	0.02451 (5)	0.93821 (4)	0.02671 (7)	0.02606 (14)
S2	0.01351 (6)	0.72499 (4)	0.46938 (7)	0.02860 (14)
S3	0.02648 (6)	0.60549 (4)	-0.04504 (7)	0.02819 (14)
O1	-0.00760 (15)	0.85153 (11)	0.0747 (2)	0.0323 (4)
O4	-0.01485 (17)	0.75606 (12)	0.3275 (2)	0.0354 (5)
C1	0.1093 (2)	0.98601 (17)	0.1650 (4)	0.0370 (6)
H1A	0.1248	1.0453	0.1416	0.044*
H1B	0.0575	0.9852	0.2479	0.044*
O7	-0.00597 (15)	0.66745 (11)	0.0653 (2)	0.0341 (4)
N3	0.2342 (2)	0.76329 (15)	-0.0215 (3)	0.0352 (5)
H3A	0.2121	0.8104	-0.0691	0.042*
H3B	0.3127	0.7721	0.0091	0.042*
O9	-0.08429 (17)	0.57634 (13)	-0.1176 (2)	0.0405 (5)
O6	-0.10014 (17)	0.70601 (14)	0.5464 (2)	0.0435 (5)
O5	0.09764 (18)	0.78116 (13)	0.5428 (2)	0.0410 (5)
N2	0.2285 (2)	0.65745 (15)	0.2430 (3)	0.0353 (5)
H2A	0.2123	0.6108	0.1915	0.042*
H2B	0.3089	0.6714	0.2269	0.042*
N1	0.2190 (2)	0.85854 (15)	0.2622 (3)	0.0406 (6)
H1C	0.1828	0.8664	0.3458	0.049*
H1D	0.2964	0.8383	0.2785	0.049*
O2	0.10597 (16)	0.93593 (13)	-0.0947 (2)	0.0335 (4)

supplementary materials

O3	-0.08694 (16)	0.98967 (12)	0.0079 (2)	0.0372 (5)
O8	0.10532 (17)	0.53775 (12)	0.0077 (2)	0.0400 (5)
C2	0.2341 (2)	0.94364 (17)	0.1992 (3)	0.0358 (7)
H2C	0.2828	0.9383	0.1144	0.043*
H2D	0.2803	0.9799	0.2628	0.043*
C4	0.2203 (3)	0.6309 (2)	0.3918 (3)	0.0405 (7)
H4A	0.2601	0.5756	0.4027	0.049*
H4B	0.2658	0.6716	0.4486	0.049*
C5	0.2410 (2)	0.69305 (19)	-0.1240 (3)	0.0388 (7)
H5A	0.2858	0.6455	-0.0830	0.047*
H5B	0.2882	0.7122	-0.2044	0.047*
C6	0.1144 (3)	0.6630 (2)	-0.1711 (3)	0.0400 (7)
H6A	0.1253	0.6269	-0.2522	0.048*
H6B	0.0660	0.7123	-0.1998	0.048*
C3	0.0887 (3)	0.62542 (17)	0.4424 (3)	0.0391 (7)
H3C	0.0881	0.5940	0.5294	0.047*
H3D	0.0403	0.5929	0.3755	0.047*
K1	-0.25179 (5)	0.58729 (4)	-0.34300 (7)	0.03377 (13)

Atomic displacement parameters (\AA^2)

	U^{11}	U^{22}	U^{33}	U^{12}	U^{13}	U^{23}
Co1	0.02482 (16)	0.02763 (16)	0.02701 (17)	-0.00158 (13)	-0.00214 (15)	0.00258 (18)
S1	0.0264 (3)	0.0271 (3)	0.0247 (3)	0.0020 (2)	0.0012 (3)	0.0002 (3)
S2	0.0305 (3)	0.0299 (3)	0.0255 (3)	-0.0029 (3)	-0.0005 (3)	0.0035 (3)
S3	0.0263 (3)	0.0273 (3)	0.0310 (3)	-0.0002 (2)	-0.0040 (3)	-0.0012 (3)
O1	0.0275 (9)	0.0322 (9)	0.0373 (11)	-0.0025 (8)	-0.0026 (8)	0.0060 (9)
O4	0.0347 (10)	0.0446 (12)	0.0269 (10)	0.0049 (9)	-0.0010 (8)	0.0091 (9)
C1	0.0433 (14)	0.0354 (13)	0.0324 (15)	0.0022 (12)	-0.0045 (13)	-0.0100 (16)
O7	0.0277 (9)	0.0327 (10)	0.0419 (12)	-0.0020 (8)	0.0015 (8)	-0.0095 (9)
N3	0.0294 (11)	0.0359 (12)	0.0403 (14)	-0.0045 (9)	0.0053 (10)	0.0031 (11)
O9	0.0356 (11)	0.0411 (11)	0.0448 (13)	-0.0029 (9)	-0.0125 (9)	-0.0060 (10)
O6	0.0357 (10)	0.0525 (13)	0.0424 (12)	-0.0065 (9)	0.0102 (9)	0.0090 (11)
O5	0.0464 (11)	0.0388 (11)	0.0378 (12)	-0.0084 (9)	-0.0056 (9)	-0.0048 (10)
N2	0.0315 (12)	0.0382 (13)	0.0362 (15)	0.0055 (10)	-0.0006 (10)	0.0017 (11)
N1	0.0420 (13)	0.0383 (13)	0.0416 (14)	-0.0097 (11)	-0.0125 (11)	0.0057 (12)
O2	0.0372 (10)	0.0369 (11)	0.0262 (10)	0.0008 (8)	0.0059 (8)	0.0018 (8)
O3	0.0343 (10)	0.0384 (11)	0.0388 (12)	0.0092 (8)	0.0005 (8)	0.0011 (9)
O8	0.0388 (11)	0.0319 (10)	0.0492 (14)	0.0076 (8)	-0.0064 (9)	0.0028 (10)
C2	0.0349 (15)	0.0383 (15)	0.0343 (16)	-0.0077 (12)	-0.0045 (11)	-0.0006 (12)
C4	0.0419 (16)	0.0382 (16)	0.0414 (18)	0.0080 (13)	-0.0072 (13)	0.0039 (13)
C5	0.0349 (15)	0.0422 (17)	0.0395 (17)	0.0007 (13)	0.0072 (12)	0.0031 (14)
C6	0.0484 (17)	0.0439 (17)	0.0277 (16)	-0.0019 (14)	-0.0015 (12)	-0.0001 (13)
C3	0.0534 (18)	0.0281 (14)	0.0357 (17)	0.0001 (12)	-0.0019 (14)	0.0061 (13)
K1	0.0351 (3)	0.0354 (3)	0.0309 (3)	-0.0048 (2)	-0.0014 (3)	0.0009 (3)

Geometric parameters (\AA , $^\circ$)

Co1—O1	2.1316 (18)	N2—C4	1.492 (4)
--------	-------------	-------	-----------

Co1—O7	2.1411 (18)	N2—H2A	0.9000
Co1—O4	2.142 (2)	N2—H2B	0.9000
Co1—N2	2.143 (2)	N1—C2	1.473 (4)
Co1—N3	2.146 (2)	N1—H1C	0.9000
Co1—N1	2.149 (2)	N1—H1D	0.9000
S1—O3	1.4500 (18)	O2—K1 ⁱⁱⁱ	2.8522 (19)
S1—O2	1.4567 (19)	O3—K1 ^{iv}	2.7140 (19)
S1—O1	1.4748 (18)	O8—K1 ^v	2.8893 (19)
S1—C1	1.775 (3)	C2—H2C	0.9700
S2—O5	1.443 (2)	C2—H2D	0.9700
S2—O6	1.4533 (19)	C4—C3	1.491 (4)
S2—O4	1.479 (2)	C4—H4A	0.9700
S2—C3	1.774 (3)	C4—H4B	0.9700
S3—O8	1.4472 (19)	C5—C6	1.503 (4)
S3—O9	1.4482 (19)	C5—H5A	0.9700
S3—O7	1.479 (2)	C5—H5B	0.9700
S3—C6	1.778 (3)	C6—H6A	0.9700
C1—C2	1.525 (4)	C6—H6B	0.9700
C1—H1A	0.9700	C3—H3C	0.9700
C1—H1B	0.9700	C3—H3D	0.9700
N3—C5	1.479 (4)	K1—O6 ^{vi}	2.687 (2)
N3—H3A	0.9000	K1—O3 ^{vii}	2.7140 (19)
N3—H3B	0.9000	K1—O5 ^{viii}	2.8361 (19)
O9—K1	2.816 (2)	K1—O2 ^{ix}	2.8522 (19)
O6—K1 ⁱ	2.687 (2)	K1—O8 ^x	2.8893 (19)
O5—K1 ⁱⁱ	2.8361 (19)	K1—O9	2.816 (2)
O1—Co1—O7	84.97 (7)	H2A—N2—H2B	106.6
O1—Co1—O4	83.80 (7)	C2—N1—Co1	121.71 (18)
O7—Co1—O4	84.46 (7)	C2—N1—H1C	106.9
O1—Co1—N2	174.95 (8)	Co1—N1—H1C	106.9
O7—Co1—N2	90.31 (8)	C2—N1—H1D	106.9
O4—Co1—N2	93.97 (8)	Co1—N1—H1D	106.9
O1—Co1—N3	91.86 (9)	H1C—N1—H1D	106.7
O7—Co1—N3	92.19 (8)	S1—O2—K1 ⁱⁱⁱ	172.74 (12)
O4—Co1—N3	174.73 (8)	S1—O3—K1 ^{iv}	140.30 (12)
N2—Co1—N3	90.11 (9)	S3—O8—K1 ^v	170.75 (13)
O1—Co1—N1	89.78 (8)	N1—C2—C1	112.7 (2)
O7—Co1—N1	172.81 (8)	N1—C2—H2C	109.0
O4—Co1—N1	90.10 (9)	C1—C2—H2C	109.0
N2—Co1—N1	94.76 (10)	N1—C2—H2D	109.0
N3—Co1—N1	92.89 (9)	C1—C2—H2D	109.0
O3—S1—O2	113.88 (12)	H2C—C2—H2D	107.8
O3—S1—O1	111.10 (11)	C3—C4—N2	112.6 (2)
O2—S1—O1	111.52 (12)	C3—C4—H4A	109.1
O3—S1—C1	106.18 (12)	N2—C4—H4A	109.1
O2—S1—C1	107.75 (13)	C3—C4—H4B	109.1

supplementary materials

O1—S1—C1	105.88 (13)	N2—C4—H4B	109.1
O5—S2—O6	113.38 (13)	H4A—C4—H4B	107.8
O5—S2—O4	112.20 (13)	N3—C5—C6	112.9 (2)
O6—S2—O4	111.44 (12)	N3—C5—H5A	109.0
O5—S2—C3	108.98 (13)	C6—C5—H5A	109.0
O6—S2—C3	105.87 (13)	N3—C5—H5B	109.0
O4—S2—C3	104.34 (14)	C6—C5—H5B	109.0
O8—S3—O9	114.43 (12)	H5A—C5—H5B	107.8
O8—S3—O7	111.54 (13)	C5—C6—S3	115.5 (2)
O9—S3—O7	111.12 (12)	C5—C6—H6A	108.4
O8—S3—C6	107.61 (13)	S3—C6—H6A	108.4
O9—S3—C6	105.31 (14)	C5—C6—H6B	108.4
O7—S3—C6	106.24 (13)	S3—C6—H6B	108.4
S1—O1—Co1	127.05 (10)	H6A—C6—H6B	107.5
S2—O4—Co1	124.53 (11)	C4—C3—S2	115.1 (2)
C2—C1—S1	115.1 (2)	C4—C3—H3C	108.5
C2—C1—H1A	108.5	S2—C3—H3C	108.5
S1—C1—H1A	108.5	C4—C3—H3D	108.5
C2—C1—H1B	108.5	S2—C3—H3D	108.5
S1—C1—H1B	108.5	H3C—C3—H3D	107.5
H1A—C1—H1B	107.5	O6 ^{vi} —K1—O3 ^{vii}	124.38 (7)
S3—O7—Co1	126.71 (10)	O6 ^{vi} —K1—O9	87.87 (6)
C5—N3—Co1	122.94 (17)	O3 ^{vii} —K1—O9	140.88 (6)
C5—N3—H3A	106.6	O6 ^{vi} —K1—O5 ^{viii}	71.71 (6)
Co1—N3—H3A	106.6	O3 ^{vii} —K1—O5 ^{viii}	81.07 (6)
C5—N3—H3B	106.6	O9—K1—O5 ^{viii}	134.66 (7)
Co1—N3—H3B	106.6	O6 ^{vi} —K1—O2 ^{ix}	137.80 (7)
H3A—N3—H3B	106.6	O3 ^{vii} —K1—O2 ^{ix}	91.78 (6)
S3—O9—K1	150.60 (13)	O9—K1—O2 ^{ix}	71.81 (6)
S2—O6—K1 ⁱ	147.95 (13)	O5 ^{viii} —K1—O2 ^{ix}	96.53 (6)
S2—O5—K1 ⁱⁱ	170.53 (14)	O6 ^{vi} —K1—O8 ^x	86.91 (7)
C4—N2—Co1	122.22 (18)	O3 ^{vii} —K1—O8 ^x	72.54 (6)
C4—N2—H2A	106.8	O9—K1—O8 ^x	89.77 (6)
Co1—N2—H2A	106.8	O5 ^{viii} —K1—O8 ^x	127.47 (7)
C4—N2—H2B	106.8	O2 ^{ix} —K1—O8 ^x	128.14 (6)
Co1—N2—H2B	106.8		
O3—S1—O1—Co1	-165.99 (13)	O4—S2—O5—K1 ⁱⁱ	24.0 (8)
O2—S1—O1—Co1	65.79 (17)	C3—S2—O5—K1 ⁱⁱ	139.1 (8)
C1—S1—O1—Co1	-51.14 (17)	O1—Co1—N2—C4	-81.5 (11)
O7—Co1—O1—S1	-148.59 (16)	O7—Co1—N2—C4	-102.3 (2)
O4—Co1—O1—S1	126.45 (16)	O4—Co1—N2—C4	-17.8 (2)
N2—Co1—O1—S1	-169.5 (9)	N3—Co1—N2—C4	165.5 (2)
N3—Co1—O1—S1	-56.56 (16)	N1—Co1—N2—C4	72.6 (2)
N1—Co1—O1—S1	36.33 (17)	O1—Co1—N1—C2	-34.1 (2)
O5—S2—O4—Co1	69.55 (17)	O7—Co1—N1—C2	-77.2 (8)

O6—S2—O4—Co1	-162.09 (13)	O4—Co1—N1—C2	-117.9 (2)
C3—S2—O4—Co1	-48.29 (17)	N2—Co1—N1—C2	148.1 (2)
O1—Co1—O4—S2	-159.21 (15)	N3—Co1—N1—C2	57.7 (2)
O7—Co1—O4—S2	115.26 (15)	O3—S1—O2—K1 ⁱⁱⁱ	-91.5 (9)
N2—Co1—O4—S2	25.34 (16)	O1—S1—O2—K1 ⁱⁱⁱ	35.3 (9)
N3—Co1—O4—S2	166.0 (8)	C1—S1—O2—K1 ⁱⁱⁱ	151.0 (9)
N1—Co1—O4—S2	-69.44 (15)	O2—S1—O3—K1 ^{iv}	-149.88 (15)
O3—S1—C1—C2	-177.7 (2)	O1—S1—O3—K1 ^{iv}	83.18 (19)
O2—S1—C1—C2	-55.3 (3)	C1—S1—O3—K1 ^{iv}	-31.5 (2)
O1—S1—C1—C2	64.1 (2)	O9—S3—O8—K1 ^v	-114.2 (7)
O8—S3—O7—Co1	71.69 (17)	O7—S3—O8—K1 ^v	13.0 (8)
O9—S3—O7—Co1	-159.33 (13)	C6—S3—O8—K1 ^v	129.1 (7)
C6—S3—O7—Co1	-45.30 (17)	Co1—N1—C2—C1	56.8 (3)
O1—Co1—O7—S3	117.83 (15)	S1—C1—C2—N1	-69.4 (3)
O4—Co1—O7—S3	-157.93 (15)	Co1—N2—C4—C3	43.5 (3)
N2—Co1—O7—S3	-63.97 (16)	Co1—N3—C5—C6	49.2 (3)
N3—Co1—O7—S3	26.15 (16)	N3—C5—C6—S3	-71.0 (3)
N1—Co1—O7—S3	161.1 (7)	O8—S3—C6—C5	-52.0 (3)
O1—Co1—N3—C5	-108.5 (2)	O9—S3—C6—C5	-174.4 (2)
O7—Co1—N3—C5	-23.5 (2)	O7—S3—C6—C5	67.6 (2)
O4—Co1—N3—C5	-73.9 (9)	N2—C4—C3—S2	-73.2 (3)
N2—Co1—N3—C5	66.8 (2)	O5—S2—C3—C4	-45.3 (3)
N1—Co1—N3—C5	161.6 (2)	O6—S2—C3—C4	-167.6 (2)
O8—S3—O9—K1	-136.5 (2)	O4—S2—C3—C4	74.7 (2)
O7—S3—O9—K1	96.0 (3)	S3—O9—K1—O6 ^{vi}	4.6 (3)
C6—S3—O9—K1	-18.6 (3)	S3—O9—K1—O3 ^{vii}	152.8 (2)
O5—S2—O6—K1 ⁱ	-123.5 (2)	S3—O9—K1—O5 ^{viii}	-56.8 (3)
O4—S2—O6—K1 ⁱ	108.8 (2)	S3—O9—K1—O2 ^{ix}	-137.8 (3)
C3—S2—O6—K1 ⁱ	-4.1 (3)	S3—O9—K1—O8 ^x	91.6 (3)
O6—S2—O5—K1 ⁱⁱ	-103.3 (8)		

Symmetry codes: (i) $x, y, z+1$; (ii) $x+1/2, -y+3/2, z+1$; (iii) $x+1/2, -y+3/2, z$; (iv) $-x-1/2, y+1/2, z+1/2$; (v) $-x, -y+1, z+1/2$; (vi) $x, y, z-1$; (vii) $-x-1/2, y-1/2, z-1/2$; (viii) $x-1/2, -y+3/2, z-1$; (ix) $x-1/2, -y+3/2, z$; (x) $-x, -y+1, z-1/2$.

Hydrogen-bond geometry ($\text{\AA}, ^\circ$)

$D-H\cdots A$	$D-H$	$H\cdots A$	$D\cdots A$	$D-H\cdots A$
N3—H3B \cdots O7 ⁱⁱⁱ	0.90	2.22	3.097 (3)	163.
N2—H2B \cdots O4 ⁱⁱⁱ	0.90	2.40	3.166 (3)	143.
N2—H2B \cdots O1 ⁱⁱⁱ	0.90	2.47	3.255 (3)	146.
N1—H1D \cdots O4 ⁱⁱⁱ	0.90	2.55	3.423 (3)	165.
N3—H3A \cdots O2	0.90	2.28	3.113 (3)	153.
N2—H2A \cdots O8	0.90	2.39	3.219 (3)	152.
N1—H1C \cdots O5	0.90	2.49	3.229 (3)	140.

Symmetry codes: (iii) $x+1/2, -y+3/2, z$.

Fig. 1




Fig. 2

