

2-Methoxy-4-methyl-1-[1-(phenylsulfonyl)propan-2-yl]benzene

Bao-Jun Shi,^a Li-Chun Ding,^b Gang Chen^b and Zhen-Ting Du^{b*}

^aState Key Laboratory of Crop Stress Biology for Arid Areas, Northwest A&F University, Yangling 712100, Shannxi Province, People's Republic of China, and

^bCollege of Science, Northwest A&F University, Yangling 712100, Shannxi Province, People's Republic of China

Correspondence e-mail: duzt@nwsuaf.edu.cn


Received 27 August 2011; accepted 3 September 2011

Key indicators: single-crystal X-ray study; $T = 298\text{ K}$; mean $\sigma(\text{C}-\text{C}) = 0.006\text{ \AA}$; R factor = 0.055; wR factor = 0.187; data-to-parameter ratio = 14.3.

The title molecule, $\text{C}_{17}\text{H}_{20}\text{O}_3\text{S}$, displays a U-shaped structure; the two benzene rings are nearly parallel and partially overlapped to each other, the dihedral angle and centroid-to-centroid distance being $15.0(2)^\circ$ and $3.723(2)\text{ \AA}$. In the crystal, weak intermolecular $\text{C}-\text{H}\cdots\text{O}$ hydrogen bonds link the molecules, forming supramolecular chains running along the a axis.

Related literature

For propargylic sulfides as precursors of indene derivatives, see: Peng *et al.* (2007). For a related structure, see: Xi *et al.* (2004).



Experimental

Crystal data

$\text{C}_{17}\text{H}_{20}\text{O}_3\text{S}$	$V = 1564.1(3)\text{ \AA}^3$
$M_r = 304.39$	$Z = 4$
Monoclinic, $P2_1/c$	Mo $K\alpha$ radiation
$a = 9.101(1)\text{ \AA}$	$\mu = 0.21\text{ mm}^{-1}$
$b = 12.2579(14)\text{ \AA}$	$T = 298\text{ K}$
$c = 15.9251(17)\text{ \AA}$	$0.42 \times 0.27 \times 0.13\text{ mm}$
$\beta = 118.307(1)^\circ$	

Data collection

Bruker SMART 1000 CCD area-detector diffractometer	7654 measured reflections
Absorption correction: multi-scan (<i>SADABS</i> ; Bruker, 2001)	2754 independent reflections
$T_{\min} = 0.915$, $T_{\max} = 0.973$	1641 reflections with $I > 2\sigma(I)$
	$R_{\text{int}} = 0.050$

Refinement

$R[F^2 > 2\sigma(F^2)] = 0.055$	193 parameters
$wR(F^2) = 0.187$	H-atom parameters constrained
$S = 1.08$	$\Delta\rho_{\max} = 0.26\text{ e \AA}^{-3}$
2754 reflections	$\Delta\rho_{\min} = -0.23\text{ e \AA}^{-3}$

Table 1
Hydrogen-bond geometry (\AA , $^\circ$).

$D-\text{H}\cdots A$	$D-\text{H}$	$\text{H}\cdots A$	$D\cdots A$	$D-\text{H}\cdots A$
C11—H11B \cdots O2 ⁱ	0.96	2.59	3.503 (6)	159

Symmetry code: (i) $x + 1, y, z$.

Data collection: *SMART* (Bruker, 2007); cell refinement: *SAINT* (Bruker, 2007); data reduction: *SAINT*; program(s) used to solve structure: *SHELXTL* (Sheldrick, 2008); program(s) used to refine structure: *SHELXTL*; molecular graphics: *SHELXTL*; software used to prepare material for publication: *SHELXTL*.

Financial support from the Fundamental Research Funds for the Central Universities in NWSUAF (QN2009048) and the Excellent Young Funds (211020712) is greatly appreciated.

Supplementary data and figures for this paper are available from the IUCr electronic archives (Reference: XU5314).

References

- Bruker (2001). *SADABS*. Bruker AXS Inc., Madison, Wisconsin, USA.
- Bruker (2007). *SMART* and *SAINT*. Bruker AXS Inc., Madison, Wisconsin, USA.
- Peng, L., Zhang, X., Zhang, S. & Wang, J. (2007). *J. Org. Chem.* **72**, 1192–1197.
- Sheldrick, G. M. (2008). *Acta Cryst. A* **64**, 112–122.
- Xi, C., Lai, C., Chen, C. & Wang, R. (2004). *Synlett*, **9**, 1595–1597.

supplementary materials

Acta Cryst. (2011). E67, o2589 [doi:10.1107/S1600536811035987]

2-Methoxy-4-methyl-1-[1-(phenylsulfonyl)propan-2-yl]benzene

B.-J. Shi, L.-C. Ding, G. Chen and Z.-T. Du

Comment

Propargylic sulfides have been studied as the precursor of indene derivatives (Peng *et al.* 2007). As a result of our program of screening of new indene derivatives, we obtained a intermediate compound C₁₇H₂₀O₃S (**I**) and the synthesis and structure are reported here.

There are two benzene rings in the title compound and they exhibit face-to-face conformation. The dihedral angle between the two benzene rings is 12.0 (2) $^{\circ}$. The molecules of **I** are crystalized in P2₁/c space group which is different from that of 2-phenyl-1-(*p*-toluenesulfonyl)propan-2-ol (*Pbca*, Xi *et al.*, 2004). In the crystal structure there is an intermolecular C—H···O hydrogen-bonding interaction (Table 1), which is helpful to the stabilization of the packing.

Experimental

A mixture of 1-(1-bromopropan-2-yl)-4-methylbenzene (1.0 g, 4.7 mmol) and sodium benzenesulfinate (0.83 g, 5.6 mmol) in dry DMF (20 mL) was stirred over night at 80°C. When the reaction was completed, 50 mL water was added to the mixture and was extracted with ethyl acetate. The ethyl acetate layer was washed by 50 mL water, then 15 mL saturated sodium chloride and over anhydrous sodium sulfate and was separated on silica gel column chromatography with a gradient of petroleum ether and ethyl acetate as eluent to yield 1.3 g the title compound. The compound was then dissolved in ethyl acetate, and colorless crystals were formed on slow evaporation at room temperature over one week.

Refinement

All H atoms were placed in geometrically calculated positions and refined using a riding model with C—H = 0.93 Å and N—H = 0.86 Å and with $U_{\text{iso}}(\text{H}) = 1.2U_{\text{eq}}(\text{C}, \text{N})$.

Figures




Fig. 1. The molecular structure of the title compound with the atom numbering scheme. Displacement ellipsoids are drawn at the 30% probability level.

supplementary materials




Fig. 2. The molecular packing of (I) viewed along the α axis, with hydrogen bonds shown as dashed lines.

2-Methoxy-4-methyl-1-[1-(phenylsulfonyl)propan-2-yl]benzene

Crystal data

C ₁₇ H ₂₀ O ₃ S	$F(000) = 648$
$M_r = 304.39$	$D_x = 1.293 \text{ Mg m}^{-3}$
Monoclinic, $P2_1/c$	Mo $K\alpha$ radiation, $\lambda = 0.71073 \text{ \AA}$
Hall symbol: -P 2ybc	Cell parameters from 1375 reflections
$a = 9.101 (1) \text{ \AA}$	$\theta = 2.5\text{--}23.0^\circ$
$b = 12.2579 (14) \text{ \AA}$	$\mu = 0.21 \text{ mm}^{-1}$
$c = 15.9251 (17) \text{ \AA}$	$T = 298 \text{ K}$
$\beta = 118.307 (1)^\circ$	Block, colorless
$V = 1564.1 (3) \text{ \AA}^3$	$0.42 \times 0.27 \times 0.13 \text{ mm}$
$Z = 4$	

Data collection

Bruker SMART 1000 CCD area-detector diffractometer	2754 independent reflections
Radiation source: fine-focus sealed tube graphite	1641 reflections with $I > 2\sigma(I)$
ϕ and ω scans	$R_{\text{int}} = 0.050$
Absorption correction: multi-scan (<i>SADABS</i> ; Bruker, 2001)	$\theta_{\text{max}} = 25.0^\circ, \theta_{\text{min}} = 2.5^\circ$
$T_{\text{min}} = 0.915, T_{\text{max}} = 0.973$	$h = -10 \rightarrow 10$
7654 measured reflections	$k = -14 \rightarrow 14$
	$l = -11 \rightarrow 18$

Refinement

Refinement on F^2	Primary atom site location: structure-invariant direct methods
Least-squares matrix: full	Secondary atom site location: difference Fourier map
$R[F^2 > 2\sigma(F^2)] = 0.055$	Hydrogen site location: inferred from neighbouring sites
$wR(F^2) = 0.187$	H-atom parameters constrained
$S = 1.08$	$w = 1/[\sigma^2(F_o^2) + (0.0032P)^2 + 0.4143P]$
2754 reflections	where $P = (F_o^2 + 2F_c^2)/3$
193 parameters	$(\Delta/\sigma)_{\text{max}} < 0.001$
	$\Delta\rho_{\text{max}} = 0.26 \text{ e \AA}^{-3}$

0 restraints

 $\Delta\rho_{\min} = -0.23 \text{ e \AA}^{-3}$ *Special details*

Geometry. All esds (except the esd in the dihedral angle between two l.s. planes) are estimated using the full covariance matrix. The cell esds are taken into account individually in the estimation of esds in distances, angles and torsion angles; correlations between esds in cell parameters are only used when they are defined by crystal symmetry. An approximate (isotropic) treatment of cell esds is used for estimating esds involving l.s. planes.

Refinement. Refinement of F^2 against ALL reflections. The weighted R-factor wR and goodness of fit S are based on F^2 , conventional R-factors R are based on F, with F set to zero for negative F^2 . The threshold expression of $F^2 > 2\text{sigma}(F^2)$ is used only for calculating R-factors(gt) etc. and is not relevant to the choice of reflections for refinement. R-factors based on F^2 are statistically about twice as large as those based on F, and R-factors based on ALL data will be even larger.

Fractional atomic coordinates and isotropic or equivalent isotropic displacement parameters (\AA^2)

	x	y	z	$U_{\text{iso}}^*/U_{\text{eq}}$
O1	0.9829 (3)	0.3534 (2)	0.33678 (17)	0.0512 (7)
O2	0.3680 (3)	0.4052 (2)	0.16790 (19)	0.0622 (8)
O3	0.4268 (4)	0.2682 (2)	0.2938 (2)	0.0677 (9)
S1	0.48398 (11)	0.33304 (8)	0.23977 (7)	0.0450 (3)
C1	0.6540 (4)	0.4104 (3)	0.3246 (2)	0.0419 (9)
H1A	0.6215	0.4415	0.3693	0.050*
H1B	0.7460	0.3608	0.3601	0.050*
C2	0.7182 (4)	0.5027 (3)	0.2863 (3)	0.0431 (9)
H2	0.6230	0.5508	0.2496	0.052*
C3	0.7823 (4)	0.4663 (3)	0.2192 (2)	0.0372 (9)
C4	0.9128 (4)	0.3909 (3)	0.2449 (3)	0.0394 (9)
C5	0.9644 (4)	0.3587 (3)	0.1804 (3)	0.0434 (9)
H5	1.0517	0.3091	0.1993	0.052*
C6	0.8891 (4)	0.3985 (3)	0.0877 (3)	0.0429 (9)
C7	0.7631 (4)	0.4745 (3)	0.0628 (3)	0.0470 (10)
H7	0.7123	0.5040	0.0016	0.056*
C8	0.7120 (4)	0.5070 (3)	0.1277 (3)	0.0453 (10)
H8	0.6270	0.5583	0.1091	0.054*
C9	0.8460 (5)	0.5702 (3)	0.3700 (3)	0.0619 (12)
H9A	0.9400	0.5253	0.4095	0.093*
H9B	0.7955	0.5973	0.4069	0.093*
H9C	0.8825	0.6305	0.3460	0.093*
C10	1.1077 (5)	0.2713 (4)	0.3645 (3)	0.0709 (13)
H10A	1.0622	0.2083	0.3246	0.106*
H10B	1.1445	0.2514	0.4298	0.106*
H10C	1.2005	0.2987	0.3579	0.106*
C11	0.9425 (5)	0.3583 (4)	0.0172 (3)	0.0630 (12)
H11A	0.9019	0.2854	-0.0021	0.095*
H11B	1.0621	0.3586	0.0463	0.095*
H11C	0.8975	0.4053	-0.0376	0.095*
C12	0.5631 (4)	0.2425 (3)	0.1847 (3)	0.0381 (9)

supplementary materials

C13	0.6598 (5)	0.1548 (3)	0.2355 (3)	0.0529 (10)
H13	0.6851	0.1441	0.2987	0.063*
C14	0.7178 (6)	0.0837 (3)	0.1916 (4)	0.0674 (13)
H14	0.7839	0.0248	0.2255	0.081*
C15	0.6790 (6)	0.0988 (4)	0.0979 (4)	0.0694 (13)
H15	0.7183	0.0496	0.0687	0.083*
C16	0.5828 (5)	0.1856 (4)	0.0468 (3)	0.0628 (12)
H16	0.5575	0.1958	-0.0165	0.075*
C17	0.5239 (5)	0.2579 (3)	0.0910 (3)	0.0481 (10)
H17	0.4578	0.3168	0.0572	0.058*

Atomic displacement parameters (\AA^2)

	U^{11}	U^{22}	U^{33}	U^{12}	U^{13}	U^{23}
O1	0.0449 (15)	0.0676 (16)	0.0393 (15)	0.0191 (13)	0.0185 (13)	0.0142 (13)
O2	0.0437 (16)	0.0711 (17)	0.0598 (18)	0.0197 (13)	0.0147 (15)	0.0003 (15)
O3	0.074 (2)	0.0761 (18)	0.077 (2)	-0.0205 (16)	0.0555 (18)	-0.0102 (17)
S1	0.0406 (6)	0.0526 (6)	0.0480 (6)	-0.0027 (4)	0.0259 (5)	-0.0040 (5)
C1	0.044 (2)	0.0461 (19)	0.039 (2)	0.0018 (16)	0.0228 (18)	-0.0041 (17)
C2	0.045 (2)	0.0388 (19)	0.047 (2)	0.0029 (16)	0.0223 (19)	0.0027 (17)
C3	0.0347 (19)	0.0327 (17)	0.043 (2)	-0.0027 (15)	0.0171 (18)	0.0024 (16)
C4	0.036 (2)	0.043 (2)	0.037 (2)	-0.0035 (16)	0.0158 (18)	0.0027 (17)
C5	0.036 (2)	0.045 (2)	0.048 (2)	0.0028 (16)	0.0190 (19)	0.0011 (18)
C6	0.040 (2)	0.046 (2)	0.044 (2)	-0.0085 (16)	0.0200 (18)	-0.0016 (17)
C7	0.046 (2)	0.054 (2)	0.036 (2)	-0.0035 (18)	0.0164 (19)	0.0102 (18)
C8	0.042 (2)	0.041 (2)	0.051 (2)	0.0014 (16)	0.021 (2)	0.0116 (18)
C9	0.069 (3)	0.054 (2)	0.065 (3)	-0.009 (2)	0.034 (3)	-0.012 (2)
C10	0.058 (3)	0.095 (3)	0.063 (3)	0.039 (2)	0.032 (2)	0.035 (3)
C11	0.067 (3)	0.082 (3)	0.046 (2)	-0.005 (2)	0.031 (2)	-0.008 (2)
C12	0.035 (2)	0.0393 (18)	0.040 (2)	-0.0058 (15)	0.0178 (17)	-0.0037 (17)
C13	0.054 (2)	0.057 (2)	0.049 (2)	0.0062 (19)	0.025 (2)	0.008 (2)
C14	0.067 (3)	0.047 (2)	0.097 (4)	0.013 (2)	0.046 (3)	0.007 (3)
C15	0.079 (3)	0.058 (3)	0.085 (4)	0.000 (2)	0.050 (3)	-0.017 (3)
C16	0.070 (3)	0.072 (3)	0.052 (3)	-0.004 (2)	0.033 (2)	-0.012 (2)
C17	0.049 (2)	0.051 (2)	0.043 (2)	-0.0013 (18)	0.021 (2)	0.0019 (19)

Geometric parameters (\AA , $^\circ$)

O1—C4	1.369 (4)	C8—H8	0.9300
O1—C10	1.422 (4)	C9—H9A	0.9600
O2—S1	1.435 (3)	C9—H9B	0.9600
O3—S1	1.438 (3)	C9—H9C	0.9600
S1—C12	1.766 (4)	C10—H10A	0.9600
S1—C1	1.771 (3)	C10—H10B	0.9600
C1—C2	1.527 (5)	C10—H10C	0.9600
C1—H1A	0.9700	C11—H11A	0.9600
C1—H1B	0.9700	C11—H11B	0.9600
C2—C3	1.507 (5)	C11—H11C	0.9600
C2—C9	1.530 (5)	C12—C17	1.372 (5)

C2—H2	0.9800	C12—C13	1.382 (5)
C3—C8	1.378 (5)	C13—C14	1.370 (6)
C3—C4	1.404 (5)	C13—H13	0.9300
C4—C5	1.375 (5)	C14—C15	1.374 (6)
C5—C6	1.389 (5)	C14—H14	0.9300
C5—H5	0.9300	C15—C16	1.373 (6)
C6—C7	1.382 (5)	C15—H15	0.9300
C6—C11	1.501 (5)	C16—C17	1.388 (5)
C7—C8	1.379 (5)	C16—H16	0.9300
C7—H7	0.9300	C17—H17	0.9300
C4—O1—C10	117.9 (3)	C2—C9—H9A	109.5
O2—S1—O3	118.67 (18)	C2—C9—H9B	109.5
O2—S1—C12	108.05 (17)	H9A—C9—H9B	109.5
O3—S1—C12	107.38 (16)	C2—C9—H9C	109.5
O2—S1—C1	108.94 (16)	H9A—C9—H9C	109.5
O3—S1—C1	105.84 (17)	H9B—C9—H9C	109.5
C12—S1—C1	107.48 (17)	O1—C10—H10A	109.5
C2—C1—S1	117.0 (2)	O1—C10—H10B	109.5
C2—C1—H1A	108.0	H10A—C10—H10B	109.5
S1—C1—H1A	108.0	O1—C10—H10C	109.5
C2—C1—H1B	108.0	H10A—C10—H10C	109.5
S1—C1—H1B	108.0	H10B—C10—H10C	109.5
H1A—C1—H1B	107.3	C6—C11—H11A	109.5
C3—C2—C1	114.4 (3)	C6—C11—H11B	109.5
C3—C2—C9	112.8 (3)	H11A—C11—H11B	109.5
C1—C2—C9	109.2 (3)	C6—C11—H11C	109.5
C3—C2—H2	106.6	H11A—C11—H11C	109.5
C1—C2—H2	106.6	H11B—C11—H11C	109.5
C9—C2—H2	106.6	C17—C12—C13	120.6 (4)
C8—C3—C4	116.8 (4)	C17—C12—S1	119.8 (3)
C8—C3—C2	120.0 (3)	C13—C12—S1	119.6 (3)
C4—C3—C2	123.2 (3)	C14—C13—C12	119.3 (4)
O1—C4—C5	123.4 (3)	C14—C13—H13	120.4
O1—C4—C3	115.7 (3)	C12—C13—H13	120.4
C5—C4—C3	120.9 (3)	C13—C14—C15	120.4 (4)
C4—C5—C6	121.5 (3)	C13—C14—H14	119.8
C4—C5—H5	119.3	C15—C14—H14	119.8
C6—C5—H5	119.3	C16—C15—C14	120.7 (4)
C7—C6—C5	117.8 (4)	C16—C15—H15	119.6
C7—C6—C11	121.7 (4)	C14—C15—H15	119.6
C5—C6—C11	120.5 (3)	C15—C16—C17	119.1 (4)
C8—C7—C6	120.6 (4)	C15—C16—H16	120.5
C8—C7—H7	119.7	C17—C16—H16	120.5
C6—C7—H7	119.7	C12—C17—C16	119.9 (4)
C3—C8—C7	122.4 (3)	C12—C17—H17	120.0
C3—C8—H8	118.8	C16—C17—H17	120.0
C7—C8—H8	118.8		
O2—S1—C1—C2	-39.9 (3)	C5—C6—C7—C8	-1.6 (5)

supplementary materials


O3—S1—C1—C2	-168.6 (3)	C11—C6—C7—C8	177.5 (3)
C12—S1—C1—C2	76.9 (3)	C4—C3—C8—C7	1.4 (5)
S1—C1—C2—C3	-60.9 (4)	C2—C3—C8—C7	-178.4 (3)
S1—C1—C2—C9	171.6 (3)	C6—C7—C8—C3	-0.1 (5)
C1—C2—C3—C8	121.4 (3)	O2—S1—C12—C17	7.0 (3)
C9—C2—C3—C8	-112.9 (4)	O3—S1—C12—C17	136.1 (3)
C1—C2—C3—C4	-58.5 (4)	C1—S1—C12—C17	-110.4 (3)
C9—C2—C3—C4	67.2 (4)	O2—S1—C12—C13	-171.0 (3)
C10—O1—C4—C5	-4.4 (5)	O3—S1—C12—C13	-41.9 (3)
C10—O1—C4—C3	175.7 (3)	C1—S1—C12—C13	71.6 (3)
C8—C3—C4—O1	178.7 (3)	C17—C12—C13—C14	0.7 (6)
C2—C3—C4—O1	-1.4 (5)	S1—C12—C13—C14	178.7 (3)
C8—C3—C4—C5	-1.2 (5)	C12—C13—C14—C15	-0.7 (6)
C2—C3—C4—C5	178.7 (3)	C13—C14—C15—C16	0.5 (7)
O1—C4—C5—C6	179.6 (3)	C14—C15—C16—C17	-0.5 (7)
C3—C4—C5—C6	-0.5 (5)	C13—C12—C17—C16	-0.7 (5)
C4—C5—C6—C7	1.9 (5)	S1—C12—C17—C16	-178.6 (3)
C4—C5—C6—C11	-177.3 (3)	C15—C16—C17—C12	0.5 (6)

Hydrogen-bond geometry (\AA , $^\circ$)

$D\text{—H}\cdots A$	$D\text{—H}$	$H\cdots A$	$D\cdots A$	$D\text{—H}\cdots A$
C11—H11B \cdots O2 ⁱ	0.96	2.59	3.503 (6)	159

Symmetry codes: (i) $x+1, y, z$.

Fig. 1



supplementary materials

Fig. 2

