

Monoclinic, $P2_1/c$
 $a = 17.7495 (13)$ Å
 $b = 9.4273 (6)$ Å
 $c = 9.4684 (7)$ Å
 $\beta = 103.052 (3)^\circ$
 $V = 1543.42 (19)$ Å³

$Z = 4$
Mo $K\alpha$ radiation
 $\mu = 4.08$ mm⁻¹
 $T = 295$ K
 $0.20 \times 0.10 \times 0.10$ mm

***N'*-[(E)-2-Hydroxy-3,5-diiodobenzylidene]cyclohexane-1-carbohydrazide**

A. Thirugnanasundar,^a J. Suresh,^b C. Meenakshi,^c G. Chakkavarthi^{d*} and G. Rajagopal^{e*}

^aDepartment of Chemistry, Velalar College of Engineering and Technology, Erode 638 009, India, ^bDepartment of Physics, The Madura College, Madurai 625 011, India, ^cDepartment of Chemistry, Government Arts College for Women (Autonomous), Madurai 625 002, India, ^dDepartment of Physics, CPCL Polytechnic College, Chennai 600 068, India, and ^eDepartment of Chemistry, Government Arts College, Melur 625 106, India

Correspondence e-mail: chakkavarthi_2005@yahoo.com, rajagopal18@yahoo.com


Received 5 September 2011; accepted 7 September 2011

Key indicators: single-crystal X-ray study; $T = 295$ K; mean $\sigma(C-C) = 0.018$ Å; R factor = 0.058; wR factor = 0.172; data-to-parameter ratio = 14.9.

In the title compound, $C_{14}H_{10}I_2N_2O_2$, the two aromatic rings are inclined at a dihedral angle of 16.72 (33)°. The molecular structure is stabilized by an intramolecular O—H···N hydrogen bond. In the crystal, intermolecular N—H···O interactions link the molecules into chains running along the c axis. C—H···O interactions also occur. The crystal used for the structure determination was a non-merohedrally twinned with a domain ratio of 0.972 (2):0.028 (2).

Related literature

For the biological activity of Schiff base derivatives, see: Daier *et al.* (2004); Golcu *et al.* (2005); Liu & Yang (2010); Zgierski & Grabowska (2000). For related structures, see: Manvizhi *et al.* (2011); Thirugnanasundar *et al.* (2011).



Experimental

Crystal data

$C_{14}H_{10}I_2N_2O_2$

$M_r = 492.04$

Data collection

Bruker Kappa APEXII diffractometer
Absorption correction: multi-scan (*SADABS*; Sheldrick, 1996)
 $T_{\min} = 0.496$, $T_{\max} = 0.686$

13146 measured reflections
2708 independent reflections
2349 reflections with $I > 2\sigma(I)$
 $R_{\text{int}} = 0.029$

Refinement

$R[F^2 > 2\sigma(F^2)] = 0.058$
 $wR(F^2) = 0.172$
 $S = 1.11$
2708 reflections

182 parameters
H-atom parameters constrained
 $\Delta\rho_{\max} = 1.48$ e Å⁻³
 $\Delta\rho_{\min} = -1.52$ e Å⁻³

Table 1
Hydrogen-bond geometry (Å, °).

$D\cdots H\cdots A$	$D-H$	$H\cdots A$	$D\cdots A$	$D-H\cdots A$
O2—H2A···N2	0.85	1.84	2.586 (13)	145
N1—H1···O1 ⁱ	0.86	2.02	2.824 (14)	155
C3—H3···O2 ⁱⁱ	0.93	2.53	3.365 (19)	150

Symmetry codes: (i) $x, -y + \frac{1}{2}, z + \frac{1}{2}$; (ii) $-x + 1, y + \frac{1}{2}, -z + \frac{3}{2}$.

Data collection: *APEX2* (Bruker, 2004); cell refinement: *SAINT* (Bruker, 2004); data reduction: *SAINT*; program(s) used to solve structure: *SHELXS97* (Sheldrick, 2008); program(s) used to refine structure: *SHELXL97* (Sheldrick, 2008); molecular graphics: *PLATON* (Spek, 2009); software used to prepare material for publication: *SHELXL97*.

The authors wish to acknowledge the SAIF, IIT, Madras, for the data collection.

Supplementary data and figures for this paper are available from the IUCr electronic archives (Reference: BT5637).

References

- Bruker (2004). *APEX2* and *SAINT*. Bruker AXS Inc., Madison, Wisconsin, USA.
- Daier, V., Biava, H., Palopoli, C., Shove, S., Tuchagues, J. P. & Signorella, S. (2004). *J. Inorg. Biochem.* **98**, 1806–1817.
- Golcu, A., Turner, M., Demirelli, H. & Wheatley, R. A. (2005). *Inorg. Chim. Acta*, **358**, 1785–1797.
- Liu, Y.-C. & Yang, Z.-Y. (2010). *J. Biochem.* **147**, 381–391.
- Manvizhi, K., Chakkavarthi, G., Anbalagan, G. & Rajagopal, G. (2011). *Acta Cryst. E67*, o2500.
- Sheldrick, G. M. (1996). *SADABS*. University of Göttingen, Germany.
- Sheldrick, G. M. (2008). *Acta Cryst. A64*, 112–122.
- Spek, A. L. (2009). *Acta Cryst. D65*, 148–155.
- Thirugnanasundar, A., Suresh, J., Ramu, A. & RajaGopal, G. (2011). *Acta Cryst. E67*, o2303.
- Zgierski, M. Z. & Grabowska, A. (2000). *J. Chem. Phys.* **113**, 7845–7852.

supplementary materials

Acta Cryst. (2011). E67, o2620 [doi:10.1107/S1600536811036488]

***N'*-[(E)-2-Hydroxy-3,5-diiodobenzylidene]cyclohexane-1-carbohydrazide**

A. Thirugnanasundar, J. Suresh, C. Meenakshi, G. Chakkaravarthi and G. Rajagopal

Comment

Schiff base ligands play an important role in the development of coordination chemistry and possess important properties such as biological activity (Golcu *et al.*, 2005; Liu & Yang, 2010), catalytic activity (Daier *et al.*, 2004) and photochromic properties (Zgierski & Grabowska, 2000).

The geometric parameters of the title compound (Fig. 1) agree well with related structures (Manvizhi *et al.*, 2011; Thirugnanasundar *et al.*, 2011). The two aromatic rings are inclined at an angle of 16.72 (33)°.

The molecular structure is stabilized by an intramolecular O—H···N hydrogen bond. The crystal structure is controlled by intermolecular N—H···O and C—H···O (Fig. 2 & Table 1) interactions.

Experimental

A methanolic solution (10 ml) of benzoic acid hydrazide (5 mmol) was magnetically stirred in a round bottom flask followed by drop wise addition of methanolic solution of 3,5-diiodosalicylaldehyde (5 mmol). The reaction mixture was then refluxed for three hours and upon cooling to 278 K, a yellow crystalline solid precipitates from the mixture. The solid which is separated out was filtered washed with ice cold ethanol and dried in vacuo over anhydrous CaCl_2 . Single crystals suitable for the X-ray diffraction were obtained by slow evaporation of a solution of the title compound in DMF at room temperature. Melting Point : 485 K.

Refinement

All H atoms were positioned geometrically with C—H = 0.93 Å, O—H = 0.82 Å and N—H = 0.86 Å and allowed to ride on their parent atoms, with $U_{\text{iso}}(\text{H}) = 1.5 U_{\text{eq}}(\text{O})$, 1.2 $U_{\text{eq}}(\text{N})$ and 1.2 $U_{\text{eq}}(\text{C})$. Initial checkCIF/PLATON results indicated possible twinning; introduction of the suggested command TWIN 1 0 0.85 0 -1 0 0 0 -1 2 during refinement gave a very modest decrease in the R-factor, from 0.0743 to 0.0584, with BASF = 0.028 (2).

Figures




Fig. 1. The molecular structure of (I), with atom labels and 30% probability displacement ellipsoids for non-H atoms.

supplementary materials




Fig. 2. The packing of (I), viewed down b axis. Intermolecular Hydrogen bonds are shown as dashed lines. H atoms not involved in hydrogen bonding have been omitted.

N¹-[(E)-2-Hydroxy-3,5-diiodobenzylidene]cyclohexane-1- carbohydrazide

Crystal data

C ₁₄ H ₁₀ I ₂ N ₂ O ₂	$F(000) = 920$
$M_r = 492.04$	$D_x = 2.118 \text{ Mg m}^{-3}$
Monoclinic, $P2_1/c$	Mo $K\alpha$ radiation, $\lambda = 0.71073 \text{ \AA}$
Hall symbol: -P 2ybc	Cell parameters from 13146 reflections
$a = 17.7495 (13) \text{ \AA}$	$\theta = 2.2\text{--}25.0^\circ$
$b = 9.4273 (6) \text{ \AA}$	$\mu = 4.08 \text{ mm}^{-1}$
$c = 9.4684 (7) \text{ \AA}$	$T = 295 \text{ K}$
$\beta = 103.052 (3)^\circ$	Block, yellow
$V = 1543.42 (19) \text{ \AA}^3$	$0.20 \times 0.10 \times 0.10 \text{ mm}$
$Z = 4$	

Data collection

Bruker Kappa APEXII diffractometer	2708 independent reflections
Radiation source: fine-focus sealed tube graphite	2349 reflections with $I > 2\sigma(I)$
ω and φ scans	$R_{\text{int}} = 0.029$
Absorption correction: multi-scan (<i>SADABS</i> ; Sheldrick, 1996)	$\theta_{\text{max}} = 25.0^\circ, \theta_{\text{min}} = 2.2^\circ$
$T_{\text{min}} = 0.496, T_{\text{max}} = 0.686$	$h = -21 \rightarrow 21$
13146 measured reflections	$k = -11 \rightarrow 8$
	$l = -11 \rightarrow 11$

Refinement

Refinement on F^2	Primary atom site location: structure-invariant direct methods
Least-squares matrix: full	Secondary atom site location: difference Fourier map
$R[F^2 > 2\sigma(F^2)] = 0.058$	Hydrogen site location: inferred from neighbouring sites
$wR(F^2) = 0.172$	H-atom parameters constrained
$S = 1.11$	$w = 1/[\sigma^2(F_o^2) + (0.0412P)^2 + 42.3729P]$ where $P = (F_o^2 + 2F_c^2)/3$

2708 reflections	$(\Delta/\sigma)_{\max} < 0.001$
182 parameters	$\Delta\rho_{\max} = 1.48 \text{ e \AA}^{-3}$
0 restraints	$\Delta\rho_{\min} = -1.52 \text{ e \AA}^{-3}$

Fractional atomic coordinates and isotropic or equivalent isotropic displacement parameters (\AA^2)

	<i>x</i>	<i>y</i>	<i>z</i>	$U_{\text{iso}}^*/U_{\text{eq}}$
I1	0.89740 (5)	0.16968 (10)	0.83785 (10)	0.0458 (3)
I2	0.91863 (7)	-0.18253 (14)	1.36905 (13)	0.0777 (5)
O1	0.5247 (5)	0.3352 (11)	0.8089 (10)	0.052 (3)
O2	0.7346 (5)	0.2255 (10)	0.9018 (10)	0.044 (2)
H2A	0.6895	0.2404	0.9145	0.065*
N1	0.5428 (6)	0.2182 (13)	1.0250 (11)	0.043 (3)
H1	0.5244	0.1906	1.0969	0.052*
N2	0.6169 (6)	0.1863 (12)	1.0158 (11)	0.040 (3)
C1	0.4189 (7)	0.3296 (14)	0.9272 (12)	0.036 (3)
C2	0.3821 (7)	0.4436 (15)	0.8509 (13)	0.043 (3)
H2	0.4075	0.4988	0.7950	0.052*
C3	0.3079 (9)	0.4755 (18)	0.8577 (16)	0.057 (4)
H3	0.2836	0.5541	0.8082	0.068*
C4	0.2694 (8)	0.3939 (16)	0.9356 (15)	0.047 (3)
H4	0.2185	0.4156	0.9369	0.057*
C5	0.3052 (7)	0.2777 (15)	1.0138 (13)	0.041 (3)
H5	0.2790	0.2217	1.0678	0.049*
C6	0.3807 (7)	0.2479 (15)	1.0092 (14)	0.041 (3)
H6	0.4060	0.1719	1.0621	0.049*
C7	0.4996 (7)	0.2964 (14)	0.9137 (11)	0.035 (3)
C8	0.6579 (7)	0.1120 (14)	1.1159 (12)	0.036 (3)
H8	0.6375	0.0800	1.1924	0.043*
C9	0.7375 (7)	0.0770 (13)	1.1107 (13)	0.034 (3)
C10	0.7797 (7)	-0.0131 (15)	1.2142 (14)	0.042 (3)
H10	0.7576	-0.0486	1.2872	0.051*
C11	0.8534 (8)	-0.0503 (15)	1.2102 (14)	0.043 (3)
C12	0.8881 (7)	0.0029 (13)	1.1027 (14)	0.040 (3)
H12	0.9385	-0.0225	1.1010	0.048*
C13	0.8471 (7)	0.0932 (13)	0.9997 (13)	0.034 (3)
C14	0.7710 (6)	0.1333 (13)	1.0022 (12)	0.031 (2)

Atomic displacement parameters (\AA^2)

	U^{11}	U^{22}	U^{33}	U^{12}	U^{13}	U^{23}
I1	0.0346 (4)	0.0620 (6)	0.0442 (5)	0.0048 (4)	0.0159 (4)	0.0154 (4)
I2	0.0733 (8)	0.0933 (9)	0.0714 (8)	0.0406 (7)	0.0266 (6)	0.0484 (7)
O1	0.041 (5)	0.074 (7)	0.047 (6)	0.002 (5)	0.021 (4)	0.010 (5)
O2	0.034 (4)	0.048 (5)	0.051 (5)	0.007 (4)	0.015 (4)	0.015 (4)
N1	0.032 (5)	0.064 (7)	0.036 (6)	0.015 (5)	0.015 (4)	0.005 (5)
N2	0.024 (5)	0.058 (7)	0.042 (6)	0.002 (5)	0.017 (4)	-0.006 (5)
C1	0.026 (6)	0.053 (8)	0.025 (6)	0.003 (5)	-0.001 (5)	-0.006 (5)

supplementary materials

C2	0.041 (7)	0.054 (8)	0.034 (7)	0.006 (6)	0.006 (6)	0.000 (6)
C3	0.052 (9)	0.066 (10)	0.051 (8)	0.018 (8)	0.010 (7)	0.002 (8)
C4	0.036 (7)	0.055 (9)	0.054 (8)	0.007 (6)	0.016 (6)	-0.002 (7)
C5	0.035 (6)	0.059 (8)	0.033 (6)	0.003 (6)	0.016 (5)	0.003 (6)
C6	0.034 (6)	0.046 (8)	0.046 (7)	0.008 (6)	0.013 (6)	0.010 (6)
C7	0.031 (6)	0.057 (8)	0.017 (5)	0.001 (5)	0.007 (4)	0.001 (5)
C8	0.034 (6)	0.052 (8)	0.023 (6)	0.005 (6)	0.011 (5)	-0.005 (5)
C9	0.034 (6)	0.039 (7)	0.031 (6)	0.003 (5)	0.010 (5)	-0.002 (5)
C10	0.042 (7)	0.053 (8)	0.036 (7)	0.007 (6)	0.016 (6)	0.006 (6)
C11	0.046 (7)	0.048 (8)	0.034 (6)	0.007 (6)	0.008 (6)	0.006 (6)
C12	0.042 (7)	0.037 (7)	0.044 (7)	0.011 (6)	0.016 (6)	0.004 (5)
C13	0.033 (6)	0.037 (7)	0.032 (6)	0.003 (5)	0.008 (5)	-0.002 (5)
C14	0.031 (6)	0.033 (6)	0.028 (6)	0.006 (5)	0.006 (5)	-0.003 (5)

Geometric parameters (\AA , $^\circ$)

I1—C13	2.070 (12)	C4—C5	1.393 (19)
I2—C11	2.090 (13)	C4—H4	0.9300
O1—C7	1.231 (14)	C5—C6	1.379 (17)
O2—C14	1.342 (14)	C5—H5	0.9300
O2—H2A	0.8480	C6—H6	0.9300
N1—C7	1.370 (15)	C8—C9	1.463 (16)
N1—N2	1.370 (13)	C8—H8	0.9300
N1—H1	0.8600	C9—C10	1.382 (17)
N2—C8	1.268 (16)	C9—C14	1.402 (16)
C1—C2	1.375 (18)	C10—C11	1.363 (18)
C1—C6	1.376 (18)	C10—H10	0.9300
C1—C7	1.500 (16)	C11—C12	1.397 (18)
C2—C3	1.366 (19)	C12—C13	1.373 (17)
C2—H2	0.9300	C12—H12	0.9300
C3—C4	1.35 (2)	C13—C14	1.409 (16)
C3—H3	0.9300		
C14—O2—H2A	108.9	O1—C7—C1	122.2 (11)
C7—N1—N2	116.6 (10)	N1—C7—C1	115.2 (10)
C7—N1—H1	121.7	N2—C8—C9	119.7 (11)
N2—N1—H1	121.7	N2—C8—H8	120.1
C8—N2—N1	118.1 (10)	C9—C8—H8	120.1
C2—C1—C6	119.8 (11)	C10—C9—C14	120.1 (11)
C2—C1—C7	117.9 (12)	C10—C9—C8	119.2 (11)
C6—C1—C7	122.2 (11)	C14—C9—C8	120.8 (11)
C3—C2—C1	119.8 (14)	C11—C10—C9	120.5 (12)
C3—C2—H2	120.1	C11—C10—H10	119.7
C1—C2—H2	120.1	C9—C10—H10	119.7
C4—C3—C2	120.7 (14)	C10—C11—C12	120.8 (12)
C4—C3—H3	119.6	C10—C11—I2	120.9 (10)
C2—C3—H3	119.6	C12—C11—I2	118.3 (9)
C3—C4—C5	120.7 (13)	C13—C12—C11	119.4 (11)
C3—C4—H4	119.6	C13—C12—H12	120.3
C5—C4—H4	119.6	C11—C12—H12	120.3

C6—C5—C4	118.3 (12)	C12—C13—C14	120.7 (11)
C6—C5—H5	120.9	C12—C13—I1	119.6 (9)
C4—C5—H5	120.9	C14—C13—I1	119.7 (9)
C1—C6—C5	120.7 (12)	O2—C14—C9	123.3 (10)
C1—C6—H6	119.7	O2—C14—C13	118.1 (10)
C5—C6—H6	119.7	C9—C14—C13	118.5 (11)
O1—C7—N1	122.6 (11)		
C7—N1—N2—C8	−178.8 (12)	N2—C8—C9—C14	4.9 (19)
C6—C1—C2—C3	0(2)	C14—C9—C10—C11	−1(2)
C7—C1—C2—C3	178.5 (12)	C8—C9—C10—C11	178.5 (13)
C1—C2—C3—C4	−2(2)	C9—C10—C11—C12	1(2)
C2—C3—C4—C5	2(2)	C9—C10—C11—I2	178.5 (10)
C3—C4—C5—C6	0(2)	C10—C11—C12—C13	0(2)
C2—C1—C6—C5	1(2)	I2—C11—C12—C13	−178.1 (10)
C7—C1—C6—C5	−177.0 (12)	C11—C12—C13—C14	0.5 (19)
C4—C5—C6—C1	−1(2)	C11—C12—C13—I1	−178.9 (10)
N2—N1—C7—O1	1.0 (19)	C10—C9—C14—O2	−177.3 (12)
N2—N1—C7—C1	−179.9 (11)	C8—C9—C14—O2	2.8 (18)
C2—C1—C7—O1	−22.7 (18)	C10—C9—C14—C13	1.5 (18)
C6—C1—C7—O1	155.4 (13)	C8—C9—C14—C13	−178.4 (11)
C2—C1—C7—N1	158.2 (12)	C12—C13—C14—O2	177.8 (11)
C6—C1—C7—N1	−23.7 (18)	I1—C13—C14—O2	−2.8 (15)
N1—N2—C8—C9	−179.4 (11)	C12—C13—C14—C9	−1.1 (18)
N2—C8—C9—C10	−175.0 (13)	I1—C13—C14—C9	178.3 (9)

Hydrogen-bond geometry (Å, °)

D—H···A	D—H	H···A	D···A	D—H···A
O2—H2A···N2	0.85	1.84	2.586 (13)	145
N1—H1···O1 ⁱ	0.86	2.02	2.824 (14)	155
C3—H3···O2 ⁱⁱ	0.93	2.53	3.365 (19)	150

Symmetry codes: (i) $x, -y+1/2, z+1/2$; (ii) $-x+1, y+1/2, -z+3/2$.

supplementary materials

Fig. 1




Fig. 2

