

(E)-N'-(3-Bromo-5-chloro-2-hydroxybenzylidene)nicotinohydrazide

M. Prabhu,^a K. Parthipan,^b A. Ramu,^c G. Chakkaravarthi^{d*} and G. Rajagopal^{e*}

^aShasun Pharmaceuticals Ltd, Chennai 600 048, India, ^bDepartment of Chemistry, Pondicherry University, Pondicherry 605014, India, ^cDepartment of Inorganic Chemistry, Madurai Kamaraj University, Madurai 625 021, India, ^dDepartment of Physics, CPCL Polytechnic College, Chennai 600 068, India, and ^eDepartment of Chemistry, Government Arts College, Melur 625 106, India
Correspondence e-mail: chakkaravarthi_2005@yahoo.com, rajagopal18@yahoo.com


Received 15 September 2011; accepted 19 September 2011

Key indicators: single-crystal X-ray study; T = 295 K; mean σ(C–C) = 0.005 Å; R factor = 0.047; wR factor = 0.125; data-to-parameter ratio = 21.5.

There are two independent molecules in the asymmetric unit of the title compound, C₁₃H₉BrClN₃O₂, in which the dihedral angles between the benzene and pyridine rings are 8.23 (9)° and 52.84 (12)°. Both the molecules exist in an *E* configuration with respect to the C=N double bond. The two molecules in the asymmetric unit are linked *via* weak C–H···O hydrogen bonds. In both the molecules, an intramolecular O–H···N hydrogen bond generate an *S*(6) graph-set motif. In the crystal, intermolecular N–H···O and C–H···O hydrogen bonds generate bifurcated *R*₂¹(7) ring motifs. The crystal packing is further stabilized by weak intermolecular N–H···O, N–H···N, C–H···O and π–π [centroid–centroid distance 3.615 (2) Å] interactions.

Related literature

For related structures, see: Naveenkumar *et al.* (2010); Su *et al.* (2010); Tecer *et al.* (2010). For hydrogen-bond motifs, see: Bernstein *et al.* (1995).



Experimental

Crystal data

C₁₃H₉BrClN₃O₂
M_r = 354.59
Monoclinic, *P*2₁/*c*
a = 18.2217 (5) Å
b = 7.4666 (2) Å
c = 23.6916 (5) Å
β = 122.685 (1)°

V = 2712.93 (12) Å³
Z = 8
Mo Kα radiation
μ = 3.23 mm⁻¹
T = 295 K
0.30 × 0.24 × 0.20 mm

Data collection

Bruker APEXII diffractometer
Absorption correction: multi-scan (SADABS; Sheldrick, 1996)
T_{min} = 0.444, T_{max} = 0.564

34991 measured reflections
8116 independent reflections
4715 reflections with I > 2σ(I)
R_{int} = 0.042

Refinement

R[F² > 2σ(F²)] = 0.047
wR(F²) = 0.125
S = 1.02
8116 reflections
377 parameters
4 restraints

H atoms treated by a mixture of independent and constrained refinement
Δρ_{max} = 1.27 e Å⁻³
Δρ_{min} = -0.84 e Å⁻³

Table 1

Hydrogen-bond geometry (Å, °).

D–H···A	D–H	H···A	D···A	D–H···A
O2–H2B···N3	0.81 (1)	1.89 (3)	2.576 (3)	142 (4)
O4–H4A···N6	0.82 (1)	1.83 (2)	2.569 (3)	149 (4)
C15–H15···O2	0.93	2.55	3.286 (4)	137
C16–H16···O1	0.93	2.41	3.185 (4)	141
N2–H2A···O3 ⁱ	0.85 (1)	2.06 (1)	2.906 (3)	171 (3)
C2–H2···O3 ⁱ	0.93	2.52	3.380 (4)	153
N5–H5A···N1 ⁱⁱ	0.86 (1)	2.16 (1)	3.009 (4)	170 (3)

Symmetry codes: (i) x, -y + ½, z - ½; (ii) -x + 1, -y + 2, -z.

Data collection: APEX2 (Bruker, 2004); cell refinement: SAINT (Bruker, 2004); data reduction: SAINT; program(s) used to solve structure: SHELXS97 (Sheldrick, 2008); program(s) used to refine structure: SHELXL97 (Sheldrick, 2008); molecular graphics: PLATON (Spek, 2009); software used to prepare material for publication: SHELXL97.

The authors wish to acknowledge the SAIF, IIT, Madras, for the data collection.

Supplementary data and figures for this paper are available from the IUCr electronic archives (Reference: NG5231).

References

Bernstein, J., Davis, R. E., Shimoni, L. & Chang, N.-L. (1995). *Angew. Chem. Int. Ed. Engl.* **34**, 1555–1573.
Bruker (2004). APEX2 and SAINT. Bruker AXS Inc., Madison, Wisconsin, USA.
Naveenkumar, H. S., Sadikun, A., Ibrahim, P., Yeap, C. S. & Fun, H.-K. (2010). *Acta Cryst.* **E66**, o1235–o1236.
Sheldrick, G. M. (1996). SADABS, University of Göttingen, Germany.
Sheldrick, G. M. (2008). *Acta Cryst.* **A64**, 112–122.
Spek, A. L. (2009). *Acta Cryst.* **D65**, 148–155.
Su, Y.-Q., Li, C. & Wang, P. (2010). *Acta Cryst.* **E66**, o670.
Tecer, E., Dege, N., Zülfiaroğlu, A., Şenyüz, N. & Batu, H. (2010). *Acta Cryst.* **E66**, o3369–o3370.

supplementary materials

Acta Cryst. (2011). E67, o2716 [doi:10.1107/S1600536811038268]

(*E*)-*N'*-(3-Bromo-5-chloro-2-hydroxybenzylidene)nicotinohydrazide

M. Prabhu, K. Parthipan, A. Ramu, G. Chakkaravarthi and G. Rajagopal

Comment

The geometric parameters of the title compound (I), agree well with the reported similar structures (Naveenkumar *et al.*, 2010; Su *et al.*, 2010; Tecer *et al.*, 2010). The dihedral angles between the (C1/C2/N1/C3/C4/C5) and (C8-C13) is 8.23 (9)° and (C14/C15/C16/C17/N4/C18) and (C21-C26) is 52.84 (12)°. Both the molecules exist in an *E* configuration with respect to the C=N double bond. The two molecules in the asymmetric unit are linked via weak C15-H15⋯O2 and C16-H16⋯O1 hydrogen bonds. In both the molecules, the intramolecular O2-H2B⋯N3 and O4-H4A⋯N6 hydrogen bonds generate six-membered rings, each with graph-set motif S(6) (see, Fig. 1).

In the crystal structure (Fig. 2), the intermolecular N2-H2A⋯O3 and C2-H2⋯O3 hydrogen bonds generates a seven-membered ring, with bifurcated R¹₂(7) ring motif. The crystal packing is further stabilized due to weak intermolecular N-H⋯N, N-H⋯O, C-H⋯O (Table 1) and $\pi\cdots\pi$ [Cg4⋯Cg4 (2-x,1-y,1-z) = 3.614 (2)Å; Cg4 is the centroid of the ring (C21-C26)] interactions.

Experimental

Nicotinoylhydrazide (5 mmol) was dissolved in 20 mL of dry methanol with stirring and warming over a period of 10 min. To the warm hydrazide solution, 3- bromo-5-chloro salicylaldehyde (5 mmol) in 20 mL of dry methanol was added and the mixture was stirred and slowly refluxed for 2 h. The mixture was then cooled down to room temperature when pale yellow crystalline compound precipitated. The compound was collected by filtration, washed well with cold methanol and dried in vacuum. Single crystals suitable for the X-ray diffraction were obtained by slow evaporation of a solution of the title compound in methanol at room temperature. Melting Point: 503 K.

Refinement

The amino and hydroxy H-atoms were located in a difference Fourier map, and were refined with N-H distance restraint of 0.86 (1) Å and O-H distance restraint of 0.82 (1) Å. All other H atoms were positioned geometrically with C-H = 0.93 Å, and allowed to ride on their parent atoms, with Uiso(H) = 1.2Ueq(C).

Figures




Fig. 1. The molecular structure of (I), with atom labels and 30% probability displacement ellipsoids for non-H atoms.




Fig. 2. The packing of (I), viewed down *b* axis. Hydrogen bonds are shown as dashed lines. H atoms not involved in hydrogen bonding have been omitted.

(*E*)-*N'*-(3-Bromo-5-chloro-2-hydroxybenzylidene)nicotinohydrazide

Crystal data

$C_{13}H_9BrClN_3O_2$

$M_r = 354.59$

Monoclinic, $P2_1/c$

Hall symbol: -P 2ybc

$a = 18.2217$ (5) Å

$b = 7.4666$ (2) Å

$c = 23.6916$ (5) Å

$\beta = 122.685$ (1)°

$V = 2712.93$ (12) Å³

$Z = 8$

$F(000) = 1408$

$D_x = 1.736$ Mg m⁻³

Mo $K\alpha$ radiation, $\lambda = 0.71073$ Å

Cell parameters from 7402 reflections

$\theta = 2.3$ – 27.8 °

$\mu = 3.23$ mm⁻¹

$T = 295$ K

Block, pale yellow

$0.30 \times 0.24 \times 0.20$ mm

Data collection

Bruker APEXII
diffractometer

Radiation source: fine-focus sealed tube
graphite

ω and ϕ scans

Absorption correction: multi-scan
(*SADABS*; Sheldrick, 1996)

$T_{\min} = 0.444$, $T_{\max} = 0.564$

34991 measured reflections

8116 independent reflections

4715 reflections with $I > 2\sigma(I)$

$R_{\text{int}} = 0.042$

$\theta_{\max} = 30.4$ °, $\theta_{\min} = 2.3$ °

$h = -25 \rightarrow 25$

$k = -10 \rightarrow 10$

$l = -31 \rightarrow 33$

Refinement

Refinement on F^2

Least-squares matrix: full

$R[F^2 > 2\sigma(F^2)] = 0.047$

$wR(F^2) = 0.125$

$S = 1.02$

8116 reflections

Primary atom site location: structure-invariant direct methods

Secondary atom site location: difference Fourier map

Hydrogen site location: inferred from neighbouring sites

H atoms treated by a mixture of independent and constrained refinement

$w = 1/[\sigma^2(F_o^2) + (0.0543P)^2 + 1.4342P]$

where $P = (F_o^2 + 2F_c^2)/3$

$(\Delta/\sigma)_{\max} < 0.001$

377 parameters

$$\Delta\rho_{\max} = 1.27 \text{ e } \text{\AA}^{-3}$$

4 restraints

$$\Delta\rho_{\min} = -0.84 \text{ e } \text{\AA}^{-3}$$

Fractional atomic coordinates and isotropic or equivalent isotropic displacement parameters (\AA^2)

	x	y	z	$U_{\text{iso}}^*/U_{\text{eq}}$
Br1	0.750800 (19)	0.43505 (5)	0.256817 (14)	0.04545 (11)
Br2	0.92847 (3)	0.37571 (8)	0.62069 (2)	0.08778 (19)
Cl1	0.97541 (6)	0.57738 (16)	0.17505 (5)	0.0727 (3)
Cl2	1.13001 (5)	0.78996 (14)	0.56676 (6)	0.0691 (3)
O1	0.41709 (16)	0.7827 (4)	-0.01578 (11)	0.0622 (7)
O2	0.62074 (13)	0.5694 (3)	0.11875 (10)	0.0427 (5)
O3	0.56100 (13)	0.6644 (3)	0.35290 (10)	0.0471 (6)
O4	0.78596 (14)	0.5308 (4)	0.49132 (11)	0.0498 (6)
N1	0.31049 (16)	1.0700 (4)	-0.22503 (12)	0.0412 (6)
N2	0.50665 (16)	0.7928 (4)	-0.05295 (12)	0.0404 (6)
N3	0.57128 (15)	0.7165 (4)	0.00520 (11)	0.0390 (6)
N4	0.34447 (16)	0.7592 (5)	0.16222 (15)	0.0613 (9)
N5	0.65084 (14)	0.7503 (4)	0.31986 (11)	0.0371 (6)
N6	0.72089 (14)	0.7123 (4)	0.38238 (11)	0.0386 (6)
C1	0.36031 (18)	0.9139 (4)	-0.12190 (14)	0.0357 (6)
C2	0.37331 (18)	0.9923 (4)	-0.16888 (14)	0.0367 (6)
H2	0.4293	0.9904	-0.1607	0.044*
C3	0.2310 (2)	1.0706 (5)	-0.23508 (16)	0.0463 (8)
H3	0.1857	1.1219	-0.2744	0.056*
C4	0.2119 (2)	1.0005 (5)	-0.19115 (17)	0.0489 (8)
H4	0.1555	1.0062	-0.2003	0.059*
C5	0.2773 (2)	0.9218 (5)	-0.13355 (16)	0.0454 (8)
H5	0.2661	0.8742	-0.1026	0.054*
C6	0.42963 (19)	0.8249 (4)	-0.05901 (14)	0.0390 (7)
C7	0.64618 (19)	0.6968 (5)	0.01529 (14)	0.0421 (7)
H7	0.6568	0.7285	-0.0176	0.051*
C8	0.71621 (18)	0.6234 (4)	0.07931 (14)	0.0384 (7)
C9	0.70035 (18)	0.5672 (4)	0.12829 (14)	0.0348 (6)
C10	0.77050 (19)	0.5086 (4)	0.18998 (14)	0.0367 (6)
C11	0.85431 (19)	0.5085 (5)	0.20395 (15)	0.0430 (7)
H11	0.9005	0.4695	0.2454	0.052*
C12	0.86859 (19)	0.5669 (5)	0.15582 (17)	0.0482 (8)
C13	0.8007 (2)	0.6208 (5)	0.09373 (17)	0.0478 (8)
H13	0.8115	0.6557	0.0612	0.057*
C14	0.49790 (16)	0.7151 (4)	0.23663 (13)	0.0323 (6)
C15	0.50623 (19)	0.6727 (4)	0.18378 (14)	0.0389 (7)
H15	0.5603	0.6442	0.1909	0.047*
C16	0.4322 (2)	0.6739 (5)	0.12011 (15)	0.0513 (9)
H16	0.4351	0.6456	0.0831	0.062*
C17	0.3545 (2)	0.7173 (6)	0.11241 (17)	0.0604 (10)
H17	0.3050	0.7176	0.0691	0.073*
C18	0.41587 (18)	0.7556 (5)	0.22332 (16)	0.0476 (8)

supplementary materials

H18	0.4106	0.7817	0.2593	0.057*
C19	0.57123 (17)	0.7085 (4)	0.30792 (13)	0.0340 (6)
C20	0.79620 (18)	0.7645 (4)	0.39838 (15)	0.0393 (7)
H20	0.8030	0.8357	0.3693	0.047*
C21	0.87182 (18)	0.7109 (4)	0.46324 (15)	0.0390 (7)
C22	0.86279 (18)	0.5946 (4)	0.50526 (15)	0.0404 (7)
C23	0.9379 (2)	0.5408 (5)	0.56524 (15)	0.0491 (8)
C24	1.0189 (2)	0.6014 (5)	0.58363 (16)	0.0510 (9)
H24	1.0682	0.5644	0.6238	0.061*
C25	1.02624 (18)	0.7172 (5)	0.54212 (17)	0.0473 (8)
C26	0.95434 (18)	0.7721 (5)	0.48211 (16)	0.0455 (8)
H26	0.9608	0.8496	0.4543	0.055*
H2A	0.517 (2)	0.813 (5)	-0.0833 (13)	0.054 (10)*
H5A	0.655 (2)	0.804 (4)	0.2895 (12)	0.041 (9)*
H2B	0.5850 (19)	0.618 (5)	0.0836 (11)	0.063 (12)*
H4A	0.7488 (19)	0.569 (5)	0.4542 (10)	0.068 (13)*

Atomic displacement parameters (\AA^2)

	U^{11}	U^{22}	U^{33}	U^{12}	U^{13}	U^{23}
Br1	0.03930 (17)	0.0614 (2)	0.03022 (15)	-0.00234 (15)	0.01524 (12)	0.00653 (14)
Br2	0.0636 (3)	0.1316 (5)	0.0522 (2)	0.0067 (3)	0.0208 (2)	0.0320 (3)
Cl1	0.0330 (4)	0.1031 (8)	0.0796 (7)	0.0043 (5)	0.0287 (4)	0.0239 (6)
Cl2	0.0245 (4)	0.0654 (6)	0.0928 (7)	-0.0062 (4)	0.0157 (4)	-0.0170 (5)
O1	0.0541 (14)	0.102 (2)	0.0400 (13)	0.0159 (14)	0.0314 (11)	0.0236 (13)
O2	0.0291 (10)	0.0656 (15)	0.0317 (11)	0.0023 (10)	0.0152 (9)	0.0069 (11)
O3	0.0347 (11)	0.0793 (17)	0.0298 (10)	-0.0008 (11)	0.0190 (9)	0.0019 (11)
O4	0.0314 (11)	0.0760 (18)	0.0374 (12)	0.0003 (11)	0.0155 (10)	0.0022 (12)
N1	0.0359 (13)	0.0505 (16)	0.0348 (13)	0.0051 (12)	0.0176 (11)	0.0087 (12)
N2	0.0339 (13)	0.0601 (17)	0.0249 (12)	-0.0015 (12)	0.0143 (10)	0.0063 (11)
N3	0.0341 (13)	0.0509 (16)	0.0251 (11)	0.0002 (11)	0.0115 (10)	0.0051 (11)
N4	0.0264 (13)	0.094 (3)	0.0499 (17)	0.0097 (15)	0.0117 (12)	0.0140 (17)
N5	0.0219 (11)	0.0573 (17)	0.0272 (12)	0.0003 (11)	0.0100 (9)	0.0064 (11)
N6	0.0232 (11)	0.0540 (16)	0.0289 (12)	0.0022 (11)	0.0077 (9)	-0.0003 (11)
C1	0.0325 (14)	0.0427 (18)	0.0319 (14)	-0.0003 (13)	0.0174 (12)	-0.0018 (12)
C2	0.0311 (14)	0.0450 (17)	0.0352 (15)	0.0014 (13)	0.0187 (12)	0.0030 (13)
C3	0.0376 (16)	0.055 (2)	0.0391 (16)	0.0046 (15)	0.0160 (13)	0.0046 (15)
C4	0.0322 (15)	0.065 (2)	0.0488 (18)	0.0031 (15)	0.0212 (14)	0.0020 (17)
C5	0.0376 (16)	0.059 (2)	0.0467 (18)	-0.0010 (15)	0.0272 (14)	0.0026 (16)
C6	0.0361 (16)	0.050 (2)	0.0295 (14)	-0.0010 (13)	0.0171 (12)	0.0006 (13)
C7	0.0371 (16)	0.058 (2)	0.0272 (14)	-0.0037 (14)	0.0148 (12)	0.0067 (13)
C8	0.0325 (15)	0.0487 (19)	0.0306 (14)	-0.0015 (13)	0.0149 (12)	0.0043 (13)
C9	0.0294 (13)	0.0389 (16)	0.0317 (13)	-0.0021 (12)	0.0135 (11)	-0.0004 (12)
C10	0.0351 (15)	0.0404 (17)	0.0312 (14)	-0.0037 (13)	0.0156 (12)	0.0011 (12)
C11	0.0305 (15)	0.0516 (19)	0.0368 (16)	-0.0020 (14)	0.0117 (13)	0.0049 (14)
C12	0.0285 (14)	0.062 (2)	0.0513 (19)	-0.0014 (15)	0.0196 (14)	0.0073 (17)
C13	0.0375 (16)	0.065 (2)	0.0461 (18)	-0.0016 (15)	0.0260 (15)	0.0093 (16)
C14	0.0234 (12)	0.0395 (17)	0.0308 (14)	0.0019 (12)	0.0125 (11)	0.0050 (12)

C15	0.0308 (14)	0.054 (2)	0.0307 (14)	-0.0013 (13)	0.0161 (12)	0.0047 (13)
C16	0.0460 (19)	0.070 (2)	0.0288 (15)	-0.0042 (17)	0.0143 (14)	0.0041 (15)
C17	0.0329 (17)	0.087 (3)	0.0378 (18)	-0.0017 (18)	0.0037 (14)	0.0145 (18)
C18	0.0293 (15)	0.067 (2)	0.0448 (18)	0.0084 (15)	0.0190 (14)	0.0077 (16)
C19	0.0279 (13)	0.0443 (17)	0.0304 (14)	0.0024 (12)	0.0162 (11)	-0.0003 (12)
C20	0.0288 (14)	0.0459 (19)	0.0368 (15)	-0.0010 (13)	0.0135 (12)	-0.0013 (13)
C21	0.0248 (13)	0.0464 (18)	0.0373 (15)	0.0014 (13)	0.0112 (12)	-0.0089 (13)
C22	0.0283 (14)	0.054 (2)	0.0342 (15)	0.0028 (13)	0.0137 (12)	-0.0076 (13)
C23	0.0391 (17)	0.065 (2)	0.0342 (15)	0.0096 (16)	0.0142 (13)	-0.0013 (15)
C24	0.0306 (15)	0.065 (2)	0.0368 (16)	0.0095 (15)	0.0047 (13)	-0.0106 (16)
C25	0.0209 (13)	0.051 (2)	0.0547 (19)	-0.0005 (13)	0.0102 (13)	-0.0177 (17)
C26	0.0285 (14)	0.048 (2)	0.0509 (19)	-0.0029 (14)	0.0152 (14)	-0.0088 (15)

Geometric parameters (Å, °)

Br1—C10	1.885 (3)	C5—H5	0.9300
Br2—C23	1.876 (4)	C7—C8	1.462 (4)
Cl1—C12	1.745 (3)	C7—H7	0.9300
Cl2—C25	1.738 (3)	C8—C13	1.384 (4)
O1—C6	1.205 (4)	C8—C9	1.403 (4)
O2—C9	1.342 (3)	C9—C10	1.394 (4)
O2—H2B	0.814 (10)	C10—C11	1.377 (4)
O3—C19	1.222 (3)	C11—C12	1.370 (5)
O4—C22	1.341 (4)	C11—H11	0.9300
O4—H4A	0.818 (10)	C12—C13	1.375 (4)
N1—C2	1.331 (4)	C13—H13	0.9300
N1—C3	1.336 (4)	C14—C15	1.377 (4)
N2—C6	1.353 (4)	C14—C18	1.384 (4)
N2—N3	1.363 (3)	C14—C19	1.484 (4)
N2—H2A	0.853 (10)	C15—C16	1.378 (4)
N3—C7	1.261 (4)	C15—H15	0.9300
N4—C18	1.325 (4)	C16—C17	1.366 (5)
N4—C17	1.325 (5)	C16—H16	0.9300
N5—C19	1.356 (4)	C17—H17	0.9300
N5—N6	1.364 (3)	C18—H18	0.9300
N5—H5A	0.860 (10)	C20—C21	1.461 (4)
N6—C20	1.272 (4)	C20—H20	0.9300
C1—C5	1.385 (4)	C21—C26	1.394 (4)
C1—C2	1.386 (4)	C21—C22	1.396 (5)
C1—C6	1.492 (4)	C22—C23	1.398 (4)
C2—H2	0.9300	C23—C24	1.371 (5)
C3—C4	1.367 (5)	C24—C25	1.369 (5)
C3—H3	0.9300	C24—H24	0.9300
C4—C5	1.368 (5)	C25—C26	1.376 (4)
C4—H4	0.9300	C26—H26	0.9300
C9—O2—H2B	112 (3)	C11—C12—C13	121.2 (3)
C22—O4—H4A	107 (3)	C11—C12—Cl1	118.9 (2)
C2—N1—C3	116.4 (3)	C13—C12—Cl1	119.9 (3)
C6—N2—N3	117.5 (2)	C12—C13—C8	120.4 (3)

supplementary materials

C6—N2—H2A	125 (2)	C12—C13—H13	119.8
N3—N2—H2A	117 (2)	C8—C13—H13	119.8
C7—N3—N2	119.8 (2)	C15—C14—C18	118.6 (3)
C18—N4—C17	116.2 (3)	C15—C14—C19	123.6 (2)
C19—N5—N6	116.5 (2)	C18—C14—C19	117.7 (3)
C19—N5—H5A	120 (2)	C14—C15—C16	118.1 (3)
N6—N5—H5A	124 (2)	C14—C15—H15	121.0
C20—N6—N5	119.1 (3)	C16—C15—H15	121.0
C5—C1—C2	117.7 (3)	C17—C16—C15	118.6 (3)
C5—C1—C6	117.6 (3)	C17—C16—H16	120.7
C2—C1—C6	124.8 (3)	C15—C16—H16	120.7
N1—C2—C1	124.0 (3)	N4—C17—C16	124.6 (3)
N1—C2—H2	118.0	N4—C17—H17	117.7
C1—C2—H2	118.0	C16—C17—H17	117.7
N1—C3—C4	124.1 (3)	N4—C18—C14	123.8 (3)
N1—C3—H3	118.0	N4—C18—H18	118.1
C4—C3—H3	118.0	C14—C18—H18	118.1
C3—C4—C5	118.7 (3)	O3—C19—N5	122.0 (3)
C3—C4—H4	120.6	O3—C19—C14	122.3 (3)
C5—C4—H4	120.6	N5—C19—C14	115.7 (2)
C4—C5—C1	119.1 (3)	N6—C20—C21	118.8 (3)
C4—C5—H5	120.4	N6—C20—H20	120.6
C1—C5—H5	120.4	C21—C20—H20	120.6
O1—C6—N2	121.7 (3)	C26—C21—C22	119.8 (3)
O1—C6—C1	121.2 (3)	C26—C21—C20	119.3 (3)
N2—C6—C1	117.1 (2)	C22—C21—C20	120.9 (3)
N3—C7—C8	119.3 (3)	O4—C22—C21	123.7 (3)
N3—C7—H7	120.3	O4—C22—C23	117.8 (3)
C8—C7—H7	120.3	C21—C22—C23	118.4 (3)
C13—C8—C9	119.3 (3)	C24—C23—C22	121.6 (3)
C13—C8—C7	119.5 (3)	C24—C23—Br2	119.0 (2)
C9—C8—C7	121.0 (3)	C22—C23—Br2	119.4 (3)
O2—C9—C10	118.4 (3)	C25—C24—C23	119.1 (3)
O2—C9—C8	122.9 (2)	C25—C24—H24	120.5
C10—C9—C8	118.6 (3)	C23—C24—H24	120.5
C11—C10—C9	121.5 (3)	C24—C25—C26	121.5 (3)
C11—C10—Br1	119.2 (2)	C24—C25—Cl2	117.8 (2)
C9—C10—Br1	119.3 (2)	C26—C25—Cl2	120.7 (3)
C12—C11—C10	119.0 (3)	C25—C26—C21	119.6 (3)
C12—C11—H11	120.5	C25—C26—H26	120.2
C10—C11—H11	120.5	C21—C26—H26	120.2
C6—N2—N3—C7	-174.5 (3)	C7—C8—C13—C12	174.4 (3)
C19—N5—N6—C20	-173.2 (3)	C18—C14—C15—C16	0.4 (5)
C3—N1—C2—C1	0.2 (5)	C19—C14—C15—C16	176.2 (3)
C5—C1—C2—N1	-1.8 (5)	C14—C15—C16—C17	0.2 (5)
C6—C1—C2—N1	179.5 (3)	C18—N4—C17—C16	-0.8 (6)
C2—N1—C3—C4	1.3 (5)	C15—C16—C17—N4	0.0 (6)
N1—C3—C4—C5	-1.1 (6)	C17—N4—C18—C14	1.4 (6)
C3—C4—C5—C1	-0.6 (5)	C15—C14—C18—N4	-1.3 (5)

C2—C1—C5—C4	2.0 (5)	C19—C14—C18—N4	-177.4 (3)
C6—C1—C5—C4	-179.2 (3)	N6—N5—C19—O3	10.6 (4)
N3—N2—C6—O1	-2.4 (5)	N6—N5—C19—C14	-167.8 (3)
N3—N2—C6—C1	178.0 (3)	C15—C14—C19—O3	-145.3 (3)
C5—C1—C6—O1	-9.7 (5)	C18—C14—C19—O3	30.6 (5)
C2—C1—C6—O1	169.1 (3)	C15—C14—C19—N5	33.1 (4)
C5—C1—C6—N2	169.9 (3)	C18—C14—C19—N5	-151.0 (3)
C2—C1—C6—N2	-11.3 (5)	N5—N6—C20—C21	-175.0 (3)
N2—N3—C7—C8	176.8 (3)	N6—C20—C21—C26	-178.0 (3)
N3—C7—C8—C13	-172.5 (3)	N6—C20—C21—C22	4.2 (5)
N3—C7—C8—C9	3.0 (5)	C26—C21—C22—O4	179.3 (3)
C13—C8—C9—O2	178.5 (3)	C20—C21—C22—O4	-2.9 (5)
C7—C8—C9—O2	3.0 (5)	C26—C21—C22—C23	-0.9 (5)
C13—C8—C9—C10	-0.6 (5)	C20—C21—C22—C23	176.9 (3)
C7—C8—C9—C10	-176.1 (3)	O4—C22—C23—C24	-179.3 (3)
O2—C9—C10—C11	-177.8 (3)	C21—C22—C23—C24	0.9 (5)
C8—C9—C10—C11	1.3 (5)	O4—C22—C23—Br2	2.9 (4)
O2—C9—C10—Br1	0.1 (4)	C21—C22—C23—Br2	-176.9 (2)
C8—C9—C10—Br1	179.3 (2)	C22—C23—C24—C25	-0.1 (5)
C9—C10—C11—C12	-0.2 (5)	Br2—C23—C24—C25	177.8 (3)
Br1—C10—C11—C12	-178.2 (3)	C23—C24—C25—C26	-0.8 (5)
C10—C11—C12—C13	-1.6 (5)	C23—C24—C25—Cl2	-179.6 (3)
C10—C11—C12—Cl1	176.8 (3)	C24—C25—C26—C21	0.9 (5)
C11—C12—C13—C8	2.4 (6)	Cl2—C25—C26—C21	179.6 (2)
Cl1—C12—C13—C8	-176.0 (3)	C22—C21—C26—C25	0.0 (5)
C9—C8—C13—C12	-1.2 (5)	C20—C21—C26—C25	-177.8 (3)

Hydrogen-bond geometry (\AA , $^\circ$)

<i>D</i> —H \cdots <i>A</i>	<i>D</i> —H	H \cdots <i>A</i>	<i>D</i> \cdots <i>A</i>	<i>D</i> —H \cdots <i>A</i>
O2—H2B \cdots N3	0.81 (1)	1.89 (3)	2.576 (3)	142 (4)
O4—H4A \cdots N6	0.82 (1)	1.83 (2)	2.569 (3)	149 (4)
C15—H15 \cdots O2	0.93	2.55	3.286 (4)	137
C16—H16 \cdots O1	0.93	2.41	3.185 (4)	141
N2—H2A \cdots O3 ⁱ	0.85 (1)	2.06 (1)	2.906 (3)	171 (3)
C2—H2 \cdots O3 ⁱ	0.93	2.52	3.380 (4)	153
N5—H5A \cdots N1 ⁱⁱ	0.86 (1)	2.16 (1)	3.009 (4)	170 (3)

Symmetry codes: (i) $x, -y+3/2, z-1/2$; (ii) $-x+1, -y+2, -z$.

Fig. 1




Fig. 2

