

Acta Crystallographica Section E

## Structure Reports

Online

ISSN 1600-5368

## 1-(2,4,6-Triisopropylphenyl)ethanone

Amber D. Blair, Arthur D. Hendsbee and Jason D. Masuda\*

The Maritimes Centre for Green Chemistry, Department of Chemistry, Saint Mary's University, 923 Robie Street, Halifax, Nova Scotia, Canada B3H 3C3

Correspondence e-mail: jason.masuda@smu.ca


Received 12 August 2011; accepted 19 September 2011

Key indicators: single-crystal X-ray study;  $T = 129$  K; mean  $\sigma(\text{C}-\text{C}) = 0.002$  Å; disorder in main residue;  $R$  factor = 0.048;  $wR$  factor = 0.120; data-to-parameter ratio = 14.2.

The title compound,  $\text{C}_{17}\text{H}_{26}\text{O}$ , is a di-*ortho*-alkyl substituted phenyl ethanone that exhibits a significant twisting of the ketone fragment relative to the aromatic ring [ $\text{O}-\text{C}-\text{C}-\text{C}$  torsion angle =  $89.32(17)^\circ$ ] due to steric pressure from the *ortho*-isopropyl groups. One *ortho*- and the *para*-isopropyl group exhibit orientational disorder with a refined site occupancy factor of 0.562 (3):0.438 (3).

## Related literature

There are two examples in the literature of crystallographically characterized *ortho*-substituted phenyl ethanones, see: van Koningsveld *et al.* (1987); Padmanabhan *et al.* (1986); De Ridder & Schenk (1995). For the preparation, see: Delair *et al.* (1996). For the use of the title molecule in the preparation of 2-ethynyl-1,3,5-triisopropylbenzene, see: Tani *et al.* (1963). For some related ligands containing *ortho*-isopropyl groups, see: Boéré & Masuda (2002); Boéré *et al.* (2008); Giffin *et al.* (2010*a,b*, 2011).



## Experimental

## Crystal data

$\text{C}_{17}\text{H}_{26}\text{O}$   
 $M_r = 246.38$

Monoclinic,  $P2_1/n$   
 $a = 5.8590(12)$  Å

$b = 20.248(4)$  Å  
 $c = 13.148(3)$  Å  
 $\beta = 92.568(2)^\circ$   
 $V = 1558.2(6)$  Å<sup>3</sup>  
 $Z = 4$

Mo  $K\alpha$  radiation  
 $\mu = 0.06$  mm<sup>-1</sup>  
 $T = 129$  K  
 $0.45 \times 0.41 \times 0.35$  mm

## Data collection

Bruker APEXII CCD diffractometer  
 Absorption correction: multi-scan (SADABS; Bruker, 2008)  
 $T_{\min} = 0.972$ ,  $T_{\max} = 0.978$

10949 measured reflections  
 3047 independent reflections  
 2460 reflections with  $I > 2\sigma(I)$   
 $R_{\text{int}} = 0.030$

## Refinement

$R[F^2 > 2\sigma(F^2)] = 0.048$   
 $wR(F^2) = 0.120$   
 $S = 1.05$   
 3047 reflections  
 214 parameters

21 restraints  
 H-atom parameters constrained  
 $\Delta\rho_{\text{max}} = 0.19$  e Å<sup>-3</sup>  
 $\Delta\rho_{\text{min}} = -0.27$  e Å<sup>-3</sup>

Data collection: APEX2 (Bruker, 2008); cell refinement: SAINT (Bruker, 2008); data reduction: SAINT; program(s) used to solve structure: SHELXS97 (Sheldrick, 2008); program(s) used to refine structure: SHELXL97 (Sheldrick, 2008); molecular graphics: ORTEP-3 for Windows (Farrugia, 1997) and Mercury (Macrae *et al.*, 2008); software used to prepare material for publication: SHELXTL (Bruker, 2008) and publCIF (Westrip, 2010).

JDM would like to acknowledge the Canadian Foundation for Innovation Leaders Opportunity Fund (CFI-LFO) for upgrades to the diffractometer and the Natural Science and Engineering Council of Canada (NSERC) for operating funds.

Supplementary data and figures for this paper are available from the IUCr electronic archives (Reference: NR2011).

## References

- Boéré, R. T., Bond, A. M., Cronin, S., Duffy, N. W., Hazendonk, P., Masuda, J. D., Pollard, K., Roemmele, T. L., Tran, P. & Zhang, Y. (2008). *New J. Chem.* **32**, 214–231.
- Boéré, R. T. & Masuda, J. D. (2002). *Can. J. Chem.* **80**, 1607–1617.
- Bruker (2008). APEX2, SAINT and SADABS. Bruker AXS Inc., Madison, Wisconsin, USA.
- Delair, P., Kanazawa, A. M., de Azevedo, M. B. M. & Greene, A. E. (1996). *Tetrahedron Asymmetry*, **7**, 2707–2710.
- De Ridder, D. J. A. & Schenk, H. (1995). *Bull. Soc. Chim. Belg.* **104**, 81–95.
- Farrugia, L. J. (1997). *J. Appl. Cryst.* **30**, 565.
- Giffin, N. A., Hendsbee, A. D. & Masuda, J. D. (2010*a*). *Acta Cryst.* **E66**, o2194.
- Giffin, N. A., Hendsbee, A. D. & Masuda, J. D. (2010*b*). *Acta Cryst.* **E66**, o2090–o2091.
- Giffin, N. A., Hendsbee, A. D. & Masuda, J. D. (2011). *J. Organomet. Chem.* **696**, 2533–2536.
- Koningsveld, H. van, Scheele, J. J. & Jansen, J. C. (1987). *Acta Cryst.* **C43**, 294–296.
- Macrae, C. F., Bruno, I. J., Chisholm, J. A., Edgington, P. R., McCabe, P., Pidcock, E., Rodriguez-Monge, L., Taylor, R., van de Streek, J. & Wood, P. A. (2008). *J. Appl. Cryst.* **41**, 466–470.
- Padmanabhan, K., Dopp, D., Venkatesan, K. & Ramamurthy, V. (1986). *J. Chem. Soc. Perkin Trans. 2*, pp. 897–906.
- Sheldrick, G. M. (2008). *Acta Cryst.* **A64**, 112–122.
- Tani, H., Tanabe, M. & Toda, F. (1963). *Chem. Ind.* pp. 1589–1590.
- Westrip, S. P. (2010). *J. Appl. Cryst.* **43**, 920–925.

**supplementary materials**

*Acta Cryst.* (2011). E67, o2731 [ doi:10.1107/S1600536811038293 ]

## 1-(2,4,6-Triisopropylphenyl)ethanone

A. D. Blair, A. D. Hendsbee and J. D. Masuda

### Comment

We have had a long standing interest in the impact of sterically bulky *ortho*-alkyl groups on the conformation of ligands containing aryl rings, and in particular, those with *ortho*-isopropyl substituents (Boéré & Masuda, 2002; Boéré *et al.*, 2008; Giffin *et al.*, 2010a; Giffin, *et al.*, 2010b; Giffin *et al.*, 2011). The steric impact of these *ortho*-isopropyl groups has proven important in the stabilization of many reactive functional groups. In our continuing interest in these systems we have prepared and studied the title compound which is an intermediate to the related acetylene, 2-ethynyl-1,3,5-triisopropylbenzene (Tani *et al.*, 1963).

There are few solid state structures of *ortho*-alkyl substituted ethanones reported in the literature. One example is 1-*tert*-Butyl-4-acetyl-3,5-dimethyl-2,6-dinitrobenzene that exhibit O—C—C<sub>Ar</sub>—C<sub>Ar</sub> angles in the range of 77.50° to 84.12° (Padmanabhan *et al.*, 1986; De Ridder & Schenk, 1995). Another example is 4-*tert*-Butyl-2,6-dimethylacetophenone that has a O—C—C<sub>Ar</sub>—C<sub>Ar</sub> angle of 79.58° (van Koningsveld *et al.*, 1987). The title compound the ketone fragment is nearly perpendicular to the aryl ring with a torsion angle of 89.32 (17)° between the ketone C=O and the aryl ring (O1—C1—C3—C4). This can be attributed to the increased steric pressure of the *ortho*-isopropyl groups relative to the *ortho*-methyl groups in the previous two examples.

One *ortho*- and the *para*-isopropyl group exhibit two-site disorder with a refined site occupancy factor of 0.562 (3):0.438 (3). A careful look at the packing of the molecule reveals no typical hydrogen bonding and lacks significant intermolecular interactions (Figure 2).

### Experimental

The title compound was prepared following literature methods (Delair *et al.*, 1996) and was crystallized as large needles from a hot methanol solution cooled to room temperature.

### Refinement

The hydrogen atoms were placed in geometrically idealized positions of 0.98Å (methyl C—H), 0.95Å (aromatic C—H) and 1.00Å (methine C—H) and constrained to ride on the parent atom with  $U_{\text{iso}}(\text{H}) = 1.2 U_{\text{eq}}(\text{C})$  for aromatic and tertiary protons as well as  $U_{\text{iso}}(\text{H}) = 1.5 U_{\text{eq}}(\text{C})$  for the methyl groups. Two of the three isopropyl groups were modeled with two-site disorder. The isopropyl groups containing atoms C12A, C13A, C14A and C12B, C13B, C14B as well as C15A, C16A, C17A and C15B, C16B, C17B were modeled with a refined site occupancy factor of 0.562 (3):0.438 (3). In order to obtain satisfactory thermal parameters for the disordered part of the molecule DELU commands were used on the atoms C12A, C12B, C13A, C13B, C14A, C14B and C15A, C15B, C16A, C16B, C17A, C17B respectively for each isopropyl group. Unit-cell parameters were determined using a  $\theta$  range of 2.54 to 27.83° however, in order to obtain a reasonable level of data completeness,  $\theta$  was limited to 26.00° for refinement.

## Figures




Fig. 1. The molecular structure of 1-(2,4,6-triisopropylphenyl)ethanone, with atom labels and 50% probability displacement ellipsoids for non-H atoms. Hydrogen atoms have been removed for clarity.




Fig. 2. A packing diagram of 1-(2,4,6-triisopropylphenyl)ethanone viewed down the *a* axis. The minor disordered isopropyl groups are omitted for clarity.

## 1-(2,4,6-Triisopropylphenyl)ethanone

### Crystal data

$C_{17}H_{26}O$

$M_r = 246.38$

Monoclinic,  $P2_1/n$

Hall symbol:  $-P\ 2_1n$

$a = 5.8590$  (12) Å

$b = 20.248$  (4) Å

$c = 13.148$  (3) Å

$\beta = 92.568$  (2)°

$V = 1558.2$  (6) Å<sup>3</sup>

$Z = 4$

$F(000) = 544$

$D_x = 1.051$  Mg m<sup>-3</sup>

Mo  $K\alpha$  radiation,  $\lambda = 0.71073$  Å

Cell parameters from 3864 reflections

$\theta = 2.5$ – $27.8$ °

$\mu = 0.06$  mm<sup>-1</sup>

$T = 129$  K

Block, colourless

$0.45 \times 0.41 \times 0.35$  mm

### Data collection

Bruker APEXII CCD  
diffractometer

Radiation source: fine-focus sealed tube  
graphite

$\varphi$  and  $\omega$  scans

Absorption correction: multi-scan  
(*SADABS*; Bruker, 2008)

$T_{\min} = 0.972$ ,  $T_{\max} = 0.978$

3047 independent reflections

2460 reflections with  $I > 2\sigma(I)$

$R_{\text{int}} = 0.030$

$\theta_{\text{max}} = 26.0$ °,  $\theta_{\text{min}} = 2.0$ °

$h = -7 \rightarrow 7$

$k = -24 \rightarrow 24$

10949 measured reflections

$l = -16 \rightarrow 13$

*Refinement*

Refinement on  $F^2$

Primary atom site location: structure-invariant direct methods

Least-squares matrix: full

Secondary atom site location: difference Fourier map

$R[F^2 > 2\sigma(F^2)] = 0.048$

Hydrogen site location: inferred from neighbouring sites

$wR(F^2) = 0.120$

H-atom parameters constrained

$S = 1.05$

$w = 1/[\sigma^2(F_o^2) + (0.0417P)^2 + 0.6687P]$

where  $P = (F_o^2 + 2F_c^2)/3$

3047 reflections

$(\Delta/\sigma)_{\max} < 0.001$

214 parameters

$\Delta\rho_{\max} = 0.19 \text{ e } \text{\AA}^{-3}$

21 restraints

$\Delta\rho_{\min} = -0.27 \text{ e } \text{\AA}^{-3}$

*Special details*

**Geometry.** All e.s.d.'s (except the e.s.d. in the dihedral angle between two l.s. planes) are estimated using the full covariance matrix. The cell e.s.d.'s are taken into account individually in the estimation of e.s.d.'s in distances, angles and torsion angles; correlations between e.s.d.'s in cell parameters are only used when they are defined by crystal symmetry. An approximate (isotropic) treatment of cell e.s.d.'s is used for estimating e.s.d.'s involving l.s. planes.

**Refinement.** Refinement of  $F^2$  against ALL reflections. The weighted  $R$ -factor  $wR$  and goodness of fit  $S$  are based on  $F^2$ , conventional  $R$ -factors  $R$  are based on  $F$ , with  $F$  set to zero for negative  $F^2$ . The threshold expression of  $F^2 > \sigma(F^2)$  is used only for calculating  $R$ -factors(gt) *etc.* and is not relevant to the choice of reflections for refinement.  $R$ -factors based on  $F^2$  are statistically about twice as large as those based on  $F$ , and  $R$ -factors based on ALL data will be even larger.

*Fractional atomic coordinates and isotropic or equivalent isotropic displacement parameters ( $\text{\AA}^2$ )*

	<i>x</i>	<i>y</i>	<i>z</i>	$U_{\text{iso}}^*/U_{\text{eq}}$	Occ. (<1)
O1	0.01390 (17)	0.19352 (5)	0.46376 (7)	0.0360 (3)	
C1	-0.0742 (2)	0.17979 (7)	0.38182 (10)	0.0274 (3)	
C2	-0.3278 (3)	0.17911 (13)	0.36357 (14)	0.0645 (7)	
H2A	-0.4011	0.1937	0.4253	0.097*	
H2B	-0.3783	0.1342	0.3462	0.097*	
H2C	-0.3705	0.2090	0.3073	0.097*	
C3	0.0664 (2)	0.16216 (7)	0.29222 (10)	0.0250 (3)	
C4	0.1265 (2)	0.09627 (7)	0.27715 (11)	0.0298 (3)	
C5	0.2545 (2)	0.08078 (8)	0.19350 (11)	0.0337 (4)	
H5A	0.2963	0.0361	0.1827	0.040*	
C6	0.3226 (2)	0.12825 (9)	0.12577 (11)	0.0340 (4)	
C7	0.2595 (2)	0.19336 (8)	0.14225 (11)	0.0345 (4)	
H7A	0.3045	0.2265	0.0960	0.041*	
C8	0.1318 (2)	0.21148 (7)	0.22491 (10)	0.0284 (3)	
C9	0.0604 (3)	0.04223 (8)	0.35037 (14)	0.0458 (5)	
H9A	-0.0378	0.0627	0.4021	0.055*	
C10	0.2720 (4)	0.01420 (11)	0.40679 (19)	0.0743 (7)	

## supplementary materials

---

H10A	0.3480	0.0492	0.4471	0.111*	
H10B	0.3773	-0.0032	0.3574	0.111*	
H10C	0.2268	-0.0215	0.4521	0.111*	
C11	-0.0781 (4)	-0.01245 (10)	0.2970 (2)	0.0781 (8)	
H11A	-0.2156	0.0065	0.2634	0.117*	
H11B	-0.1222	-0.0453	0.3473	0.117*	
H11C	0.0146	-0.0337	0.2461	0.117*	
C12A	0.4554 (7)	0.09901 (19)	0.0358 (4)	0.0262 (8)	0.562 (3)
H12A	0.4883	0.0512	0.0487	0.031*	0.562 (3)
C13A	0.3133 (6)	0.1066 (2)	-0.0634 (3)	0.0268 (7)	0.562 (3)
H13A	0.4034	0.0925	-0.1205	0.040*	0.562 (3)
H13B	0.2693	0.1530	-0.0726	0.040*	0.562 (3)
H13C	0.1756	0.0793	-0.0609	0.040*	0.562 (3)
C14A	0.6817 (4)	0.13736 (14)	0.0294 (2)	0.0299 (7)	0.562 (3)
H14A	0.7695	0.1193	-0.0259	0.045*	0.562 (3)
H14B	0.7706	0.1331	0.0940	0.045*	0.562 (3)
H14C	0.6488	0.1841	0.0160	0.045*	0.562 (3)
C12B	0.4652 (9)	0.1237 (2)	0.0327 (5)	0.0238 (10)	0.438 (3)
H12B	0.5066	0.1689	0.0090	0.029*	0.438 (3)
C14B	0.6776 (5)	0.08476 (18)	0.0610 (3)	0.0307 (9)	0.438 (3)
H14D	0.7730	0.1098	0.1105	0.046*	0.438 (3)
H14E	0.7633	0.0766	-0.0001	0.046*	0.438 (3)
H14F	0.6347	0.0425	0.0910	0.046*	0.438 (3)
C13B	0.3143 (10)	0.0881 (3)	-0.0497 (4)	0.0287 (14)*	0.438 (3)
H13D	0.4046	0.0791	-0.1090	0.043*	0.438 (3)
H13E	0.1841	0.1163	-0.0700	0.043*	0.438 (3)
H13F	0.2584	0.0465	-0.0222	0.043*	0.438 (3)
C15A	0.0614 (13)	0.2804 (4)	0.2368 (7)	0.0403 (10)	0.562 (3)
H15A	-0.0094	0.2858	0.3041	0.048*	0.562 (3)
C16A	-0.1203 (7)	0.2960 (2)	0.1501 (3)	0.0578 (10)	0.562 (3)
H16A	-0.1752	0.3414	0.1576	0.087*	0.562 (3)
H16B	-0.2488	0.2653	0.1541	0.087*	0.562 (3)
H16C	-0.0511	0.2912	0.0840	0.087*	0.562 (3)
C17A	0.2662 (6)	0.32461 (14)	0.2330 (3)	0.0495 (9)	0.562 (3)
H17A	0.3818	0.3106	0.2847	0.074*	0.562 (3)
H17B	0.2208	0.3703	0.2462	0.074*	0.562 (3)
H17C	0.3298	0.3218	0.1654	0.074*	0.562 (3)
C15B	0.0787 (18)	0.2866 (5)	0.2488 (10)	0.044 (2)	0.438 (3)
H15B	-0.0508	0.2864	0.2958	0.053*	0.438 (3)
C16B	0.0019 (8)	0.3298 (2)	0.1582 (3)	0.0445 (10)	0.438 (3)
H16D	-0.1275	0.3088	0.1211	0.067*	0.438 (3)
H16E	0.1286	0.3350	0.1126	0.067*	0.438 (3)
H16F	-0.0445	0.3732	0.1827	0.067*	0.438 (3)
C17B	0.2869 (6)	0.32093 (18)	0.3094 (4)	0.0402 (10)	0.438 (3)
H17D	0.3228	0.2964	0.3723	0.060*	0.438 (3)
H17E	0.2459	0.3665	0.3260	0.060*	0.438 (3)
H17F	0.4206	0.3211	0.2671	0.060*	0.438 (3)

Atomic displacement parameters ( $\text{\AA}^2$ )

	$U^{11}$	$U^{22}$	$U^{33}$	$U^{12}$	$U^{13}$	$U^{23}$
O1	0.0370 (6)	0.0477 (7)	0.0232 (5)	-0.0009 (5)	-0.0008 (4)	-0.0086 (5)
C1	0.0263 (7)	0.0320 (7)	0.0240 (7)	-0.0024 (6)	0.0019 (5)	-0.0061 (6)
C2	0.0251 (8)	0.131 (2)	0.0382 (10)	-0.0011 (10)	0.0074 (7)	-0.0324 (11)
C3	0.0169 (6)	0.0364 (8)	0.0215 (7)	-0.0047 (5)	-0.0007 (5)	-0.0064 (6)
C4	0.0251 (7)	0.0345 (8)	0.0302 (8)	-0.0103 (6)	0.0061 (6)	-0.0112 (6)
C5	0.0268 (7)	0.0419 (8)	0.0330 (8)	-0.0057 (6)	0.0067 (6)	-0.0162 (7)
C6	0.0176 (6)	0.0633 (10)	0.0212 (7)	-0.0002 (6)	0.0001 (5)	-0.0076 (7)
C7	0.0210 (7)	0.0589 (10)	0.0235 (7)	0.0008 (6)	-0.0013 (5)	0.0088 (7)
C8	0.0197 (6)	0.0397 (7)	0.0253 (7)	0.0006 (5)	-0.0048 (5)	0.0021 (6)
C9	0.0527 (10)	0.0298 (8)	0.0574 (11)	-0.0150 (7)	0.0299 (9)	-0.0112 (7)
C10	0.0848 (16)	0.0629 (14)	0.0770 (16)	-0.0128 (12)	0.0222 (13)	0.0352 (12)
C11	0.0690 (14)	0.0466 (11)	0.123 (2)	-0.0340 (10)	0.0532 (14)	-0.0408 (12)
C12A	0.0276 (14)	0.027 (2)	0.0246 (13)	-0.0014 (14)	0.0059 (10)	-0.002 (2)
C13A	0.0273 (14)	0.032 (2)	0.0215 (12)	-0.0052 (15)	0.0069 (10)	-0.0046 (15)
C14A	0.0219 (11)	0.0399 (17)	0.0283 (13)	0.0005 (10)	0.0048 (9)	-0.0030 (12)
C12B	0.0224 (18)	0.027 (3)	0.0221 (16)	-0.0016 (17)	0.0031 (11)	-0.003 (3)
C14B	0.0245 (15)	0.038 (2)	0.0303 (18)	0.0031 (12)	0.0035 (12)	-0.0010 (14)
C15A	0.039 (3)	0.0478 (17)	0.035 (2)	0.0202 (17)	0.0160 (19)	0.017 (2)
C16A	0.059 (2)	0.072 (3)	0.0433 (19)	0.0391 (18)	0.0063 (15)	0.0148 (18)
C17A	0.077 (2)	0.0236 (12)	0.049 (2)	0.0030 (13)	0.0153 (17)	-0.0030 (14)
C15B	0.036 (4)	0.0340 (19)	0.061 (5)	0.006 (2)	-0.013 (3)	0.011 (2)
C16B	0.049 (2)	0.0377 (19)	0.047 (2)	0.0158 (19)	0.0026 (18)	0.0073 (17)
C17B	0.044 (2)	0.0297 (17)	0.047 (3)	-0.0006 (14)	0.0028 (16)	-0.0076 (16)

Geometric parameters ( $\text{\AA}$ ,  $^\circ$ )

O1—C1	1.2060 (17)	C13A—H13B	0.9800
C1—C2	1.495 (2)	C13A—H13C	0.9800
C1—C3	1.5102 (19)	C14A—H14A	0.9800
C2—H2A	0.9800	C14A—H14B	0.9800
C2—H2B	0.9800	C14A—H14C	0.9800
C2—H2C	0.9800	C12B—C14B	1.506 (6)
C3—C4	1.396 (2)	C12B—C13B	1.546 (8)
C3—C8	1.399 (2)	C12B—H12B	1.0000
C4—C5	1.3942 (19)	C14B—H14D	0.9800
C4—C9	1.519 (2)	C14B—H14E	0.9800
C5—C6	1.381 (2)	C14B—H14F	0.9800
C5—H5A	0.9500	C13B—H13D	0.9800
C6—C7	1.389 (2)	C13B—H13E	0.9800
C6—C12B	1.515 (7)	C13B—H13F	0.9800
C6—C12A	1.561 (5)	C15A—C17A	1.500 (8)
C7—C8	1.396 (2)	C15A—C16A	1.557 (8)
C7—H7A	0.9500	C15A—H15A	1.0000
C8—C15A	1.466 (8)	C16A—H16A	0.9800
C8—C15B	1.587 (11)	C16A—H16B	0.9800

## supplementary materials

---

C9—C11	1.525 (2)	C16A—H16C	0.9800
C9—C10	1.526 (3)	C17A—H17A	0.9800
C9—H9A	1.0000	C17A—H17B	0.9800
C10—H10A	0.9800	C17A—H17C	0.9800
C10—H10B	0.9800	C15B—C16B	1.528 (11)
C10—H10C	0.9800	C15B—C17B	1.587 (10)
C11—H11A	0.9800	C15B—H15B	1.0000
C11—H11B	0.9800	C16B—H16D	0.9800
C11—H11C	0.9800	C16B—H16E	0.9800
C12A—C13A	1.524 (7)	C16B—H16F	0.9800
C12A—C14A	1.542 (5)	C17B—H17D	0.9800
C12A—H12A	1.0000	C17B—H17E	0.9800
C13A—H13A	0.9800	C17B—H17F	0.9800
O1—C1—C2	121.88 (13)	C12A—C14A—H14A	109.5
O1—C1—C3	121.65 (12)	C12A—C14A—H14B	109.5
C2—C1—C3	116.47 (12)	H14A—C14A—H14B	109.5
C1—C2—H2A	109.5	C12A—C14A—H14C	109.5
C1—C2—H2B	109.5	H14A—C14A—H14C	109.5
H2A—C2—H2B	109.5	H14B—C14A—H14C	109.5
C1—C2—H2C	109.5	C6—C12B—C14B	108.1 (4)
H2A—C2—H2C	109.5	C6—C12B—C13B	106.0 (4)
H2B—C2—H2C	109.5	C14B—C12B—C13B	111.7 (4)
C4—C3—C8	120.95 (13)	C6—C12B—H12B	110.3
C4—C3—C1	119.11 (12)	C14B—C12B—H12B	110.3
C8—C3—C1	119.94 (13)	C13B—C12B—H12B	110.3
C5—C4—C3	118.35 (14)	C12B—C14B—H14D	109.5
C5—C4—C9	119.96 (14)	C12B—C14B—H14E	109.5
C3—C4—C9	121.68 (13)	H14D—C14B—H14E	109.5
C6—C5—C4	122.21 (14)	C12B—C14B—H14F	109.5
C6—C5—H5A	118.9	H14D—C14B—H14F	109.5
C4—C5—H5A	118.9	H14E—C14B—H14F	109.5
C5—C6—C7	118.26 (13)	C12B—C13B—H13D	109.5
C5—C6—C12B	131.6 (2)	C12B—C13B—H13E	109.5
C7—C6—C12B	110.1 (2)	H13D—C13B—H13E	109.5
C5—C6—C12A	113.13 (19)	C12B—C13B—H13F	109.5
C7—C6—C12A	128.56 (19)	H13D—C13B—H13F	109.5
C6—C7—C8	121.78 (14)	H13E—C13B—H13F	109.5
C6—C7—H7A	119.1	C8—C15A—C17A	109.6 (5)
C8—C7—H7A	119.1	C8—C15A—C16A	107.6 (5)
C7—C8—C3	118.45 (14)	C17A—C15A—C16A	112.1 (5)
C7—C8—C15A	119.8 (3)	C8—C15A—H15A	109.2
C3—C8—C15A	121.7 (3)	C17A—C15A—H15A	109.2
C7—C8—C15B	121.5 (5)	C16A—C15A—H15A	109.2
C3—C8—C15B	119.8 (5)	C15A—C16A—H16A	109.5
C4—C9—C11	112.16 (17)	C15A—C16A—H16B	109.5
C4—C9—C10	110.56 (13)	H16A—C16A—H16B	109.5
C11—C9—C10	110.95 (17)	C15A—C16A—H16C	109.5
C4—C9—H9A	107.7	H16A—C16A—H16C	109.5
C11—C9—H9A	107.7	H16B—C16A—H16C	109.5



C10—C9—H9A	107.7	C15A—C17A—H17A	109.5
C9—C10—H10A	109.5	C15A—C17A—H17B	109.5
C9—C10—H10B	109.5	H17A—C17A—H17B	109.5
H10A—C10—H10B	109.5	C15A—C17A—H17C	109.5
C9—C10—H10C	109.5	H17A—C17A—H17C	109.5
H10A—C10—H10C	109.5	H17B—C17A—H17C	109.5
H10B—C10—H10C	109.5	C16B—C15B—C17B	109.5 (7)
C9—C11—H11A	109.5	C16B—C15B—C8	116.6 (8)
C9—C11—H11B	109.5	C17B—C15B—C8	111.5 (6)
H11A—C11—H11B	109.5	C16B—C15B—H15B	106.2
C9—C11—H11C	109.5	C17B—C15B—H15B	106.2
H11A—C11—H11C	109.5	C8—C15B—H15B	106.2
H11B—C11—H11C	109.5	C15B—C16B—H16D	109.5
C13A—C12A—C14A	109.9 (4)	C15B—C16B—H16E	109.5
C13A—C12A—C6	109.8 (3)	H16D—C16B—H16E	109.5
C14A—C12A—C6	108.0 (3)	C15B—C16B—H16F	109.5
C13A—C12A—H12A	109.7	H16D—C16B—H16F	109.5
C14A—C12A—H12A	109.7	H16E—C16B—H16F	109.5
C6—C12A—H12A	109.7	C15B—C17B—H17D	109.5
C12A—C13A—H13A	109.5	C15B—C17B—H17E	109.5
C12A—C13A—H13B	109.5	H17D—C17B—H17E	109.5
H13A—C13A—H13B	109.5	C15B—C17B—H17F	109.5
C12A—C13A—H13C	109.5	H17D—C17B—H17F	109.5
H13A—C13A—H13C	109.5	H17E—C17B—H17F	109.5
H13B—C13A—H13C	109.5		
O1—C1—C3—C4	89.32 (17)	C5—C4—C9—C10	66.0 (2)
C2—C1—C3—C4	-90.50 (18)	C3—C4—C9—C10	-112.65 (17)
O1—C1—C3—C8	-91.58 (17)	C5—C6—C12A—C13A	112.2 (3)
C2—C1—C3—C8	88.60 (18)	C7—C6—C12A—C13A	-64.9 (4)
C8—C3—C4—C5	0.3 (2)	C12B—C6—C12A—C13A	-77.6 (11)
C1—C3—C4—C5	179.40 (12)	C5—C6—C12A—C14A	-128.0 (2)
C8—C3—C4—C9	178.99 (13)	C7—C6—C12A—C14A	55.0 (4)
C1—C3—C4—C9	-1.9 (2)	C12B—C6—C12A—C14A	42.2 (10)
C3—C4—C5—C6	-0.1 (2)	C5—C6—C12B—C14B	-48.4 (4)
C9—C4—C5—C6	-178.78 (14)	C7—C6—C12B—C14B	130.1 (3)
C4—C5—C6—C7	-0.3 (2)	C12A—C6—C12B—C14B	-60.5 (11)
C4—C5—C6—C12B	178.1 (3)	C5—C6—C12B—C13B	71.5 (5)
C4—C5—C6—C12A	-177.6 (2)	C7—C6—C12B—C13B	-110.1 (3)
C5—C6—C7—C8	0.4 (2)	C12A—C6—C12B—C13B	59.3 (10)
C12B—C6—C7—C8	-178.4 (3)	C7—C8—C15A—C17A	-52.9 (6)
C12A—C6—C7—C8	177.3 (3)	C3—C8—C15A—C17A	129.8 (4)
C6—C7—C8—C3	-0.1 (2)	C15B—C8—C15A—C17A	53 (6)
C6—C7—C8—C15A	-177.5 (4)	C7—C8—C15A—C16A	69.2 (5)
C6—C7—C8—C15B	174.6 (4)	C3—C8—C15A—C16A	-108.1 (5)
C4—C3—C8—C7	-0.21 (19)	C15B—C8—C15A—C16A	175 (7)
C1—C3—C8—C7	-179.30 (12)	C7—C8—C15B—C16B	44.8 (8)
C4—C3—C8—C15A	177.1 (4)	C3—C8—C15B—C16B	-140.5 (6)
C1—C3—C8—C15A	-1.9 (4)	C15A—C8—C15B—C16B	-33 (6)
C4—C3—C8—C15B	-175.0 (4)	C7—C8—C15B—C17B	-81.9 (8)

## supplementary materials

---

C1—C3—C8—C15B	5.9 (4)	C3—C8—C15B—C17B	92.8 (8)
C5—C4—C9—C11	-58.4 (2)	C15A—C8—C15B—C17B	-160 (7)
C3—C4—C9—C11	122.94 (16)		

Fig. 1




Fig. 2

