

N-(2-Chlorophenyl)-4-methylbenzamide

Vinola Z. Rodrigues,^a Peter Herich,^b B. Thimme Gowda^{a*} and Jozef Kožíšek^b

^aDepartment of Chemistry, Mangalore University, Mangalagangotri 574 199, Mangalore, India, and ^bInstitute of Physical Chemistry and Chemical Physics, Slovak University of Technology, Radlinského 9, SK-812 37 Bratislava, Slovak Republic
Correspondence e-mail: gowdabt@yahoo.com


Received 20 September 2011; accepted 21 September 2011

Key indicators: single-crystal X-ray study; $T = 293\text{ K}$; mean $\sigma(\text{C}-\text{C}) = 0.003\text{ \AA}$; R factor = 0.043; wR factor = 0.126; data-to-parameter ratio = 22.9.

The asymmetric unit of the title compound, $C_{14}H_{12}\text{ClNO}$, contains two independent molecules in which the dihedral angles between the two aromatic rings are $51.76(6)$ and $51.48(7)^\circ$. The crystal structure is stabilized by intermolecular $\text{N}-\text{H}\cdots\text{O}$ hydrogen bonds, which link the molecules into chains running along the c axis.

Related literature

For preparation of the title compound, see: Gowda *et al.* (2003). For our studies on the effects of substituents on the structures and other aspects of *N*-(aryl)-amides, see: Arjunan *et al.* (2004); Bowes *et al.* (2003); Gowda *et al.* (2001); Rodrigues *et al.* (2011); Saeed *et al.* (2010) on *N*-(aryl)-methanesulfonamides, see: Gowda *et al.* (2007) and on *N*-(aryl)-arylsulfonamides, see: Gowda *et al.* (2005).



Experimental

Crystal data

$C_{14}H_{12}\text{ClNO}$

$M_r = 245.70$

Monoclinic, $P2_1/n$

$a = 9.6940(5)\text{ \AA}$

$b = 27.4495(9)\text{ \AA}$

$c = 9.9025(4)\text{ \AA}$

$\beta = 106.730(5)^\circ$

$V = 2523.48(19)\text{ \AA}^3$

$Z = 8$

Mo $K\alpha$ radiation

$\mu = 0.29\text{ mm}^{-1}$

$T = 293\text{ K}$

$0.97 \times 0.13 \times 0.10\text{ mm}$

Data collection

Oxford Diffraction Xcalibur Ruby

Gemini diffractometer

Absorption correction: analytical [CrysAlis RED (Oxford Diffraction, 2009) based on expressions derived by Clark &

Reid (1995)]

$T_{\min} = 0.957$, $T_{\max} = 0.972$

47577 measured reflections

7045 independent reflections

2850 reflections with $I > 2\sigma(I)$

$R_{\text{int}} = 0.046$

Refinement

$R[F^2 > 2\sigma(F^2)] = 0.043$

$wR(F^2) = 0.126$

$S = 0.85$

7045 reflections

307 parameters

H-atom parameters constrained

$\Delta\rho_{\max} = 0.24\text{ e \AA}^{-3}$

$\Delta\rho_{\min} = -0.30\text{ e \AA}^{-3}$

Table 1
Hydrogen-bond geometry (\AA , $^\circ$).

$D-\text{H}\cdots A$	$D-\text{H}$	$\text{H}\cdots A$	$D\cdots A$	$D-\text{H}\cdots A$
N15—H15A \cdots O34	0.86	2.02	2.8408 (16)	159
N32—H32A \cdots O17 ⁱ	0.86	2.01	2.8455 (16)	165

Symmetry code: (i) $x, y, z - 1$.

Data collection: CrysAlis CCD (Oxford Diffraction, 2009); cell refinement: CrysAlis CCD; data reduction: CrysAlis RED (Oxford Diffraction, 2009); program(s) used to solve structure: SHELXS97 (Sheldrick, 2008); program(s) used to refine structure: SHELXL97 (Sheldrick, 2008); molecular graphics: DIAMOND (Brandenburg, 2002); software used to prepare material for publication: enCIFer (Allen *et al.*, 2004).

MF and JK thank the VEGA Grant Agency of the Slovak Ministry of Education (1/0679/11) and the Research and Development Agency of Slovakia (APVV-0202-10) for their financial support and the Structural Funds, Interreg IIIA, for financial support in purchasing the diffractometer. VZR thanks the University Grants Commission, Government of India, New Delhi, for award of an RFSMS research fellowship.

Supplementary data and figures for this paper are available from the IUCr electronic archives (Reference: BT5650).

References

- Allen, F. H., Johnson, O., Shields, G. P., Smith, B. R. & Towler, M. (2004). *J. Appl. Cryst.* **37**, 335–338.
- Arjunan, V., Mohan, S., Subramanian, S. & Gowda, B. T. (2004). *Spectrochim. Acta Part A*, **60**, 1141–1159.
- Bowes, K. F., Glidewell, C., Low, J. N., Skakle, J. M. S. & Wardell, J. L. (2003). *Acta Cryst.* **C59**, o1–o3.
- Brandenburg, K. (2002). DIAMOND. Crystal Impact GbR, Bonn, Germany.
- Clark, R. C. & Reid, J. S. (1995). *Acta Cryst.* **A51**, 887–897.
- Gowda, B. T., Foro, S. & Fuess, H. (2007). *Acta Cryst.* **E63**, o2597.
- Gowda, B. T., Jyothi, K., Paulus, H. & Fuess, H. (2003). *Z. Naturforsch. Teil A*, **58**, 225–230.
- Gowda, B. T., Paulus, H. & Fuess, H. (2001). *Z. Naturforsch. Teil A*, **56**, 386–394.
- Gowda, B. T., Shetty, M. & Jayalakshmi, K. L. (2005). *Z. Naturforsch. Teil A*, **60**, 106–112.
- Oxford Diffraction (2009). CrysAlis CCD and CrysAlis RED. Oxford Diffraction Ltd, Yarnton, England.
- Rodrigues, V. Z., Fronc, M., Gowda, B. T. & Kožíšek, J. (2011). *Acta Cryst.* **E67**, o1500.
- Saeed, A., Arshad, M. & Simpson, J. (2010). *Acta Cryst.* **E66**, o2808–o2809.
- Sheldrick, G. M. (2008). *Acta Cryst.* **A64**, 112–122.

supplementary materials

Acta Cryst. (2011). E67, o2767 [doi:10.1107/S1600536811038840]

N-(2-Chlorophenyl)-4-methylbenzamide

V. Z. Rodrigues, P. Herich, B. T. Gowda and J. Kozísek

Comment

The amide and sulfonamide moieties are the constituents of many biologically significant compounds. As part of our work on the substituent effects on the structures and other aspects of *N*-(aryl)-amides (Arjunan *et al.*, 2004; Bowes *et al.*, 2003; Gowda *et al.*, 2001; Saeed *et al.*, 2010; Rodrigues *et al.*, 2011), *N*-(aryl)-methanesulfonamides (Gowda *et al.*, 2007) and *N*-(aryl)-arylsulfonamides (Gowda *et al.*, 2005), in the present work, the crystal structure of *N*-(2-Chlorophenyl)-4-methylbenzamide (I) has been determined (Fig. 1). The asymmetric unit of (I) contains two independent molecules. In the crystal, the *ortho*-Cl substituent in the anilino ring is positioned *syn* to the N—H bond in one of the molecules and *anti* in the other molecule. Further, the N—H and C=O bonds in the C—NH—C(O)—C segment are *anti* to each other in both the molecules, similar to that observed in *N*-(2-methylphenyl)-4-methylbenzamide (II) (Rodrigues *et al.*, 2011).

The packing of molecules linked by N—H···O hydrogen bonds into infinite chains is shown in Fig. 2.

Experimental

The title compound was prepared according to the method described by Gowda *et al.* (2003). The purity of the compound was checked by determining its melting point. It was characterized by recording its infrared and NMR spectra. cuboid-like colourless single crystals of the title compound were obtained by slow evaporation from an ethanol solution of the compound (0.5 g in about 30 ml of ethanol) at room temperature.

Refinement

All H atoms were visible in difference maps and then treated as riding atoms with C—H distances of 0.93 Å (C-aromatic), 0.96 Å (C-methyl) and N—H = 0.86 Å. The $U_{\text{iso}}(\text{H})$ values were set at 1.2 U_{eq} (C-aromatic, N) and 1.5 U_{eq} (C-methyl).

Figures




Fig. 1. Molecular structure of the title compound showing the atom labelling scheme. Displacement ellipsoids are drawn at the 50% probability level. H atoms are represented as small spheres of arbitrary radii.




Fig. 2. Part of the crystal structure of the title compound. Molecular chains are generated by N—H···O hydrogen bonds which are shown by dashed lines. H atoms not involved in intermolecular bonding have been omitted.

supplementary materials

N-(2-Chlorophenyl)-4-methylbenzamide

Crystal data

C ₁₄ H ₁₂ ClNO	<i>F</i> (000) = 1024
<i>M_r</i> = 245.70	<i>D_x</i> = 1.293 Mg m ⁻³
Monoclinic, <i>P2₁/n</i>	Mo <i>Kα</i> radiation, λ = 0.71073 Å
Hall symbol: -P 2yn	Cell parameters from 13361 reflections
<i>a</i> = 9.6940 (5) Å	θ = 3.6–29.4°
<i>b</i> = 27.4495 (9) Å	μ = 0.29 mm ⁻¹
<i>c</i> = 9.9025 (4) Å	<i>T</i> = 293 K
β = 106.730 (5)°	Cuboid, colorless
<i>V</i> = 2523.48 (19) Å ³	0.97 × 0.13 × 0.10 mm
<i>Z</i> = 8	

Data collection

Oxford Diffraction Xcalibur Ruby Gemini diffractometer	7045 independent reflections
Radiation source: fine-focus sealed tube graphite	2850 reflections with $I > 2\sigma(I)$
Detector resolution: 10.4340 pixels mm ⁻¹	$R_{\text{int}} = 0.046$
ω scans	$\theta_{\text{max}} = 29.4^\circ$, $\theta_{\text{min}} = 3.6^\circ$
Absorption correction: analytical [CrysAlis RED (Oxford Diffraction, 2009) based on expressions derived by Clark & Reid (1995)]	$h = -13 \rightarrow 13$
$T_{\text{min}} = 0.957$, $T_{\text{max}} = 0.972$	$k = -37 \rightarrow 37$
47577 measured reflections	$l = -13 \rightarrow 12$

Refinement

Refinement on F^2	Primary atom site location: structure-invariant direct methods
Least-squares matrix: full	Secondary atom site location: difference Fourier map
$R[F^2 > 2\sigma(F^2)] = 0.043$	Hydrogen site location: inferred from neighbouring sites
$wR(F^2) = 0.126$	H-atom parameters constrained
$S = 0.85$	$w = 1/[\sigma^2(F_o^2) + (0.0704P)^2]$
7045 reflections	where $P = (F_o^2 + 2F_c^2)/3$
307 parameters	$(\Delta/\sigma)_{\text{max}} < 0.001$
0 restraints	$\Delta\rho_{\text{max}} = 0.24 \text{ e \AA}^{-3}$
	$\Delta\rho_{\text{min}} = -0.30 \text{ e \AA}^{-3}$

Special details

Geometry. All e.s.d.'s (except the e.s.d. in the dihedral angle between two l.s. planes) are estimated using the full covariance matrix. The cell e.s.d.'s are taken into account individually in the estimation of e.s.d.'s in distances, angles and torsion angles; correlations between e.s.d.'s in cell parameters are only used when they are defined by crystal symmetry. An approximate (isotropic) treatment of cell e.s.d.'s is used for estimating e.s.d.'s involving l.s. planes.

Refinement. Refinement of F^2 against ALL reflections. The weighted R -factor wR and goodness of fit S are based on F^2 , conventional R -factors R are based on F , with F set to zero for negative F^2 . The threshold expression of $F^2 > \sigma(F^2)$ is used only for calculating R -factors(gt) etc. and is not relevant to the choice of reflections for refinement. R -factors based on F^2 are statistically about twice as large as those based on F , and R -factors based on ALL data will be even larger.

Fractional atomic coordinates and isotropic or equivalent isotropic displacement parameters (\AA^2)

	<i>x</i>	<i>y</i>	<i>z</i>	$U_{\text{iso}}^*/U_{\text{eq}}$
C1	0.4465 (2)	0.18661 (6)	0.78372 (15)	0.0492 (4)
C2	0.5890 (2)	0.19536 (7)	0.85616 (18)	0.0567 (5)
C3	0.6412 (3)	0.24255 (8)	0.8811 (2)	0.0712 (6)
H3A	0.7374	0.2481	0.9296	0.085*
C4	0.5494 (3)	0.28101 (8)	0.8336 (2)	0.0794 (7)
H4A	0.5839	0.3128	0.8495	0.095*
C5	0.4082 (3)	0.27296 (8)	0.7632 (2)	0.0827 (7)
H5A	0.3463	0.2991	0.7320	0.099*
C6	0.3572 (2)	0.22594 (7)	0.7386 (2)	0.0676 (5)
H6A	0.2608	0.2207	0.6906	0.081*
C7	0.3647 (2)	0.10713 (6)	0.83990 (16)	0.0497 (4)
C8	0.30994 (18)	0.05833 (6)	0.78704 (16)	0.0466 (4)
C9	0.2405 (2)	0.04866 (7)	0.64638 (17)	0.0555 (5)
H9A	0.2249	0.0738	0.5808	0.067*
C10	0.1947 (2)	0.00242 (7)	0.6028 (2)	0.0653 (5)
H10A	0.1467	-0.0031	0.5083	0.078*
C11	0.2182 (2)	-0.03609 (7)	0.6964 (2)	0.0677 (5)
C12	0.2852 (2)	-0.02635 (7)	0.8368 (2)	0.0693 (6)
H12A	0.3009	-0.0516	0.9021	0.083*
C13	0.3290 (2)	0.02018 (7)	0.88200 (18)	0.0583 (5)
H13A	0.3719	0.0260	0.9774	0.070*
C14	0.1746 (4)	-0.08743 (8)	0.6468 (3)	0.1105 (10)
H14C	0.1994	-0.1092	0.7260	0.133*
H14B	0.2242	-0.0971	0.5800	0.133*
H14A	0.0725	-0.0886	0.6028	0.133*
C18	0.3176 (2)	0.05619 (6)	0.25095 (15)	0.0472 (4)
C19	0.1723 (2)	0.04811 (6)	0.18847 (17)	0.0556 (5)
C20	0.1129 (2)	0.00237 (8)	0.1859 (2)	0.0703 (6)
H20A	0.0150	-0.0025	0.1435	0.084*
C21	0.1995 (3)	-0.03562 (8)	0.2465 (2)	0.0755 (6)
H21A	0.1605	-0.0667	0.2437	0.091*
C22	0.3426 (3)	-0.02856 (7)	0.3110 (2)	0.0755 (6)
H22A	0.4003	-0.0546	0.3535	0.091*

supplementary materials

C23	0.4020 (2)	0.01741 (7)	0.31306 (19)	0.0624 (5)
H23A	0.4997	0.0221	0.3568	0.075*
C24	0.41322 (19)	0.13420 (6)	0.35262 (16)	0.0511 (4)
C25	0.44828 (19)	0.18501 (6)	0.32278 (15)	0.0476 (4)
C26	0.4024 (2)	0.20526 (6)	0.18911 (17)	0.0621 (5)
H26A	0.3556	0.1858	0.1129	0.074*
C27	0.4250 (2)	0.25370 (7)	0.1674 (2)	0.0683 (6)
H27A	0.3930	0.2663	0.0765	0.082*
C28	0.4937 (2)	0.28404 (7)	0.2766 (2)	0.0618 (5)
C29	0.5419 (2)	0.26368 (7)	0.4093 (2)	0.0707 (6)
H29A	0.5907	0.2832	0.4848	0.085*
C30	0.5195 (2)	0.21513 (7)	0.43291 (18)	0.0606 (5)
H30A	0.5526	0.2025	0.5237	0.073*
C31	0.5128 (3)	0.33727 (7)	0.2505 (3)	0.0915 (7)
H31C	0.5625	0.3529	0.3377	0.110*
H31B	0.4202	0.3522	0.2124	0.110*
H31A	0.5681	0.3407	0.1847	0.110*
N15	0.39382 (16)	0.13853 (5)	0.74748 (13)	0.0507 (4)
H15A	0.3799	0.1290	0.6618	0.061*
N32	0.37672 (17)	0.10304 (5)	0.24535 (13)	0.0534 (4)
H32A	0.3900	0.1120	0.1668	0.064*
O17	0.38313 (17)	0.11887 (5)	0.96324 (11)	0.0771 (5)
O34	0.4147 (2)	0.12111 (5)	0.47127 (12)	0.0924 (5)
Cl16	0.70621 (7)	0.14715 (2)	0.91490 (7)	0.0936 (2)
Cl33	0.06234 (7)	0.09610 (2)	0.10827 (7)	0.0956 (2)

Atomic displacement parameters (\AA^2)

	U^{11}	U^{22}	U^{33}	U^{12}	U^{13}	U^{23}
C1	0.0709 (13)	0.0453 (10)	0.0353 (8)	-0.0042 (9)	0.0213 (8)	0.0003 (7)
C2	0.0676 (14)	0.0507 (11)	0.0551 (10)	-0.0015 (10)	0.0228 (9)	-0.0006 (8)
C3	0.0802 (15)	0.0639 (14)	0.0698 (13)	-0.0166 (12)	0.0220 (11)	-0.0058 (10)
C4	0.114 (2)	0.0492 (13)	0.0813 (14)	-0.0168 (14)	0.0383 (15)	-0.0030 (11)
C5	0.105 (2)	0.0487 (13)	0.0948 (16)	0.0094 (13)	0.0286 (15)	0.0157 (11)
C6	0.0756 (14)	0.0593 (13)	0.0651 (12)	0.0025 (11)	0.0155 (10)	0.0129 (10)
C7	0.0668 (12)	0.0505 (10)	0.0354 (9)	-0.0018 (9)	0.0203 (8)	-0.0013 (7)
C8	0.0533 (11)	0.0490 (10)	0.0408 (9)	-0.0024 (8)	0.0187 (7)	-0.0022 (8)
C9	0.0668 (12)	0.0555 (11)	0.0452 (9)	-0.0073 (9)	0.0175 (9)	-0.0013 (8)
C10	0.0708 (14)	0.0716 (14)	0.0548 (11)	-0.0152 (11)	0.0202 (9)	-0.0157 (10)
C11	0.0751 (14)	0.0558 (12)	0.0801 (14)	-0.0189 (10)	0.0351 (11)	-0.0147 (11)
C12	0.0872 (16)	0.0539 (12)	0.0719 (13)	-0.0079 (11)	0.0309 (11)	0.0086 (10)
C13	0.0735 (13)	0.0547 (12)	0.0478 (9)	-0.0055 (10)	0.0192 (9)	0.0040 (9)
C14	0.145 (3)	0.0753 (16)	0.122 (2)	-0.0446 (17)	0.0568 (19)	-0.0319 (15)
C18	0.0676 (13)	0.0425 (10)	0.0359 (8)	-0.0033 (9)	0.0218 (8)	-0.0009 (7)
C19	0.0702 (14)	0.0515 (11)	0.0455 (9)	-0.0003 (10)	0.0172 (9)	0.0017 (8)
C20	0.0780 (15)	0.0693 (14)	0.0619 (12)	-0.0175 (12)	0.0171 (10)	-0.0064 (10)
C21	0.106 (2)	0.0494 (12)	0.0799 (14)	-0.0162 (13)	0.0409 (14)	-0.0102 (11)
C22	0.103 (2)	0.0470 (13)	0.0871 (14)	0.0161 (12)	0.0443 (14)	0.0128 (10)

C23	0.0660 (13)	0.0630 (13)	0.0610 (11)	0.0055 (11)	0.0228 (9)	0.0065 (9)
C24	0.0676 (12)	0.0538 (11)	0.0357 (9)	-0.0019 (9)	0.0210 (8)	-0.0024 (8)
C25	0.0581 (11)	0.0484 (10)	0.0398 (9)	-0.0006 (8)	0.0194 (8)	-0.0032 (8)
C26	0.0951 (16)	0.0470 (11)	0.0432 (10)	-0.0061 (10)	0.0185 (9)	-0.0058 (8)
C27	0.1027 (17)	0.0506 (12)	0.0544 (11)	-0.0028 (11)	0.0273 (11)	0.0041 (9)
C28	0.0671 (13)	0.0489 (11)	0.0769 (13)	-0.0062 (10)	0.0326 (10)	-0.0050 (10)
C29	0.0736 (15)	0.0660 (13)	0.0696 (13)	-0.0172 (11)	0.0159 (11)	-0.0206 (11)
C30	0.0692 (13)	0.0615 (12)	0.0469 (10)	-0.0078 (10)	0.0100 (9)	-0.0059 (9)
C31	0.109 (2)	0.0522 (13)	0.1262 (19)	-0.0105 (13)	0.0550 (16)	-0.0083 (13)
N15	0.0737 (10)	0.0472 (8)	0.0339 (7)	-0.0094 (7)	0.0200 (7)	-0.0038 (6)
N32	0.0805 (11)	0.0490 (9)	0.0340 (7)	-0.0119 (7)	0.0215 (7)	-0.0006 (6)
O17	0.1376 (14)	0.0618 (8)	0.0409 (7)	-0.0250 (8)	0.0399 (7)	-0.0084 (6)
O34	0.1802 (17)	0.0653 (9)	0.0414 (7)	-0.0260 (10)	0.0475 (9)	-0.0063 (6)
Cl16	0.0746 (4)	0.0705 (4)	0.1285 (5)	0.0101 (3)	0.0178 (3)	0.0039 (3)
Cl33	0.0833 (4)	0.0885 (4)	0.1050 (5)	0.0137 (3)	0.0112 (3)	0.0323 (3)

Geometric parameters (Å, °)

C1—C6	1.375 (3)	C18—C19	1.384 (3)
C1—C2	1.382 (3)	C18—N32	1.415 (2)
C1—N15	1.423 (2)	C19—C20	1.379 (3)
C2—C3	1.387 (3)	C19—Cl33	1.7365 (19)
C2—Cl16	1.732 (2)	C20—C21	1.365 (3)
C3—C4	1.374 (3)	C20—H20A	0.9300
C3—H3A	0.9300	C21—C22	1.364 (3)
C4—C5	1.364 (3)	C21—H21A	0.9300
C4—H4A	0.9300	C22—C23	1.385 (3)
C5—C6	1.378 (3)	C22—H22A	0.9300
C5—H5A	0.9300	C23—H23A	0.9300
C6—H6A	0.9300	C24—O34	1.2248 (17)
C7—O17	1.2255 (17)	C24—N32	1.330 (2)
C7—N15	1.345 (2)	C24—C25	1.485 (2)
C7—C8	1.480 (2)	C25—C30	1.384 (2)
C8—C13	1.384 (2)	C25—C26	1.386 (2)
C8—C9	1.387 (2)	C26—C27	1.374 (3)
C9—C10	1.373 (3)	C26—H26A	0.9300
C9—H9A	0.9300	C27—C28	1.375 (3)
C10—C11	1.381 (3)	C27—H27A	0.9300
C10—H10A	0.9300	C28—C29	1.380 (3)
C11—C12	1.380 (3)	C28—C31	1.505 (3)
C11—C14	1.512 (3)	C29—C30	1.381 (3)
C12—C13	1.379 (3)	C29—H29A	0.9300
C12—H12A	0.9300	C30—H30A	0.9300
C13—H13A	0.9300	C31—H31C	0.9600
C14—H14C	0.9600	C31—H31B	0.9600
C14—H14B	0.9600	C31—H31A	0.9600
C14—H14A	0.9600	N15—H15A	0.8600
C18—C23	1.375 (2)	N32—H32A	0.8600
C6—C1—C2	118.26 (17)	C20—C19—C18	121.37 (18)

supplementary materials

C6—C1—N15	119.97 (17)	C20—C19—Cl33	118.95 (17)
C2—C1—N15	121.63 (16)	C18—C19—Cl33	119.67 (14)
C1—C2—C3	120.95 (18)	C21—C20—C19	119.1 (2)
C1—C2—Cl16	120.17 (14)	C21—C20—H20A	120.4
C3—C2—Cl16	118.88 (17)	C19—C20—H20A	120.4
C4—C3—C2	119.3 (2)	C22—C21—C20	120.8 (2)
C4—C3—H3A	120.3	C22—C21—H21A	119.6
C2—C3—H3A	120.3	C20—C21—H21A	119.6
C5—C4—C3	120.4 (2)	C21—C22—C23	119.9 (2)
C5—C4—H4A	119.8	C21—C22—H22A	120.0
C3—C4—H4A	119.8	C23—C22—H22A	120.0
C4—C5—C6	119.9 (2)	C18—C23—C22	120.5 (2)
C4—C5—H5A	120.1	C18—C23—H23A	119.8
C6—C5—H5A	120.1	C22—C23—H23A	119.8
C1—C6—C5	121.2 (2)	O34—C24—N32	120.29 (15)
C1—C6—H6A	119.4	O34—C24—C25	121.75 (14)
C5—C6—H6A	119.4	N32—C24—C25	117.94 (13)
O17—C7—N15	120.71 (15)	C30—C25—C26	117.58 (16)
O17—C7—C8	121.67 (14)	C30—C25—C24	119.67 (14)
N15—C7—C8	117.62 (13)	C26—C25—C24	122.55 (15)
C13—C8—C9	117.97 (16)	C27—C26—C25	121.04 (16)
C13—C8—C7	118.71 (14)	C27—C26—H26A	119.5
C9—C8—C7	123.32 (15)	C25—C26—H26A	119.5
C10—C9—C8	120.80 (17)	C26—C27—C28	121.73 (17)
C10—C9—H9A	119.6	C26—C27—H27A	119.1
C8—C9—H9A	119.6	C28—C27—H27A	119.1
C9—C10—C11	121.40 (17)	C27—C28—C29	117.26 (17)
C9—C10—H10A	119.3	C27—C28—C31	120.59 (19)
C11—C10—H10A	119.3	C29—C28—C31	122.15 (19)
C12—C11—C10	117.82 (17)	C28—C29—C30	121.68 (17)
C12—C11—C14	121.0 (2)	C28—C29—H29A	119.2
C10—C11—C14	121.2 (2)	C30—C29—H29A	119.2
C13—C12—C11	121.17 (17)	C29—C30—C25	120.69 (17)
C13—C12—H12A	119.4	C29—C30—H30A	119.7
C11—C12—H12A	119.4	C25—C30—H30A	119.7
C12—C13—C8	120.78 (17)	C28—C31—H31C	109.5
C12—C13—H13A	119.6	C28—C31—H31B	109.5
C8—C13—H13A	119.6	H31C—C31—H31B	109.5
C11—C14—H14C	109.5	C28—C31—H31A	109.5
C11—C14—H14B	109.5	H31C—C31—H31A	109.5
H14C—C14—H14B	109.5	H31B—C31—H31A	109.5
C11—C14—H14A	109.5	C7—N15—C1	123.57 (12)
H14C—C14—H14A	109.5	C7—N15—H15A	118.2
H14B—C14—H14A	109.5	C1—N15—H15A	118.2
C23—C18—C19	118.30 (16)	C24—N32—C18	124.80 (13)
C23—C18—N32	121.70 (17)	C24—N32—H32A	117.6
C19—C18—N32	119.96 (16)	C18—N32—H32A	117.6
C6—C1—C2—C3	-0.8 (2)	C18—C19—C20—C21	0.0 (3)
N15—C1—C2—C3	174.95 (14)	Cl33—C19—C20—C21	178.76 (15)

C6—C1—C2—Cl16	−179.78 (13)	C19—C20—C21—C22	1.1 (3)
N15—C1—C2—Cl16	−4.0 (2)	C20—C21—C22—C23	−1.3 (3)
C1—C2—C3—C4	0.3 (3)	C19—C18—C23—C22	0.9 (2)
Cl16—C2—C3—C4	179.27 (15)	N32—C18—C23—C22	−176.82 (15)
C2—C3—C4—C5	0.4 (3)	C21—C22—C23—C18	0.3 (3)
C3—C4—C5—C6	−0.5 (3)	O34—C24—C25—C30	17.0 (3)
C2—C1—C6—C5	0.7 (3)	N32—C24—C25—C30	−164.36 (17)
N15—C1—C6—C5	−175.18 (16)	O34—C24—C25—C26	−157.66 (19)
C4—C5—C6—C1	0.0 (3)	N32—C24—C25—C26	21.0 (3)
O17—C7—C8—C13	−23.7 (3)	C30—C25—C26—C27	−1.0 (3)
N15—C7—C8—C13	156.29 (17)	C24—C25—C26—C27	173.79 (19)
O17—C7—C8—C9	156.49 (18)	C25—C26—C27—C28	0.0 (3)
N15—C7—C8—C9	−23.5 (3)	C26—C27—C28—C29	1.2 (3)
C13—C8—C9—C10	−1.2 (3)	C26—C27—C28—C31	−178.1 (2)
C7—C8—C9—C10	178.60 (17)	C27—C28—C29—C30	−1.5 (3)
C8—C9—C10—C11	−1.2 (3)	C31—C28—C29—C30	177.8 (2)
C9—C10—C11—C12	2.3 (3)	C28—C29—C30—C25	0.5 (3)
C9—C10—C11—C14	−176.7 (2)	C26—C25—C30—C29	0.7 (3)
C10—C11—C12—C13	−0.9 (3)	C24—C25—C30—C29	−174.21 (18)
C14—C11—C12—C13	178.0 (2)	O17—C7—N15—C1	−0.2 (3)
C11—C12—C13—C8	−1.4 (3)	C8—C7—N15—C1	179.78 (16)
C9—C8—C13—C12	2.5 (3)	C6—C1—N15—C7	−107.11 (19)
C7—C8—C13—C12	−177.30 (18)	C2—C1—N15—C7	77.2 (2)
C23—C18—C19—C20	−1.0 (2)	O34—C24—N32—C18	9.7 (3)
N32—C18—C19—C20	176.71 (15)	C25—C24—N32—C18	−168.96 (16)
C23—C18—C19—Cl33	−179.72 (12)	C23—C18—N32—C24	−77.9 (2)
N32—C18—C19—Cl33	−2.0 (2)	C19—C18—N32—C24	104.46 (19)

Hydrogen-bond geometry (Å, °)

D—H···A	D—H	H···A	D···A	D—H···A
N15—H15A···O34	0.86	2.02	2.8408 (16)	159.
N32—H32A···O17 ⁱ	0.86	2.01	2.8455 (16)	165.

Symmetry codes: (i) $x, y, z-1$.

supplementary materials

Fig. 1




Fig. 2

