

## 8-Thia-1,6-diazabicyclo[4.3.0]nonane-7,9-dione

Bin-bin Xi,<sup>a</sup> Lu Shi,<sup>a</sup> Hai Huang,<sup>a</sup> Qin Wang<sup>a</sup> and Hong-Jun Zhu<sup>b\*</sup>

<sup>a</sup>Department of Organic Chemistry, College of Science, Nanjing University of Technology, Nanjing 210009, People's Republic of China, and <sup>b</sup>Department of Applied Chemistry, College of Science, Nanjing University of Technology, Nanjing 210009, People's Republic of China  
Correspondence e-mail: zhuhj@njut.edu.cn


Received 24 July 2011; accepted 21 September 2011

Key indicators: single-crystal X-ray study;  $T = 293\text{ K}$ ; mean  $\sigma(\text{C}-\text{C}) = 0.005\text{ \AA}$ ;  $R$  factor = 0.044;  $wR$  factor = 0.125; data-to-parameter ratio = 14.0.

There are two independent molecules, *A* and *B*, in the asymmetric unit of the title compound,  $\text{C}_6\text{H}_8\text{N}_2\text{O}_2\text{S}$ . In the crystal, pairs of intermolecular  $\text{S}\cdots\text{O}$  contacts [3.286 (1)  $\text{\AA}$ ] link the *B* molecules into inversion dimers.

### Related literature

For applications of the title compound, see: Yamaguchi *et al.* (1989). For the synthesis, see: Zhu *et al.* (2011). For bond-length data, see: Allen *et al.* (1987). For a review of carbonyl–carbonyl interactions, see: Allen *et al.* (1998).



### Experimental

#### Crystal data

$\text{C}_6\text{H}_8\text{N}_2\text{O}_2\text{S}$

$M_r = 172.20$

Triclinic,  $P\bar{1}$   
 $a = 7.8400 (16)\text{ \AA}$   
 $b = 10.464 (2)\text{ \AA}$   
 $c = 10.514 (2)\text{ \AA}$   
 $\alpha = 63.84 (3)^\circ$   
 $\beta = 79.62 (3)^\circ$   
 $\gamma = 89.42 (3)^\circ$

$V = 759.2 (3)\text{ \AA}^3$   
 $Z = 4$   
Mo  $K\alpha$  radiation  
 $\mu = 0.37\text{ mm}^{-1}$   
 $T = 293\text{ K}$   
 $0.30 \times 0.20 \times 0.10\text{ mm}$

#### Data collection

Enraf–Nonius CAD-4  
diffractometer  
Absorption correction:  $\psi$  scan  
(North *et al.*, 1968)  
 $T_{\min} = 0.896$ ,  $T_{\max} = 0.964$   
3018 measured reflections

2795 independent reflections  
2092 reflections with  $I > 2\sigma(I)$   
 $R_{\text{int}} = 0.041$   
3 standard reflections every 200  
reflections  
intensity decay: 1%

#### Refinement

$R[F^2 > 2\sigma(F^2)] = 0.044$   
 $wR(F^2) = 0.125$   
 $S = 1.00$   
2795 reflections

200 parameters  
H-atom parameters constrained  
 $\Delta\rho_{\max} = 0.22\text{ e \AA}^{-3}$   
 $\Delta\rho_{\min} = -0.32\text{ e \AA}^{-3}$

Data collection: *CAD-4 EXPRESS* (Enraf–Nonius, 1994); cell refinement: *CAD-4 EXPRESS*; data reduction: *XCAD4* (Harms & Wocadlo, 1995); program(s) used to solve structure: *SHELXS97* (Sheldrick, 2008); program(s) used to refine structure: *SHELXL97* (Sheldrick, 2008); molecular graphics: *SHELXTL* (Sheldrick, 2008); software used to prepare material for publication: *SHELXTL*.

The authors thank the Center of Testing and Analysis, Nanjing University, for support.

Supplementary data and figures for this paper are available from the IUCr electronic archives (Reference: LX2196).

### References

- Allen, F. H., Baalham, C. A., Lommerse, J. P. M. & Raithby, P. R. (1998). *Acta Cryst. B* **54**, 320–329.
- Allen, F. H., Kennard, O., Watson, D. G., Brammer, L., Orpen, A. G. & Taylor, R. (1987). *J. Chem. Soc. Perkin Trans. 2*, pp. S1–19.
- Enraf–Nonius (1994). *CAD-4 EXPRESS*. Enraf–Nonius, Delft, The Netherlands.
- Harms, K. & Wocadlo, S. (1995). *XCAD4*. University of Marburg, Germany.
- North, A. C. T., Phillips, D. C. & Mathews, F. S. (1968). *Acta Cryst. A* **24**, 351–359.
- Sheldrick, G. M. (2008). *Acta Cryst. A* **64**, 112–122.
- Yamaguchi, M., Suzuki, C., Matsunari, K., Miyazawa, T. & Nakamura, Y. (1989). EP Patent 0312064A2.
- Zhu, H. J., Xi, B. B., Feng, M. L., Wang, K., Li, Y. F., Shi, L. & Chen, C. (2011). CN Patent Appl. CN 201110091918.8.

## **supplementary materials**

*Acta Cryst.* (2011). E67, o2771 [doi:10.1107/S1600536811038785]

## 8-Thia-1,6-diazabicyclo[4.3.0]nonane-7,9-dione

B. Xi, L. Shi, H. Huang, Q. Wang and H.-J. Zhu

### Comment

The title compound, 8-thia-1,6-diazabicyclo[4.3.0]nonane-7,9-dione, is an important intermediate for a kind of manufacturing agrochemical, especially for Fluthiacet-methyl as a super-effective, wide-spectral and safe herbicide (Yamaguchi *et al.*, 1989). We report herein the crystal structure of the title compound.

The asymmetric unit of the title compound is shown in Fig. 1 and there are two independent unique molecules [labeled A & B]. The bond lengths and angles are within normal ranges (Allen *et al.*, 1987). The crystal packing (Fig. 2) is stabilized by an intermolecular S···O interaction between the sulfur and the O atom of the carbonyl group interpreted as similar to type-II carbonyl–carbonyl interaction (Allen *et al.*, 1998), with  $S1\cdots O2^{\dagger}$  distance of 3.286 (1) Å.

### Experimental

The title compound was prepared according to reported in literature (Zhu *et al.*, 2011). Single crystals suitable for X-ray diffraction were prepared by slow evaporation of a solution of the title compound in ethanol at room temperature for ca. 6 d.

### Refinement

All H atoms were positioned geometrically and refined using a riding model, with C—H = 0.95 Å and  $U_{\text{iso}}(\text{H})=1.5U_{\text{eq}}(\text{C})$ .

### Figures




Fig. 1. The molecular structure of the title compound with the atom numbering scheme. Displacement ellipsoids are drawn at the 50% probability level. H atoms are presented as small spheres of arbitrary radius.




Fig. 2. A view of the S···O interactions (dotted lines) in the crystal structure of the title compound. [Symmetry codes: (i)  $-x, -y + 2, -z$ .]

# supplementary materials

---

## 8-Thia-1,6-diazabicyclo[4.3.0]nonane-7,9-dione

### Crystal data

C <sub>6</sub> H <sub>8</sub> N <sub>2</sub> O <sub>2</sub> S	Z = 4
M <sub>r</sub> = 172.20	F(000) = 360
Triclinic, PT	D <sub>x</sub> = 1.507 Mg m <sup>-3</sup>
Hall symbol: -P 1	Mo K $\alpha$ radiation, $\lambda$ = 0.71073 Å
a = 7.8400 (16) Å	Cell parameters from 25 reflections
b = 10.464 (2) Å	$\theta$ = 10–13°
c = 10.514 (2) Å	$\mu$ = 0.37 mm <sup>-1</sup>
$\alpha$ = 63.84 (3)°	T = 293 K
$\beta$ = 79.62 (3)°	Block, colourless
$\gamma$ = 89.42 (3)°	0.30 × 0.20 × 0.10 mm
V = 759.2 (3) Å <sup>3</sup>	

### Data collection

Enraf–Nonius CAD-4 diffractometer	2092 reflections with $I > 2\sigma(I)$
Radiation source: fine-focus sealed tube	$R_{\text{int}} = 0.041$
graphite	$\theta_{\text{max}} = 25.4^\circ$ , $\theta_{\text{min}} = 2.2^\circ$
$\omega/2\theta$ scans	$h = 0 \rightarrow 9$
Absorption correction: $\psi$ scan (North <i>et al.</i> , 1968)	$k = -12 \rightarrow 12$
$T_{\text{min}} = 0.896$ , $T_{\text{max}} = 0.964$	$l = -12 \rightarrow 12$
3018 measured reflections	3 standard reflections every 200 reflections
2795 independent reflections	intensity decay: 1%

### Refinement

Refinement on $F^2$	Secondary atom site location: difference Fourier map
Least-squares matrix: full	Hydrogen site location: difference Fourier map
$R[F^2 > 2\sigma(F^2)] = 0.044$	H-atom parameters constrained
$wR(F^2) = 0.125$	$w = 1/[\sigma^2(F_o^2) + (0.077P)^2]$
$S = 1.00$	where $P = (F_o^2 + 2F_c^2)/3$
2795 reflections	$(\Delta/\sigma)_{\text{max}} < 0.001$
200 parameters	$\Delta\rho_{\text{max}} = 0.22 \text{ e \AA}^{-3}$
0 restraints	$\Delta\rho_{\text{min}} = -0.32 \text{ e \AA}^{-3}$
Primary atom site location: structure-invariant direct methods	Extinction correction: <i>SHELXL</i> , $F_c^* = kFc[1 + 0.001xFc^2\lambda^3/\sin(2\theta)]^{-1/4}$
	Extinction coefficient: 0.327 (16)

*Special details*

**Geometry.** All e.s.d.'s (except the e.s.d. in the dihedral angle between two l.s. planes) are estimated using the full covariance matrix. The cell e.s.d.'s are taken into account individually in the estimation of e.s.d.'s in distances, angles and torsion angles; correlations between e.s.d.'s in cell parameters are only used when they are defined by crystal symmetry. An approximate (isotropic) treatment of cell e.s.d.'s is used for estimating e.s.d.'s involving l.s. planes.

**Refinement.** Refinement of  $F^2$  against ALL reflections. The weighted  $R$ -factor  $wR$  and goodness of fit  $S$  are based on  $F^2$ , conventional  $R$ -factors  $R$  are based on  $F$ , with  $F$  set to zero for negative  $F^2$ . The threshold expression of  $F^2 > \sigma(F^2)$  is used only for calculating  $R$ -factors(gt) etc. and is not relevant to the choice of reflections for refinement.  $R$ -factors based on  $F^2$  are statistically about twice as large as those based on  $F$ , and  $R$ -factors based on ALL data will be even larger.

*Fractional atomic coordinates and isotropic or equivalent isotropic displacement parameters ( $\text{\AA}^2$ )*

	<i>x</i>	<i>y</i>	<i>z</i>	$U_{\text{iso}}^*/U_{\text{eq}}$
S1	-0.09731 (11)	0.82576 (8)	0.04086 (8)	0.0598 (3)
O1	-0.3185 (3)	0.5944 (3)	0.1791 (3)	0.0737 (7)
O2	0.2479 (3)	0.8734 (2)	-0.0219 (2)	0.0632 (6)
N1	-0.0292 (2)	0.5625 (2)	0.1507 (2)	0.0416 (5)
N2	0.1318 (3)	0.6432 (2)	0.1083 (2)	0.0411 (5)
C1	0.2837 (3)	0.5721 (3)	0.0768 (3)	0.0464 (6)
H1A	0.3892	0.6300	0.0591	0.056*
H1B	0.2803	0.5604	-0.0092	0.056*
C2	0.2842 (4)	0.4278 (3)	0.2034 (3)	0.0547 (7)
H2A	0.3027	0.4404	0.2860	0.066*
H2B	0.3789	0.3771	0.1791	0.066*
C3	0.1134 (4)	0.3408 (3)	0.2418 (3)	0.0529 (7)
H3A	0.1002	0.3205	0.1623	0.063*
H3B	0.1137	0.2505	0.3261	0.063*
C4	-0.0370 (4)	0.4199 (3)	0.2726 (3)	0.0491 (7)
H4A	-0.1456	0.3669	0.2878	0.059*
H4B	-0.0326	0.4290	0.3599	0.059*
C5	-0.1691 (3)	0.6421 (3)	0.1356 (3)	0.0497 (7)
C6	0.1243 (4)	0.7859 (3)	0.0365 (3)	0.0462 (6)
S2	0.16047 (9)	0.00770 (8)	0.62686 (8)	0.0517 (3)
O3	0.3930 (3)	-0.1856 (2)	0.6988 (2)	0.0625 (6)
O4	0.1494 (2)	0.2796 (2)	0.5788 (2)	0.0585 (5)
N3	0.4698 (2)	0.0396 (2)	0.6615 (2)	0.0396 (5)
N4	0.3934 (3)	0.1675 (2)	0.6431 (2)	0.0379 (5)
C7	0.5149 (3)	0.2948 (3)	0.5815 (3)	0.0462 (6)
H7A	0.4534	0.3749	0.5837	0.055*
H7B	0.5648	0.3187	0.4817	0.055*
C8	0.6568 (4)	0.2672 (3)	0.6669 (3)	0.0567 (8)
H8A	0.6083	0.2567	0.7629	0.068*
H8B	0.7430	0.3483	0.6203	0.068*
C9	0.7431 (3)	0.1341 (3)	0.6786 (3)	0.0541 (7)
H9A	0.8298	0.1158	0.7384	0.065*
H9B	0.8015	0.1481	0.5833	0.065*

## supplementary materials

---

C10	0.6102 (3)	0.0070 (3)	0.7438 (3)	0.0502 (7)
H10A	0.6655	-0.0756	0.7418	0.060*
H10B	0.5628	-0.0153	0.8437	0.060*
C11	0.3614 (3)	-0.0634 (3)	0.6686 (3)	0.0433 (6)
C12	0.2314 (3)	0.1752 (3)	0.6121 (3)	0.0412 (6)

*Atomic displacement parameters ( $\text{\AA}^2$ )*

	$U^{11}$	$U^{22}$	$U^{33}$	$U^{12}$	$U^{13}$	$U^{23}$
S1	0.0698 (5)	0.0513 (5)	0.0626 (5)	0.0262 (4)	-0.0203 (4)	-0.0268 (4)
O1	0.0382 (12)	0.0922 (17)	0.0976 (17)	0.0078 (11)	-0.0077 (11)	-0.0508 (14)
O2	0.0749 (14)	0.0403 (11)	0.0593 (13)	-0.0102 (10)	0.0012 (11)	-0.0138 (10)
N1	0.0357 (11)	0.0393 (11)	0.0448 (12)	0.0011 (9)	-0.0052 (9)	-0.0153 (10)
N2	0.0369 (11)	0.0349 (11)	0.0453 (12)	0.0006 (9)	-0.0038 (9)	-0.0139 (9)
C1	0.0375 (14)	0.0472 (15)	0.0502 (15)	0.0045 (11)	0.0008 (11)	-0.0213 (13)
C2	0.0508 (17)	0.0486 (16)	0.0617 (18)	0.0155 (13)	-0.0147 (14)	-0.0209 (14)
C3	0.073 (2)	0.0344 (14)	0.0467 (15)	0.0053 (13)	-0.0151 (14)	-0.0130 (12)
C4	0.0545 (16)	0.0431 (15)	0.0406 (14)	-0.0101 (12)	-0.0044 (12)	-0.0120 (12)
C5	0.0418 (16)	0.0638 (18)	0.0531 (16)	0.0122 (13)	-0.0109 (13)	-0.0343 (15)
C6	0.0585 (17)	0.0418 (15)	0.0362 (13)	0.0039 (13)	-0.0048 (12)	-0.0172 (12)
S2	0.0443 (4)	0.0511 (4)	0.0589 (5)	-0.0031 (3)	-0.0175 (3)	-0.0208 (3)
O3	0.0769 (15)	0.0407 (11)	0.0712 (14)	0.0081 (10)	-0.0188 (11)	-0.0247 (10)
O4	0.0494 (11)	0.0536 (12)	0.0682 (13)	0.0176 (10)	-0.0212 (10)	-0.0200 (10)
N3	0.0358 (11)	0.0354 (11)	0.0451 (12)	0.0058 (9)	-0.0080 (9)	-0.0159 (9)
N4	0.0328 (10)	0.0335 (11)	0.0445 (12)	0.0028 (8)	-0.0076 (9)	-0.0150 (9)
C7	0.0446 (15)	0.0356 (13)	0.0543 (16)	-0.0040 (11)	-0.0011 (12)	-0.0195 (12)
C8	0.0418 (15)	0.0667 (19)	0.0708 (19)	-0.0067 (14)	-0.0050 (14)	-0.0409 (16)
C9	0.0312 (13)	0.076 (2)	0.0632 (18)	0.0066 (13)	-0.0108 (13)	-0.0377 (16)
C10	0.0381 (14)	0.0576 (16)	0.0520 (16)	0.0119 (12)	-0.0132 (12)	-0.0204 (14)
C11	0.0467 (15)	0.0393 (15)	0.0413 (14)	0.0016 (12)	-0.0056 (11)	-0.0167 (11)
C12	0.0386 (13)	0.0453 (14)	0.0359 (13)	0.0049 (12)	-0.0095 (11)	-0.0138 (11)

*Geometric parameters ( $\text{\AA}$ ,  $^\circ$ )*

S1—C5	1.771 (3)	S2—C12	1.776 (3)
S1—C6	1.780 (3)	S2—C11	1.779 (3)
O1—C5	1.203 (3)	O3—C11	1.208 (3)
O2—C6	1.209 (3)	O4—C12	1.208 (3)
N1—C5	1.360 (3)	N3—C11	1.347 (3)
N1—N2	1.411 (3)	N3—N4	1.409 (3)
N1—C4	1.470 (3)	N3—C10	1.467 (3)
N2—C6	1.351 (3)	N4—C12	1.360 (3)
N2—C1	1.463 (3)	N4—C7	1.464 (3)
C1—C2	1.510 (4)	C7—C8	1.501 (4)
C1—H1A	0.9700	C7—H7A	0.9700
C1—H1B	0.9700	C7—H7B	0.9700
C2—C3	1.514 (4)	C8—C9	1.508 (4)
C2—H2A	0.9700	C8—H8A	0.9700
C2—H2B	0.9700	C8—H8B	0.9700


C3—C4	1.503 (4)	C9—C10	1.514 (4)
C3—H3A	0.9700	C9—H9A	0.9700
C3—H3B	0.9700	C9—H9B	0.9700
C4—H4A	0.9700	C10—H10A	0.9700
C4—H4B	0.9700	C10—H10B	0.9700
C5—S1—C6	91.50 (13)	C12—S2—C11	91.50 (12)
C5—N1—N2	114.0 (2)	C11—N3—N4	114.76 (19)
C5—N1—C4	121.3 (2)	C11—N3—C10	122.2 (2)
N2—N1—C4	114.5 (2)	N4—N3—C10	114.6 (2)
C6—N2—N1	114.8 (2)	C12—N4—N3	114.5 (2)
C6—N2—C1	121.8 (2)	C12—N4—C7	121.1 (2)
N1—N2—C1	115.11 (19)	N3—N4—C7	115.35 (19)
N2—C1—C2	109.5 (2)	N4—C7—C8	109.7 (2)
N2—C1—H1A	109.8	N4—C7—H7A	109.7
C2—C1—H1A	109.8	C8—C7—H7A	109.7
N2—C1—H1B	109.8	N4—C7—H7B	109.7
C2—C1—H1B	109.8	C8—C7—H7B	109.7
H1A—C1—H1B	108.2	H7A—C7—H7B	108.2
C1—C2—C3	110.6 (2)	C7—C8—C9	111.1 (2)
C1—C2—H2A	109.5	C7—C8—H8A	109.4
C3—C2—H2A	109.5	C9—C8—H8A	109.4
C1—C2—H2B	109.5	C7—C8—H8B	109.4
C3—C2—H2B	109.5	C9—C8—H8B	109.4
H2A—C2—H2B	108.1	H8A—C8—H8B	108.0
C4—C3—C2	110.8 (2)	C8—C9—C10	110.7 (2)
C4—C3—H3A	109.5	C8—C9—H9A	109.5
C2—C3—H3A	109.5	C10—C9—H9A	109.5
C4—C3—H3B	109.5	C8—C9—H9B	109.5
C2—C3—H3B	109.5	C10—C9—H9B	109.5
H3A—C3—H3B	108.1	H9A—C9—H9B	108.1
N1—C4—C3	109.9 (2)	N3—C10—C9	109.7 (2)
N1—C4—H4A	109.7	N3—C10—H10A	109.7
C3—C4—H4A	109.7	C9—C10—H10A	109.7
N1—C4—H4B	109.7	N3—C10—H10B	109.7
C3—C4—H4B	109.7	C9—C10—H10B	109.7
H4A—C4—H4B	108.2	H10A—C10—H10B	108.2
O1—C5—N1	125.0 (3)	O3—C11—N3	126.2 (3)
O1—C5—S1	125.5 (2)	O3—C11—S2	124.5 (2)
N1—C5—S1	109.5 (2)	N3—C11—S2	109.34 (18)
O2—C6—N2	125.7 (3)	O4—C12—N4	125.5 (2)
O2—C6—S1	125.1 (2)	O4—C12—S2	125.4 (2)
N2—C6—S1	109.2 (2)	N4—C12—S2	109.10 (18)
C5—N1—N2—C6	-11.3 (3)	C11—N3—N4—C12	-10.1 (3)
C4—N1—N2—C6	-157.2 (2)	C10—N3—N4—C12	-159.0 (2)
C5—N1—N2—C1	-160.4 (2)	C11—N3—N4—C7	-157.8 (2)
C4—N1—N2—C1	53.6 (3)	C10—N3—N4—C7	53.3 (3)
C6—N2—C1—C2	160.1 (2)	C12—N4—C7—C8	162.2 (2)
N1—N2—C1—C2	-53.2 (3)	N3—N4—C7—C8	-52.4 (3)

## supplementary materials

---

N2—C1—C2—C3	54.1 (3)	N4—C7—C8—C9	53.6 (3)
C1—C2—C3—C4	-56.3 (3)	C7—C8—C9—C10	-56.1 (3)
C5—N1—C4—C3	164.3 (2)	C11—N3—C10—C9	161.2 (2)
N2—N1—C4—C3	-52.4 (3)	N4—N3—C10—C9	-52.4 (3)
C2—C3—C4—N1	53.8 (3)	C8—C9—C10—N3	53.8 (3)
N2—N1—C5—O1	-171.7 (3)	N4—N3—C11—O3	-171.8 (2)
C4—N1—C5—O1	-28.3 (4)	C10—N3—C11—O3	-25.6 (4)
N2—N1—C5—S1	8.8 (3)	N4—N3—C11—S2	7.9 (3)
C4—N1—C5—S1	152.26 (19)	C10—N3—C11—S2	154.20 (19)
C6—S1—C5—O1	176.8 (3)	C12—S2—C11—O3	176.4 (2)
C6—S1—C5—N1	-3.76 (19)	C12—S2—C11—N3	-3.33 (19)
N1—N2—C6—O2	-172.2 (2)	N3—N4—C12—O4	-172.5 (2)
C1—N2—C6—O2	-25.3 (4)	C7—N4—C12—O4	-26.9 (4)
N1—N2—C6—S1	7.8 (3)	N3—N4—C12—S2	6.9 (3)
C1—N2—C6—S1	154.63 (19)	C7—N4—C12—S2	152.55 (18)
C5—S1—C6—O2	177.7 (3)	C11—S2—C12—O4	177.4 (2)
C5—S1—C6—N2	-2.26 (19)	C11—S2—C12—N4	-2.03 (19)

Fig. 1



## supplementary materials

---

Fig. 2

