

## 1-[*(Z*)-8-(4-Chlorophenoxy)-3-(2,4-di-fluorophenyl)-4-oxaocta-2-en-2-yl]-1*H*-1,2,4-triazol-4-i um nitrate

Jian-long Chen,<sup>a\*</sup> Kai Wang,<sup>b</sup> Yan Shen,<sup>a</sup> Li-li Ren<sup>a</sup> and Guo-guang Chen<sup>a</sup>

<sup>a</sup>College of Pharmaceutical Science, Nanjing University of Technology, Xinmofan Road No.5 Nanjing, Nanjing 210009, People's Republic of China, and <sup>b</sup>Jiangsu Engineering Technology Research Center of Polypeptide Pharmaceuticals, College of Life Science and Pharmaceutical Engineering, Nanjing University of Technology, Xinmofan Road No.5 Nanjing, Nanjing 210009, People's Republic of China  
Correspondence e-mail: guoguangchen@163.com


Received 7 September 2011; accepted 21 September 2011

Key indicators: single-crystal X-ray study;  $T = 293\text{ K}$ ; mean  $\sigma(\text{C}-\text{C}) = 0.005\text{ \AA}$ ;  $R$  factor = 0.066;  $wR$  factor = 0.140; data-to-parameter ratio = 14.1.

In the title compound  $\text{C}_{21}\text{H}_{21}\text{ClF}_2\text{N}_3\text{O}_2^+\cdot\text{NO}_3^-$ , the triazole ring makes dihedral angles of 40.7 (3) and 30.2 (4) $^\circ$  with the 4-chlorophenyl and 2,4-difluorophenyl rings, respectively. The cation adopts a *Z*-configuration about the  $\text{C}=\text{C}$  double bond which links the triazole ring to the 4-chlorophenoxy unit *via* a butyloxy chain. In the crystal, the cations and the anions are linked by  $\text{N}-\text{H}\cdots\text{O}$ ,  $\text{C}-\text{H}\cdots\text{O}$  and  $\text{C}-\text{H}\cdots\text{F}$  hydrogen bonding.

### Related literature

For the antifungal activity of related compounds, see: Jeu *et al.* (2003). For details of the synthesis, see: Ludwig & Kurt (1985). For a related structure, see: Kurt *et al.* (1987).



### Experimental

#### Crystal data

$\text{C}_{21}\text{H}_{21}\text{ClF}_2\text{N}_3\text{O}_2^+\cdot\text{NO}_3^-$	$V = 2293.6(8)\text{ \AA}^3$
$M_r = 482.87$	$Z = 4$
Monoclinic, $P2_1/c$	Mo $K\alpha$ radiation
$a = 7.9580(16)\text{ \AA}$	$\mu = 0.22\text{ mm}^{-1}$
$b = 31.635(6)\text{ \AA}$	$T = 293\text{ K}$
$c = 9.1850(18)\text{ \AA}$	$0.30 \times 0.10 \times 0.10\text{ mm}$
$\beta = 97.29(3)^\circ$	

#### Data collection

Enraf–Nonius CAD-4	4216 independent reflections
diffractometer	1952 reflections with $I > 2\sigma(I)$
Absorption correction: $\psi$ scan	$R_{\text{int}} = 0.092$
(North <i>et al.</i> , 1968)	3 standard reflections every 200
$T_{\min} = 0.936$ , $T_{\max} = 0.978$	reflections
4521 measured reflections	intensity decay: 1%

#### Refinement

$R[F^2 > 2\sigma(F^2)] = 0.066$	298 parameters
$wR(F^2) = 0.140$	H-atom parameters constrained
$S = 1.00$	$\Delta\rho_{\max} = 0.20\text{ e \AA}^{-3}$
4216 reflections	$\Delta\rho_{\min} = -0.24\text{ e \AA}^{-3}$

**Table 1**  
Hydrogen-bond geometry ( $\text{\AA}$ ,  $^\circ$ ).

$D-\text{H}\cdots A$	$D-\text{H}$	$\text{H}\cdots A$	$D\cdots A$	$D-\text{H}\cdots A$
N3—H3A $\cdots$ O5	0.86	1.80	2.659 (4)	174
C9—H9A $\cdots$ F1 <sup>i</sup>	0.96	2.53	3.371 (5)	146
C10—H10A $\cdots$ O4 <sup>ii</sup>	0.93	2.28	3.033 (5)	137
C1—H1A $\cdots$ O3 <sup>iii</sup>	0.93	2.57	3.434 (5)	155
C11—H11A $\cdots$ O5 <sup>iii</sup>	0.93	2.25	3.180 (5)	174

Symmetry codes: (i)  $-x + 1, -y + 1, -z + 2$ ; (ii)  $x, -y + \frac{3}{2}, z - \frac{1}{2}$ ; (iii)  $x, -y + \frac{3}{2}, z + \frac{1}{2}$ .

Data collection: *CAD-4 EXPRESS* (Enraf–Nonius, 1994); cell refinement: *CAD-4 EXPRESS*; data reduction: *XCAD4* (Harms & Wocadlo, 1995); program(s) used to solve structure: *SHELXS97* (Sheldrick, 2008); program(s) used to refine structure: *SHELXL97* (Sheldrick, 2008); molecular graphics: *SHELXTL* (Sheldrick, 2008); software used to prepare material for publication: *PLATON* (Spek, 2009).

Supplementary data and figures for this paper are available from the IUCr electronic archives (Reference: PV2445).

### References

- Enraf–Nonius (1994). *CAD-4 EXPRESS*. Enraf–Nonius, Delft, The Netherlands.
- Harms, K. & Wocadlo, S. (1995). *XCAD4*. University of Marburg, Germany.
- Jeu, L., Piacenti, F. J., Lyakhovetskiy, A. G. & Fung, H. B. (2003). *Clin. Ther.* **25**, 1321–1381.
- Kurt, T., Ludwig, Z., Max, H. P., Martin, E. & Max, D. (1987). *Helv. Chim. Acta* **70**, 441–444.
- Ludwig, Z. & Kurt, T. (1985). US Patent Appl. US4554356.
- North, A. C. T., Phillips, D. C. & Mathews, F. S. (1968). *Acta Cryst. A* **24**, 351–359.
- Sheldrick, G. M. (2008). *Acta Cryst. A* **64**, 112–122.
- Spek, A. L. (2009). *Acta Cryst. D* **65**, 148–155.

## **supplementary materials**

*Acta Cryst.* (2011). E67, o2800 [doi:10.1107/S1600536811038761]

### **1-[(Z)-8-(4-Chlorophenoxy)-3-(2,4-difluorophenyl)-4-oxaocta-2-en-2-yl]-1H-1,2,4-triazol-4-i um nitrate**

**J. Chen, K. Wang, Y. Shen, L. Ren and G. Chen**

#### **Comment**

Triazole derivatives such as voriconazole ((2*R*,3*S*)-2-(2,4-difluorophenyl)-3-(5-fluoropyrimidin-4-yl)-1-(1*H*-1,2,4-triazol-1-yl) butan-2-ol) and posaconazole (4-(4-(4-(4-(((3*R*,5*R*)-5-(2,4-difluorophenyl)-5-(1,2,4-triazol-1-ylmethyl)oxolan-3-yl)methoxy)phenyl)piperazin-1-yl)phenyl)-2-((2*S*,3*S*)-2-hydroxypentan-3-yl)-1,2,4-triazol-3-one) are safe and effective antifungal agents. (Jeu *et al.*, 2003) As part of our studies on the synthesis of new triazole derivatives, the crystal structure of the title compound, having similar structure with omoconazol (Kurt *et al.*, 1987), was determined.

In the title compound (Fig. 1), the cation adopts a Z conformation about the C=C double bond. In the crystal structure the anions and cations are connected via N—H···O, C—H···O and C—H···F hydrogen bonding (Table 1 and Fig. 2).

#### **Experimental**

1-(2,4-Difluorophenyl)-2-(1,2,4-triazol-1-yl)propan-1-one (3.0 g, 0.01 mol) 10 g of a 50% aqueous sodium hydroxide, toluene (15 ml) and 1.5 ml of a 40% aqueous solution of tetrabutyl ammonium hydroxide were mixed and heated to 323 K under vigorous stirring. 1-Bromo-4-(4-chlorophenoxy)-butane (2.6 g, 0.01 mol) dissolved in 10 ml toluene, was instilled into the stirred and warmed solution in the course of 10 h. The mixture was subsequently stirred for another 20 h at 323 K. The reaction mixture was mixed with as much water and chloroform so that the aqueous phase becomes lighter than the organic phase. Thereafter, the organic and aqueous phases were separated. The organic phase was dried with sodium sulfate. The solvents were distilled under reduced pressure. The remaining residue was a dark oil that was diluted with 10 ml 2-propanol and then adjusted to a pH-value of 2 by 30% aqueous nitric acid. The resulting nitric acid solution was then cooled in the refrigerator. The impure precipitated product herein was subsequently crystallized from a 1:1 mixture of ethyl acetate and ethanol. The purified product was analytically identified as an approximately pure *Z*-isomer of the title compound. Crystals of the title compound suitable for X-ray diffraction were obtained by slow evaporation of an ethanol solution. Details on the synthesis can be found in the literature (Ludwig & Kurt, 1985).

#### **Refinement**

H atoms were positioned geometrically with C—H = 0.93 and 0.97 Å for aromatic and methylene H atoms, respectively, and with N—H = 0.86 Å for triazole H atom, and constrained to ride on their parent atoms, with  $U_{\text{iso}}(\text{H}) = 1.2$  (or 1.5 for methyl groups) times  $U_{\text{eq}}(\text{C})$ .

# supplementary materials

---

## Figures




Fig. 1. The molecular structure of the title compound, with the atom numbering scheme. Displacement ellipsoids are drawn at 30% probability levels.




Fig. 2. The packing diagram of the title compound. Hydrogen bonds are shown as dashed lines.

## 1-[*(Z*)-8-(4-Chlorophenoxy)-3-(2,4-difluorophenyl)-4-oxaocta- 2-en-2-yl]-1*H*-1,2,4-triazol-4-i um nitrate

### Crystal data

$C_{21}H_{21}ClF_2N_3O_2^+ \cdot NO_3^-$	$F(000) = 1000$
$M_r = 482.87$	$D_x = 1.398 \text{ Mg m}^{-3}$
Monoclinic, $P2_1/c$	Mo $K\alpha$ radiation, $\lambda = 0.71073 \text{ \AA}$
Hall symbol: -P 2ybc	Cell parameters from 25 reflections
$a = 7.9580 (16) \text{ \AA}$	$\theta = 9-12^\circ$
$b = 31.635 (6) \text{ \AA}$	$\mu = 0.22 \text{ mm}^{-1}$
$c = 9.1850 (18) \text{ \AA}$	$T = 293 \text{ K}$
$\beta = 97.29 (3)^\circ$	Block, yellow
$V = 2293.6 (8) \text{ \AA}^3$	$0.30 \times 0.10 \times 0.10 \text{ mm}$
$Z = 4$	

### Data collection

Enraf–Nonius CAD-4 diffractometer	1952 reflections with $I > 2\sigma(I)$
Radiation source: fine-focus sealed tube	$R_{\text{int}} = 0.092$
graphite	$\theta_{\max} = 25.4^\circ, \theta_{\min} = 1.3^\circ$
$\omega/2\theta$ scans	$h = 0 \rightarrow 9$
Absorption correction: $\psi$ scan (North <i>et al.</i> , 1968)	$k = 0 \rightarrow 38$
$T_{\min} = 0.936, T_{\max} = 0.978$	$l = -11 \rightarrow 10$
4521 measured reflections	3 standard reflections every 200 reflections
4216 independent reflections	intensity decay: 1%

### Refinement

Refinement on $F^2$	Primary atom site location: structure-invariant direct methods
---------------------	--

Least-squares matrix: full	Secondary atom site location: difference Fourier map
$R[F^2 > 2\sigma(F^2)] = 0.066$	Hydrogen site location: inferred from neighbouring sites
$wR(F^2) = 0.140$	H-atom parameters constrained
$S = 1.00$	$w = 1/[\sigma^2(F_o^2) + (0.048P)^2]$ where $P = (F_o^2 + 2F_c^2)/3$
4216 reflections	$(\Delta/\sigma)_{\max} < 0.001$
298 parameters	$\Delta\rho_{\max} = 0.20 \text{ e \AA}^{-3}$
0 restraints	$\Delta\rho_{\min} = -0.24 \text{ e \AA}^{-3}$

### Special details

**Geometry.** All e.s.d.'s (except the e.s.d. in the dihedral angle between two l.s. planes) are estimated using the full covariance matrix. The cell e.s.d.'s are taken into account individually in the estimation of e.s.d.'s in distances, angles and torsion angles; correlations between e.s.d.'s in cell parameters are only used when they are defined by crystal symmetry. An approximate (isotropic) treatment of cell e.s.d.'s is used for estimating e.s.d.'s involving l.s. planes.

**Refinement.** Refinement of  $F^2$  against ALL reflections. The weighted  $R$ -factor  $wR$  and goodness of fit  $S$  are based on  $F^2$ , conventional  $R$ -factors  $R$  are based on  $F$ , with  $F$  set to zero for negative  $F^2$ . The threshold expression of  $F^2 > \sigma(F^2)$  is used only for calculating  $R$ -factors(gt) etc. and is not relevant to the choice of reflections for refinement.  $R$ -factors based on  $F^2$  are statistically about twice as large as those based on  $F$ , and  $R$ -factors based on ALL data will be even larger.

### Fractional atomic coordinates and isotropic or equivalent isotropic displacement parameters ( $\text{\AA}^2$ )

	$x$	$y$	$z$	$U_{\text{iso}}^*/U_{\text{eq}}$
Cl	1.06795 (17)	0.98117 (4)	0.78129 (14)	0.0884 (4)
O1	0.5208 (3)	0.66925 (8)	0.7812 (2)	0.0537 (7)
N1	0.4023 (4)	0.64572 (10)	0.5034 (3)	0.0492 (8)
F1	0.6286 (3)	0.51310 (8)	1.2146 (3)	0.0929 (9)
C1	0.4331 (5)	0.59449 (12)	0.9814 (4)	0.0600 (11)
H1A	0.3389	0.6121	0.9757	0.072*
O2	0.8631 (4)	0.80660 (8)	0.8732 (3)	0.0665 (8)
F2	0.7867 (3)	0.57220 (8)	0.7858 (3)	0.0912 (9)
N2	0.4433 (5)	0.63841 (11)	0.3638 (3)	0.0667 (10)
C2	0.4625 (6)	0.56669 (14)	1.0964 (4)	0.0680 (13)
H2B	0.3904	0.5656	1.1684	0.082*
N3	0.3111 (4)	0.69903 (10)	0.3756 (3)	0.0534 (9)
H3A	0.2657	0.7229	0.3485	0.064*
C3	0.5993 (7)	0.54084 (13)	1.1027 (4)	0.0622 (12)
C4	0.7106 (6)	0.54183 (13)	0.9996 (4)	0.0668 (12)
H4A	0.8045	0.5241	1.0054	0.080*
C5	0.6748 (6)	0.57053 (13)	0.8877 (4)	0.0569 (11)
C6	0.5399 (5)	0.59716 (12)	0.8733 (4)	0.0463 (10)
C7	0.5057 (5)	0.62726 (12)	0.7499 (4)	0.0444 (9)
C8	0.4462 (5)	0.61470 (11)	0.6153 (4)	0.0498 (10)
C9	0.4189 (6)	0.57050 (13)	0.5668 (4)	0.0864 (16)
H9A	0.4507	0.5518	0.6480	0.130*
H9B	0.4867	0.5645	0.4899	0.130*

## supplementary materials

---

H9C	0.3014	0.5664	0.5307	0.130*
C10	0.3847 (6)	0.67162 (14)	0.2922 (4)	0.0652 (12)
H10A	0.3928	0.6760	0.1932	0.078*
C11	0.3222 (5)	0.68190 (12)	0.5088 (4)	0.0506 (10)
H11A	0.2811	0.6934	0.5905	0.061*
C12	0.6628 (5)	0.68174 (11)	0.8866 (4)	0.0484 (10)
H12A	0.7679	0.6721	0.8546	0.058*
H12B	0.6527	0.6691	0.9813	0.058*
C13	0.6635 (5)	0.72830 (11)	0.8991 (4)	0.0528 (10)
H13A	0.5584	0.7374	0.9325	0.063*
H13B	0.6681	0.7405	0.8028	0.063*
C14	0.8110 (5)	0.74505 (12)	1.0040 (4)	0.0503 (10)
H14A	0.8013	0.7342	1.1014	0.060*
H14B	0.9152	0.7339	0.9746	0.060*
C15	0.8246 (5)	0.79196 (12)	1.0125 (4)	0.0542 (10)
H15A	0.7186	0.8041	1.0342	0.065*
H15B	0.9135	0.8002	1.0895	0.065*
C16	0.9051 (5)	0.84798 (12)	0.8603 (4)	0.0526 (10)
C17	0.9642 (5)	0.85865 (13)	0.7305 (4)	0.0591 (11)
H17A	0.9710	0.8380	0.6593	0.071*
C18	1.0133 (5)	0.89934 (14)	0.7049 (4)	0.0617 (11)
H18A	1.0538	0.9062	0.6172	0.074*
C19	1.0020 (5)	0.93006 (13)	0.8108 (5)	0.0617 (11)
C20	0.9427 (6)	0.91986 (14)	0.9391 (5)	0.0724 (13)
H20A	0.9355	0.9405	1.0100	0.087*
C21	0.8929 (5)	0.87874 (13)	0.9644 (4)	0.0667 (12)
H21A	0.8514	0.8720	1.0517	0.080*
N4	0.1836 (4)	0.79947 (13)	0.3777 (4)	0.0604 (9)
O3	0.1244 (4)	0.83434 (11)	0.3510 (4)	0.0941 (11)
O4	0.2476 (4)	0.78801 (11)	0.5018 (3)	0.0883 (11)
O5	0.1781 (3)	0.77208 (8)	0.2747 (3)	0.0588 (7)

Atomic displacement parameters ( $\text{\AA}^2$ )

	$U^{11}$	$U^{22}$	$U^{33}$	$U^{12}$	$U^{13}$	$U^{23}$
Cl	0.1080 (10)	0.0611 (8)	0.0955 (9)	-0.0204 (7)	0.0108 (7)	0.0053 (7)
O1	0.0688 (19)	0.0470 (16)	0.0404 (14)	-0.0018 (13)	-0.0116 (13)	-0.0013 (12)
N1	0.066 (2)	0.046 (2)	0.0340 (17)	0.0003 (17)	-0.0015 (15)	0.0002 (15)
F1	0.126 (2)	0.0812 (19)	0.0663 (16)	-0.0072 (17)	-0.0075 (15)	0.0369 (14)
C1	0.075 (3)	0.054 (3)	0.052 (2)	0.003 (2)	0.011 (2)	0.010 (2)
O2	0.102 (2)	0.0489 (18)	0.0517 (17)	-0.0183 (16)	0.0205 (15)	-0.0070 (14)
F2	0.090 (2)	0.106 (2)	0.0833 (18)	0.0335 (17)	0.0354 (16)	0.0291 (15)
N2	0.095 (3)	0.072 (3)	0.0321 (18)	0.014 (2)	0.0053 (18)	-0.0024 (17)
C2	0.094 (4)	0.066 (3)	0.047 (3)	-0.008 (3)	0.021 (2)	0.011 (2)
N3	0.067 (2)	0.051 (2)	0.0388 (18)	-0.0046 (18)	-0.0070 (16)	0.0064 (16)
C3	0.098 (4)	0.045 (3)	0.040 (2)	-0.009 (3)	-0.006 (2)	0.0116 (19)
C4	0.082 (3)	0.061 (3)	0.054 (3)	0.016 (2)	-0.004 (2)	0.007 (2)
C5	0.077 (3)	0.056 (3)	0.039 (2)	0.001 (2)	0.011 (2)	-0.001 (2)

C6	0.058 (3)	0.045 (2)	0.034 (2)	0.002 (2)	0.0029 (19)	0.0002 (17)
C7	0.051 (2)	0.039 (2)	0.041 (2)	-0.0049 (18)	-0.0012 (17)	-0.0023 (17)
C8	0.069 (3)	0.040 (2)	0.038 (2)	-0.001 (2)	-0.0001 (19)	0.0038 (18)
C9	0.132 (5)	0.057 (3)	0.062 (3)	-0.017 (3)	-0.022 (3)	-0.003 (2)
C10	0.093 (4)	0.066 (3)	0.035 (2)	0.001 (3)	0.000 (2)	0.001 (2)
C11	0.059 (3)	0.054 (3)	0.036 (2)	-0.007 (2)	-0.0015 (18)	0.0002 (19)
C12	0.049 (2)	0.051 (2)	0.043 (2)	-0.002 (2)	-0.0027 (18)	-0.0036 (18)
C13	0.063 (3)	0.047 (3)	0.047 (2)	-0.006 (2)	-0.0001 (19)	-0.0017 (19)
C14	0.045 (2)	0.057 (3)	0.048 (2)	-0.001 (2)	0.0024 (19)	-0.0049 (19)
C15	0.057 (3)	0.056 (3)	0.048 (2)	-0.003 (2)	-0.0002 (19)	-0.008 (2)
C16	0.057 (3)	0.050 (3)	0.051 (2)	-0.008 (2)	0.010 (2)	0.000 (2)
C17	0.067 (3)	0.054 (3)	0.056 (3)	-0.002 (2)	0.005 (2)	-0.008 (2)
C18	0.063 (3)	0.066 (3)	0.057 (3)	-0.004 (2)	0.010 (2)	0.003 (2)
C19	0.069 (3)	0.050 (3)	0.066 (3)	-0.011 (2)	0.006 (2)	0.001 (2)
C20	0.093 (4)	0.052 (3)	0.075 (3)	-0.014 (3)	0.022 (3)	-0.020 (2)
C21	0.087 (3)	0.059 (3)	0.059 (3)	-0.024 (2)	0.030 (2)	-0.018 (2)
N4	0.055 (2)	0.075 (3)	0.053 (2)	-0.007 (2)	0.0119 (18)	-0.002 (2)
O3	0.089 (3)	0.062 (2)	0.132 (3)	0.011 (2)	0.014 (2)	-0.007 (2)
O4	0.097 (2)	0.130 (3)	0.0366 (17)	-0.010 (2)	0.0009 (16)	-0.0060 (18)
O5	0.077 (2)	0.0595 (18)	0.0377 (14)	-0.0015 (15)	0.0004 (13)	-0.0056 (14)

*Geometric parameters (Å, °)*

Cl—C19	1.732 (4)	C9—H9C	0.9600
O1—C7	1.361 (4)	C10—H10A	0.9300
O1—C12	1.447 (4)	C11—H11A	0.9300
N1—C11	1.314 (4)	C12—C13	1.477 (4)
N1—N2	1.383 (4)	C12—H12A	0.9700
N1—C8	1.432 (4)	C12—H12B	0.9700
F1—C3	1.349 (4)	C13—C14	1.517 (5)
C1—C2	1.371 (5)	C13—H13A	0.9700
C1—C6	1.389 (5)	C13—H13B	0.9700
C1—H1A	0.9300	C14—C15	1.489 (5)
O2—C16	1.360 (4)	C14—H14A	0.9700
O2—C15	1.430 (4)	C14—H14B	0.9700
F2—C5	1.372 (4)	C15—H15A	0.9700
N2—C10	1.295 (5)	C15—H15B	0.9700
C2—C3	1.357 (6)	C16—C21	1.376 (5)
C2—H2B	0.9300	C16—C17	1.378 (5)
N3—C11	1.330 (4)	C17—C18	1.374 (5)
N3—C10	1.340 (5)	C17—H17A	0.9300
N3—H3A	0.8600	C18—C19	1.386 (5)
C3—C4	1.377 (6)	C18—H18A	0.9300
C4—C5	1.374 (5)	C19—C20	1.362 (5)
C4—H4A	0.9300	C20—C21	1.388 (5)
C5—C6	1.358 (5)	C20—H20A	0.9300
C6—C7	1.479 (5)	C21—H21A	0.9300
C7—C8	1.327 (5)	N4—O3	1.212 (4)
C8—C9	1.475 (5)	N4—O4	1.243 (4)

## supplementary materials

---

C9—H9A	0.9600	N4—O5	1.279 (4)
C9—H9B	0.9600		
C7—O1—C12	116.6 (3)	O1—C12—C13	108.6 (3)
C11—N1—N2	110.7 (3)	O1—C12—H12A	110.0
C11—N1—C8	130.2 (3)	C13—C12—H12A	110.0
N2—N1—C8	119.1 (3)	O1—C12—H12B	110.0
C2—C1—C6	122.0 (4)	C13—C12—H12B	110.0
C2—C1—H1A	119.0	H12A—C12—H12B	108.3
C6—C1—H1A	119.0	C12—C13—C14	113.1 (3)
C16—O2—C15	118.2 (3)	C12—C13—H13A	109.0
C10—N2—N1	102.9 (3)	C14—C13—H13A	109.0
C3—C2—C1	118.5 (4)	C12—C13—H13B	109.0
C3—C2—H2B	120.8	C14—C13—H13B	109.0
C1—C2—H2B	120.8	H13A—C13—H13B	107.8
C11—N3—C10	106.2 (3)	C15—C14—C13	115.2 (3)
C11—N3—H3A	126.9	C15—C14—H14A	108.5
C10—N3—H3A	126.9	C13—C14—H14A	108.5
F1—C3—C2	119.0 (4)	C15—C14—H14B	108.5
F1—C3—C4	118.3 (4)	C13—C14—H14B	108.5
C2—C3—C4	122.7 (4)	H14A—C14—H14B	107.5
C5—C4—C3	116.0 (4)	O2—C15—C14	107.3 (3)
C5—C4—H4A	122.0	O2—C15—H15A	110.2
C3—C4—H4A	122.0	C14—C15—H15A	110.2
C6—C5—F2	118.9 (3)	O2—C15—H15B	110.2
C6—C5—C4	124.7 (4)	C14—C15—H15B	110.2
F2—C5—C4	116.4 (4)	H15A—C15—H15B	108.5
C5—C6—C1	116.1 (4)	O2—C16—C21	125.5 (4)
C5—C6—C7	122.7 (3)	O2—C16—C17	115.2 (3)
C1—C6—C7	121.2 (4)	C21—C16—C17	119.3 (4)
C8—C7—O1	120.0 (3)	C18—C17—C16	120.9 (4)
C8—C7—C6	122.0 (3)	C18—C17—H17A	119.5
O1—C7—C6	117.6 (3)	C16—C17—H17A	119.5
C7—C8—N1	119.3 (3)	C17—C18—C19	119.4 (4)
C7—C8—C9	125.9 (3)	C17—C18—H18A	120.3
N1—C8—C9	114.8 (3)	C19—C18—H18A	120.3
C8—C9—H9A	109.5	C20—C19—C18	120.1 (4)
C8—C9—H9B	109.5	C20—C19—Cl	120.1 (3)
H9A—C9—H9B	109.5	C18—C19—Cl	119.8 (3)
C8—C9—H9C	109.5	C19—C20—C21	120.3 (4)
H9A—C9—H9C	109.5	C19—C20—H20A	119.9
H9B—C9—H9C	109.5	C21—C20—H20A	119.9
N2—C10—N3	113.0 (4)	C16—C21—C20	120.0 (4)
N2—C10—H10A	123.5	C16—C21—H21A	120.0
N3—C10—H10A	123.5	C20—C21—H21A	120.0
N1—C11—N3	107.2 (3)	O3—N4—O4	123.7 (4)
N1—C11—H11A	126.4	O3—N4—O5	119.4 (4)
N3—C11—H11A	126.4	O4—N4—O5	116.9 (4)
C11—N1—N2—C10	0.3 (4)	N2—N1—C8—C7	139.3 (4)

C8—N1—N2—C10	179.2 (3)	C11—N1—C8—C9	137.7 (4)
C6—C1—C2—C3	-0.8 (6)	N2—N1—C8—C9	-40.9 (5)
C1—C2—C3—F1	-179.4 (4)	N1—N2—C10—N3	0.2 (5)
C1—C2—C3—C4	0.9 (7)	C11—N3—C10—N2	-0.6 (5)
F1—C3—C4—C5	179.8 (3)	N2—N1—C11—N3	-0.7 (4)
C2—C3—C4—C5	-0.5 (6)	C8—N1—C11—N3	-179.4 (3)
C3—C4—C5—C6	0.0 (6)	C10—N3—C11—N1	0.8 (4)
C3—C4—C5—F2	179.0 (3)	C7—O1—C12—C13	176.6 (3)
F2—C5—C6—C1	-178.8 (3)	O1—C12—C13—C14	-178.3 (3)
C4—C5—C6—C1	0.2 (6)	C12—C13—C14—C15	175.8 (3)
F2—C5—C6—C7	1.6 (6)	C16—O2—C15—C14	-171.0 (3)
C4—C5—C6—C7	-179.4 (4)	C13—C14—C15—O2	-66.4 (4)
C2—C1—C6—C5	0.3 (6)	C15—O2—C16—C21	-8.9 (6)
C2—C1—C6—C7	179.8 (4)	C15—O2—C16—C17	171.0 (3)
C12—O1—C7—C8	-145.4 (4)	O2—C16—C17—C18	-179.0 (4)
C12—O1—C7—C6	41.0 (4)	C21—C16—C17—C18	1.0 (6)
C5—C6—C7—C8	74.0 (5)	C16—C17—C18—C19	-0.5 (6)
C1—C6—C7—C8	-105.6 (5)	C17—C18—C19—C20	0.1 (6)
C5—C6—C7—O1	-112.6 (4)	C17—C18—C19—Cl	178.6 (3)
C1—C6—C7—O1	67.8 (5)	C18—C19—C20—C21	-0.2 (7)
O1—C7—C8—N1	1.5 (6)	Cl—C19—C20—C21	-178.7 (4)
C6—C7—C8—N1	174.8 (3)	O2—C16—C21—C20	178.9 (4)
O1—C7—C8—C9	-178.3 (4)	C17—C16—C21—C20	-1.1 (6)
C6—C7—C8—C9	-5.0 (7)	C19—C20—C21—C16	0.7 (7)
C11—N1—C8—C7	-42.1 (6)		

*Hydrogen-bond geometry (Å, °)*

<i>D</i> —H··· <i>A</i>	<i>D</i> —H	H··· <i>A</i>	<i>D</i> ··· <i>A</i>	<i>D</i> —H··· <i>A</i>
N3—H3A···O5	0.86	1.80	2.659 (4)	174
C9—H9A···F1 <sup>i</sup>	0.96	2.53	3.371 (5)	146
C10—H10A···O4 <sup>ii</sup>	0.93	2.28	3.033 (5)	137
C1—H1A···O3 <sup>iii</sup>	0.93	2.57	3.434 (5)	155
C11—H11A···O5 <sup>iii</sup>	0.93	2.25	3.180 (5)	174

Symmetry codes: (i)  $-x+1, -y+1, -z+2$ ; (ii)  $x, -y+3/2, z-1/2$ ; (iii)  $x, -y+3/2, z+1/2$ .

## supplementary materials

---

Fig. 1




Fig. 2

