

Hexaaquacobalt(II) bis[4-(pyridin-2-yl-methoxy)benzoate] dihydrate

Li-Wei Zhang,^a Shan Gao^a and Seik Weng Ng^{b,c,*}

^aKey Laboratory of Functional Inorganic Material Chemistry, Ministry of Education, Heilongjiang University, Harbin 150080, People's Republic of China, ^bDepartment of Chemistry, University of Malaya, 50603 Kuala Lumpur, Malaysia, and ^cChemistry Department, Faculty of Science, King Abdulaziz University, PO Box 80203 Jeddah, Saudi Arabia

Correspondence e-mail: seikweng@um.edu.my


Received 3 October 2011; accepted 5 October 2011

Key indicators: single-crystal X-ray study; $T = 293$ K; mean $\sigma(\text{C}-\text{C}) = 0.004$ Å; R factor = 0.036; wR factor = 0.138; data-to-parameter ratio = 14.5.

The Co^{II} atom in the title salt, $[\text{Co}(\text{H}_2\text{O})_6](\text{C}_{13}\text{H}_{10}\text{NO}_3)_2 \cdot 2\text{H}_2\text{O}$, lies on a center of inversion in an octahedron of water molecules. The cations, anions and uncoordinated water molecules are linked by $\text{O}-\text{H} \cdots \text{O}$ and $\text{O}-\text{H} \cdots \text{N}$ hydrogen bonds into a three-dimensional network. The anion is essentially planar, with an r.m.s. deviation of all non-H atoms of 0.066 Å.

Related literature

There are many examples of hexaaquacobalt(II) benzoates; these benzoates possess substituents capable of serving as hydrogen-bond acceptors/donors, see: Deng *et al.* (2006). For the isotypic Mn(II) salt, see: Zhang *et al.* (2011).



Experimental

Crystal data

 $[\text{Co}(\text{H}_2\text{O})_6](\text{C}_{13}\text{H}_{10}\text{NO}_3)_2 \cdot 2\text{H}_2\text{O}$ $M_r = 659.50$ Triclinic, $P\bar{1}$ $a = 7.4349$ (3) Å $b = 7.5431$ (3) Å $c = 13.7531$ (6) Å $\alpha = 84.307$ (1)° $\beta = 83.153$ (1)° $\gamma = 73.333$ (1)° $V = 731.91$ (5) Å³ $Z = 1$ Mo $K\alpha$ radiation $\mu = 0.66$ mm⁻¹ $T = 293$ K

0.21 × 0.15 × 0.13 mm

Data collection

Rigaku R-AXIS RAPID IP diffractometer
Absorption correction: multi-scan (*ABSCOR*; Higashi, 1995)
 $T_{\text{min}} = 0.874$, $T_{\text{max}} = 0.919$

7251 measured reflections
3314 independent reflections
2401 reflections with $I > 2\sigma(I)$
 $R_{\text{int}} = 0.026$

Refinement

 $R[F^2 > 2\sigma(F^2)] = 0.036$ $wR(F^2) = 0.138$ $S = 1.15$

3314 reflections

228 parameters

18 restraints

H atoms treated by a mixture of independent and constrained refinement

 $\Delta\rho_{\text{max}} = 0.63$ e Å⁻³ $\Delta\rho_{\text{min}} = -0.69$ e Å⁻³

Table 1

Hydrogen-bond geometry (Å, °).

$D-\text{H} \cdots A$	$D-\text{H}$	$\text{H} \cdots A$	$D \cdots A$	$D-\text{H} \cdots A$
$\text{O1w}-\text{H11} \cdots \text{O1}$	0.84 (1)	1.94 (1)	2.762 (3)	169 (3)
$\text{O1w}-\text{H12} \cdots \text{O4w}^{\text{i}}$	0.84 (1)	1.87 (1)	2.696 (4)	171 (4)
$\text{O2w}-\text{H21} \cdots \text{O2}$	0.84 (1)	1.84 (1)	2.678 (3)	179 (3)
$\text{O2w}-\text{H22} \cdots \text{O2}^{\text{ii}}$	0.84 (1)	1.97 (2)	2.755 (3)	156 (3)
$\text{O3w}-\text{H31} \cdots \text{O1}^{\text{iii}}$	0.85 (1)	1.96 (1)	2.798 (3)	170 (4)
$\text{O3w}-\text{H32} \cdots \text{N1}^{\text{iv}}$	0.85 (1)	1.95 (1)	2.799 (3)	177 (4)
$\text{O4w}-\text{H41} \cdots \text{O2}$	0.84 (1)	2.22 (4)	2.891 (4)	137 (5)
$\text{O4w}-\text{H42} \cdots \text{O3w}^{\text{v}}$	0.84 (1)	2.42 (4)	3.176 (4)	151 (7)

Symmetry codes: (i) $-x, -y + 2, -z + 1$; (ii) $-x + 1, -y + 2, -z + 1$; (iii) $-x + 1, -y + 1, -z + 1$; (iv) $x, y - 1, z + 1$; (v) $x, y + 1, z$.

Data collection: *RAPID-AUTO* (Rigaku, 1998); cell refinement: *RAPID-AUTO*; data reduction: *CrystalClear* (Rigaku/MSK, 2002); program(s) used to solve structure: *SHELXS97* (Sheldrick, 2008); program(s) used to refine structure: *SHELXL97* (Sheldrick, 2008); molecular graphics: *X-SEED* (Barbour, 2001); software used to prepare material for publication: *publCIF* (Westrip, 2010).

This work was supported by the Key Project of the Natural Science Foundation of Heilongjiang Province (No. ZD200903), the Innovation Team of the Education Bureau of Heilongjiang Province (No. 2010 t d03), the Key Project of the Education Bureau of Heilongjiang Province (No. 12511z023) and the University of Malaya.

Supplementary data and figures for this paper are available from the IUCr electronic archives (Reference: BT5664).

References

- Barbour, L. J. (2001). *J. Supramol. Chem.* **1**, 189–191.
Deng, Z.-P., Gao, S. & Ng, S. W. (2006). *Acta Cryst.* **E62**, m3423–m3424.
Higashi, T. (1995). *ABSCOR*. Rigaku Corporation, Tokyo, Japan.
Rigaku (1998). *RAPID-AUTO*. Rigaku Corporation, Tokyo, Japan.
Rigaku/MSK (2002). *CrystalClear*. Rigaku/MSK Inc., The Woodlands, Texas, USA.
Sheldrick, G. M. (2008). *Acta Cryst.* **A64**, 112–122.
Westrip, S. P. (2010). *J. Appl. Cryst.* **43**, 920–925.
Zhang, L.-W., Gao, S. & Ng, S. W. (2011). *Acta Cryst.* **E67**, m1520.

supplementary materials

Acta Cryst. (2011). E67, m1519 [doi:10.1107/S1600536811040931]

Hexaaquacobalt(II) bis[4-(pyridin-2-ylmethoxy)benzoate] dihydrate

L.-W. Zhang, S. Gao and S. W. Ng

Comment

First-row transition metal dications form a plethora of metal dicarboxylates; however, occasionally, no direct metal–carboxylate bond is formed, and the product consists of hexaaquametal cations and carboxylate ions, the anion interacting indirectly in an outer-sphere type of coordination. 4-(Pyridin-2-ylmethoxy)benzoic acid is a commercially available carboxylic acid but there are no reports on its metal carboxylates. The reaction of the deprotonated acid with cobalt(II) ions gives the hexaaquacobalt(II) salt (Scheme I, Fig. 1). The Co^{II} atom in the salt lies on a center-of-inversion in an octahedron of water molecules. The metal atom interacts with the carboxylate ion indirectly, through the coordinated water molecules, in an outer-sphere type of coordination. The cations, anions and lattice water molecules are linked by $\text{O}\cdots\text{H}\cdots\text{O}$ and $\text{O}-\text{H}\cdots\text{N}$ hydrogen bonds into a three-dimensional network (Table 1).

Experimental

Cobalt nitrate (1 mmol) was added to an aqueous solution of 4-(pyridin-2-ylmethoxy)benzoic acid (2 mmol) that was earlier been treated with 1M sodium hydroxide to a pH of 6. The filtered solution was set aside for several days, after which pink prismatic crystals separated from solution.

Refinement

Carbon-bound H-atoms were placed in calculated positions (C–H 0.93 Å) and were included in the refinement in the riding model approximation, with $U(\text{H})$ set to $1.2U(\text{C})$. The water H-atoms were located in a difference Fourier map, and were refined with distance restraints of O–H 0.84 ± 0.01 Å and H \cdots H 1.37 ± 0.01 Å; their displacement factors were refined.

The anisotropic displacement ellipsoids of the lattice water O were restrained to an isotropic behaviour.

The (2 2 7) reflection was omitted owing to bad agreement.

Figures




Fig. 1. Anisotropic displacement ellipsoid plot (Barbour, 2001) of $\text{Co}(\text{H}_2\text{O})_6 \cdot 2(\text{C}_{13}\text{H}_{10}\text{NO}_3) \cdot 2\text{H}_2\text{O}$ at the 50% probability level; hydrogen atoms are drawn as spheres of arbitrary radius.

Hexaaquacobalt(II) bis[4-(pyridin-2-ylmethoxy)benzoate] dihydrate

Crystal data

$[\text{Co}(\text{H}_2\text{O})_6](\text{C}_{13}\text{H}_{10}\text{NO}_3)_2 \cdot 2\text{H}_2\text{O}$	$Z = 1$
$M_r = 659.50$	$F(000) = 345$
Triclinic, PT	$D_x = 1.496 \text{ Mg m}^{-3}$
Hall symbol: $-P 1$	Mo $K\alpha$ radiation, $\lambda = 0.71073 \text{ \AA}$
$a = 7.4349 (3) \text{ \AA}$	Cell parameters from 5547 reflections
$b = 7.5431 (3) \text{ \AA}$	$\theta = 3.1\text{--}27.4^\circ$
$c = 13.7531 (6) \text{ \AA}$	$\mu = 0.66 \text{ mm}^{-1}$
$\alpha = 84.307 (1)^\circ$	$T = 293 \text{ K}$
$\beta = 83.153 (1)^\circ$	Prism, pink
$\gamma = 73.333 (1)^\circ$	$0.21 \times 0.15 \times 0.13 \text{ mm}$
$V = 731.91 (5) \text{ \AA}^3$	

Data collection

Rigaku R-Axis RAPID IP diffractometer	3314 independent reflections
Radiation source: fine-focus sealed tube graphite	2401 reflections with $I > 2\sigma(I)$
ω scans	$R_{\text{int}} = 0.026$
Absorption correction: multi-scan (<i>ABSCOR</i> ; Higashi, 1995)	$\theta_{\text{max}} = 27.4^\circ$, $\theta_{\text{min}} = 3.1^\circ$
$T_{\text{min}} = 0.874$, $T_{\text{max}} = 0.919$	$h = -9 \rightarrow 9$
7251 measured reflections	$k = -9 \rightarrow 9$
	$l = -17 \rightarrow 17$

Refinement

Refinement on F^2	Primary atom site location: structure-invariant direct methods
Least-squares matrix: full	Secondary atom site location: difference Fourier map
$R[F^2 > 2\sigma(F^2)] = 0.036$	Hydrogen site location: inferred from neighbouring sites
$wR(F^2) = 0.138$	H atoms treated by a mixture of independent and constrained refinement
$S = 1.15$	$w = 1/[\sigma^2(F_o^2) + (0.0658P)^2 + 0.3695P]$
3314 reflections	where $P = (F_o^2 + 2F_c^2)/3$
228 parameters	$(\Delta/\sigma)_{\text{max}} = 0.001$
18 restraints	$\Delta\rho_{\text{max}} = 0.63 \text{ e \AA}^{-3}$
	$\Delta\rho_{\text{min}} = -0.69 \text{ e \AA}^{-3}$

Fractional atomic coordinates and isotropic or equivalent isotropic displacement parameters (\AA^2)

	x	y	z	$U_{\text{iso}}^*/U_{\text{eq}}$
Co1	0.5000	0.5000	0.5000	0.02876 (18)

O1	0.4152 (3)	0.8175 (3)	0.26860 (15)	0.0429 (5)
O2	0.3638 (3)	1.0264 (3)	0.37899 (14)	0.0415 (5)
O3	0.2477 (4)	1.5730 (3)	-0.00100 (15)	0.0493 (6)
O1W	0.2738 (3)	0.5989 (3)	0.41333 (15)	0.0368 (5)
O2W	0.4601 (3)	0.7721 (2)	0.52791 (14)	0.0368 (5)
O3W	0.3196 (3)	0.4485 (3)	0.62650 (14)	0.0376 (5)
O4W	0.0843 (4)	1.2664 (5)	0.5083 (3)	0.0673 (8)
N1	0.1751 (4)	1.7407 (3)	-0.25030 (18)	0.0403 (6)
C1	0.3773 (4)	0.9820 (4)	0.2913 (2)	0.0321 (6)
C2	0.3421 (4)	1.1356 (4)	0.21207 (19)	0.0301 (6)
C3	0.3111 (5)	1.3191 (4)	0.2343 (2)	0.0373 (7)
H3	0.3107	1.3462	0.2989	0.045*
C4	0.2809 (5)	1.4610 (4)	0.1611 (2)	0.0403 (7)
H4	0.2604	1.5830	0.1764	0.048*
C5	0.2812 (4)	1.4210 (4)	0.0650 (2)	0.0345 (6)
C6	0.3114 (5)	1.2396 (4)	0.0414 (2)	0.0366 (7)
H6	0.3109	1.2130	-0.0232	0.044*
C7	0.3425 (4)	1.0985 (4)	0.1154 (2)	0.0338 (6)
H7	0.3641	0.9765	0.0998	0.041*
C8	0.2360 (5)	1.5490 (4)	-0.1010 (2)	0.0363 (7)
H8A	0.1397	1.4877	-0.1062	0.044*
H8B	0.3557	1.4734	-0.1294	0.044*
C9	0.1869 (4)	1.7387 (4)	-0.15412 (19)	0.0317 (6)
C10	0.1258 (5)	1.9066 (5)	-0.3007 (2)	0.0452 (8)
H10	0.1169	1.9094	-0.3678	0.054*
C11	0.0878 (5)	2.0720 (5)	-0.2581 (3)	0.0490 (8)
H11A	0.0534	2.1840	-0.2955	0.059*
C12	0.1014 (5)	2.0687 (5)	-0.1596 (3)	0.0502 (8)
H12A	0.0765	2.1787	-0.1288	0.060*
C13	0.1528 (5)	1.8992 (4)	-0.1062 (2)	0.0434 (7)
H13	0.1642	1.8936	-0.0393	0.052*
H11	0.304 (5)	0.677 (4)	0.372 (2)	0.065 (13)*
H12	0.167 (3)	0.646 (5)	0.442 (3)	0.069 (14)*
H21	0.430 (5)	0.853 (3)	0.4814 (16)	0.048 (10)*
H22	0.503 (5)	0.816 (4)	0.5708 (17)	0.053 (11)*
H31	0.390 (5)	0.361 (3)	0.660 (2)	0.065 (12)*
H32	0.271 (5)	0.536 (3)	0.664 (2)	0.065 (12)*
H41	0.117 (7)	1.227 (7)	0.4522 (18)	0.10 (2)*
H42	0.149 (10)	1.336 (11)	0.518 (5)	0.24 (5)*

Atomic displacement parameters (\AA^2)

	U^{11}	U^{22}	U^{33}	U^{12}	U^{13}	U^{23}
Co1	0.0379 (3)	0.0210 (3)	0.0273 (3)	-0.0075 (2)	-0.0068 (2)	0.00113 (19)
O1	0.0675 (15)	0.0234 (9)	0.0361 (11)	-0.0079 (10)	-0.0129 (10)	0.0013 (8)
O2	0.0683 (15)	0.0294 (10)	0.0284 (10)	-0.0143 (10)	-0.0137 (10)	0.0032 (8)
O3	0.0880 (18)	0.0294 (10)	0.0294 (11)	-0.0134 (11)	-0.0167 (11)	0.0073 (8)
O1W	0.0426 (13)	0.0308 (10)	0.0367 (11)	-0.0092 (9)	-0.0095 (10)	0.0021 (9)

supplementary materials

O2W	0.0612 (14)	0.0230 (9)	0.0277 (10)	-0.0109 (9)	-0.0136 (9)	-0.0005 (8)
O3W	0.0462 (13)	0.0312 (10)	0.0294 (10)	-0.0035 (9)	-0.0002 (9)	0.0000 (9)
O4W	0.0531 (17)	0.0683 (18)	0.077 (2)	-0.0156 (14)	0.0123 (15)	-0.0117 (16)
N1	0.0455 (15)	0.0375 (13)	0.0312 (12)	-0.0026 (11)	-0.0027 (11)	0.0020 (10)
C1	0.0416 (17)	0.0266 (13)	0.0296 (14)	-0.0098 (12)	-0.0121 (12)	0.0026 (11)
C2	0.0354 (15)	0.0248 (12)	0.0294 (13)	-0.0071 (11)	-0.0063 (11)	0.0022 (10)
C3	0.0537 (19)	0.0300 (14)	0.0274 (13)	-0.0080 (13)	-0.0094 (13)	-0.0021 (11)
C4	0.060 (2)	0.0231 (13)	0.0366 (15)	-0.0077 (13)	-0.0107 (14)	0.0000 (11)
C5	0.0428 (17)	0.0305 (14)	0.0286 (13)	-0.0084 (12)	-0.0074 (12)	0.0057 (11)
C6	0.0532 (19)	0.0309 (14)	0.0244 (13)	-0.0088 (13)	-0.0065 (12)	-0.0016 (11)
C7	0.0404 (16)	0.0253 (13)	0.0336 (15)	-0.0044 (12)	-0.0067 (12)	-0.0026 (11)
C8	0.0474 (18)	0.0315 (14)	0.0273 (13)	-0.0076 (13)	-0.0042 (12)	0.0019 (11)
C9	0.0348 (15)	0.0329 (14)	0.0251 (13)	-0.0078 (12)	-0.0024 (11)	0.0040 (11)
C10	0.0486 (19)	0.0471 (18)	0.0302 (15)	-0.0013 (15)	-0.0044 (13)	0.0087 (13)
C11	0.058 (2)	0.0365 (16)	0.0501 (19)	-0.0103 (15)	-0.0161 (16)	0.0150 (14)
C12	0.071 (2)	0.0336 (15)	0.0473 (19)	-0.0124 (16)	-0.0194 (17)	0.0016 (14)
C13	0.060 (2)	0.0358 (15)	0.0352 (15)	-0.0120 (15)	-0.0145 (14)	0.0008 (12)

Geometric parameters (Å, °)

Co1—O2W	2.0559 (18)	C2—C7	1.385 (4)
Co1—O2W ⁱ	2.0559 (18)	C2—C3	1.395 (4)
Co1—O1W ⁱ	2.094 (2)	C3—C4	1.383 (4)
Co1—O1W	2.094 (2)	C3—H3	0.9300
Co1—O3W ⁱ	2.141 (2)	C4—C5	1.384 (4)
Co1—O3W	2.141 (2)	C4—H4	0.9300
O1—C1	1.254 (3)	C5—C6	1.387 (4)
O2—C1	1.268 (3)	C6—C7	1.387 (4)
O3—C5	1.372 (3)	C6—H6	0.9300
O3—C8	1.420 (3)	C7—H7	0.9300
O1W—H11	0.837 (10)	C8—C9	1.507 (4)
O1W—H12	0.838 (10)	C8—H8A	0.9700
O2W—H21	0.841 (10)	C8—H8B	0.9700
O2W—H22	0.841 (10)	C9—C13	1.381 (4)
O3W—H31	0.847 (10)	C10—C11	1.372 (5)
O3W—H32	0.846 (10)	C10—H10	0.9300
O4W—H41	0.837 (10)	C11—C12	1.369 (5)
O4W—H42	0.839 (10)	C11—H11A	0.9300
N1—C9	1.334 (4)	C12—C13	1.386 (4)
N1—C10	1.344 (4)	C12—H12A	0.9300
C1—C2	1.497 (4)	C13—H13	0.9300
O2W—Co1—O2W ⁱ	180.000 (1)	C2—C3—H3	119.8
O2W—Co1—O1W ⁱ	93.59 (8)	C3—C4—C5	119.9 (3)
O2W ⁱ —Co1—O1W ⁱ	86.41 (8)	C3—C4—H4	120.1
O2W—Co1—O1W	86.41 (8)	C5—C4—H4	120.1
O2W ⁱ —Co1—O1W	93.59 (8)	O3—C5—C4	114.6 (2)
O1W ⁱ —Co1—O1W	180.0	O3—C5—C6	124.8 (3)

O2W—Co1—O3W ⁱ	86.46 (8)	C4—C5—C6	120.5 (2)
O2W ⁱ —Co1—O3W ⁱ	93.54 (8)	C5—C6—C7	119.0 (3)
O1W ⁱ —Co1—O3W ⁱ	92.45 (9)	C5—C6—H6	120.5
O1W—Co1—O3W ⁱ	87.55 (8)	C7—C6—H6	120.5
O2W—Co1—O3W	93.54 (8)	C2—C7—C6	121.3 (3)
O2W ⁱ —Co1—O3W	86.46 (8)	C2—C7—H7	119.3
O1W ⁱ —Co1—O3W	87.55 (8)	C6—C7—H7	119.3
O1W—Co1—O3W	92.45 (9)	O3—C8—C9	107.6 (2)
O3W ⁱ —Co1—O3W	180.000 (1)	O3—C8—H8A	110.2
C5—O3—C8	119.4 (2)	C9—C8—H8A	110.2
Co1—O1W—H11	106 (3)	O3—C8—H8B	110.2
Co1—O1W—H12	117 (3)	C9—C8—H8B	110.2
H11—O1W—H12	109.7 (17)	H8A—C8—H8B	108.5
Co1—O2W—H21	117 (2)	N1—C9—C13	122.4 (3)
Co1—O2W—H22	130 (2)	N1—C9—C8	115.3 (2)
H21—O2W—H22	109.4 (16)	C13—C9—C8	122.3 (2)
Co1—O3W—H31	104 (3)	N1—C10—C11	123.2 (3)
Co1—O3W—H32	117 (3)	N1—C10—H10	118.4
H31—O3W—H32	107.6 (16)	C11—C10—H10	118.4
H41—O4W—H42	109 (6)	C12—C11—C10	118.7 (3)
C9—N1—C10	117.8 (3)	C12—C11—H11A	120.7
O1—C1—O2	123.3 (2)	C10—C11—H11A	120.7
O1—C1—C2	119.1 (2)	C11—C12—C13	119.1 (3)
O2—C1—C2	117.6 (2)	C11—C12—H12A	120.4
C7—C2—C3	118.8 (2)	C13—C12—H12A	120.4
C7—C2—C1	120.9 (2)	C9—C13—C12	118.8 (3)
C3—C2—C1	120.4 (2)	C9—C13—H13	120.6
C4—C3—C2	120.5 (3)	C12—C13—H13	120.6
C4—C3—H3	119.8		
O1—C1—C2—C7	3.0 (4)	C1—C2—C7—C6	-179.5 (3)
O2—C1—C2—C7	-175.9 (3)	C5—C6—C7—C2	0.6 (5)
O1—C1—C2—C3	-176.1 (3)	C5—O3—C8—C9	176.6 (3)
O2—C1—C2—C3	5.1 (4)	C10—N1—C9—C13	-0.8 (5)
C7—C2—C3—C4	0.1 (5)	C10—N1—C9—C8	177.9 (3)
C1—C2—C3—C4	179.2 (3)	O3—C8—C9—N1	178.4 (3)
C2—C3—C4—C5	0.1 (5)	O3—C8—C9—C13	-2.9 (4)
C8—O3—C5—C4	-177.1 (3)	C9—N1—C10—C11	0.1 (5)
C8—O3—C5—C6	2.2 (5)	N1—C10—C11—C12	0.3 (6)
C3—C4—C5—O3	179.3 (3)	C10—C11—C12—C13	0.0 (6)
C3—C4—C5—C6	0.1 (5)	N1—C9—C13—C12	1.1 (5)
O3—C5—C6—C7	-179.6 (3)	C8—C9—C13—C12	-177.6 (3)
C4—C5—C6—C7	-0.4 (5)	C11—C12—C13—C9	-0.6 (5)
C3—C2—C7—C6	-0.5 (5)		

Symmetry codes: (i) $-x+1, -y+1, -z+1$.

supplementary materials

Hydrogen-bond geometry (\AA , $^\circ$)

$D-H\cdots A$	$D-H$	$H\cdots A$	$D\cdots A$	$D-H\cdots A$
O1w—H11 \cdots O1	0.84 (1)	1.94 (1)	2.762 (3)	169 (3)
O1w—H12 \cdots O4w ⁱⁱ	0.84 (1)	1.87 (1)	2.696 (4)	171 (4)
O2w—H21 \cdots O2	0.84 (1)	1.84 (1)	2.678 (3)	179 (3)
O2w—H22 \cdots O2 ⁱⁱⁱ	0.84 (1)	1.97 (2)	2.755 (3)	156 (3)
O3w—H31 \cdots O1 ⁱ	0.85 (1)	1.96 (1)	2.798 (3)	170 (4)
O3w—H32 \cdots N1 ^{iv}	0.85 (1)	1.95 (1)	2.799 (3)	177 (4)
O4w—H41 \cdots O2	0.84 (1)	2.22 (4)	2.891 (4)	137 (5)
O4w—H42 \cdots O3w ^v	0.84 (1)	2.42 (4)	3.176 (4)	151 (7)

Symmetry codes: (ii) $-x, -y+2, -z+1$; (iii) $-x+1, -y+2, -z+1$; (i) $-x+1, -y+1, -z+1$; (iv) $x, y-1, z+1$; (v) $x, y+1, z$.

Fig. 1

