

**catena-Poly[[[bis(4-bromobenzoato- $\kappa$ O)-zinc]- $\mu$ -1,2-bis(4-pyridyl)ethene- $\kappa^2$ N:N']acetonitrile monosolvate]**

Ni-Ya Li\* and Deng-Ming Sun

College of Chemistry and Materials Science, HuaiBei Normal University, HuaiBei 235000, Anhui, People's Republic of China  
Correspondence e-mail: niyali@chnu.edu.cn


Received 10 October 2011; accepted 12 October 2011

Key indicators: single-crystal X-ray study;  $T = 223\text{ K}$ ; mean  $\sigma(\text{C}-\text{C}) = 0.007\text{ \AA}$ ;  $R$  factor = 0.053;  $wR$  factor = 0.110; data-to-parameter ratio = 17.9.

In the title coordination compound,  $\{[\text{Zn}(\text{C}_7\text{H}_4\text{BrO}_2)_2(\text{C}_{12}\text{H}_{10}\text{N}_2)]\cdot\text{CH}_3\text{CN}\}_n$ , the  $\text{Zn}^{II}$  atom is four-coordinated in a distorted tetrahedral environment by two carboxylate O atoms from two different 4-bromobenzoate (bpe) ligands and two N atoms from two symmetry-related 1,2-bis(4-pyridyl)ethene ligands. The  $\text{Zn}^{II}$  atoms are bridged by the bpe ligands, which lie across centres of inversion, forming a zigzag chain along [001]. The void space of each unit cell is occupied by an acetonitrile solvent molecule, which is connected to the complex molecule by a weak C—H···N hydrogen bond.

## Related literature

For zigzag chains constructed by  $\text{Zn}^{II}$ , mono-carboxylate ligands and dipyridyl ligands, see: Gao *et al.* (2010); Kwak *et al.* (2009); Ng *et al.* (2004).



## Experimental

### Crystal data

$[\text{Zn}(\text{C}_7\text{H}_4\text{BrO}_2)_2(\text{C}_{12}\text{H}_{10}\text{N}_2)]\cdot\text{CH}_3\text{CN}$	$\beta = 94.63 (3)^\circ$
$M_r = 688.67$	$\gamma = 99.87 (3)^\circ$
Triclinic, $P\bar{1}$	$V = 1365.8 (5)\text{ \AA}^3$
$a = 6.2738 (13)\text{ \AA}$	$Z = 2$
$b = 11.852 (2)\text{ \AA}$	Mo $K\alpha$ radiation
$c = 19.496 (4)\text{ \AA}$	$\mu = 3.86\text{ mm}^{-1}$
$\alpha = 105.25 (3)^\circ$	$T = 223\text{ K}$
	$0.20 \times 0.10 \times 0.10\text{ mm}$

### Data collection

Rigaku MercuryCCD area-detector diffractometer	13203 measured reflections
Absorption correction: multi-scan ( <i>REQAB</i> ; Jacobson, 1998)	6152 independent reflections
$T_{\min} = 0.512$ , $T_{\max} = 0.699$	3934 reflections with $I > 2\sigma(I)$
	$R_{\text{int}} = 0.045$

### Refinement

$R[F^2 > 2\sigma(F^2)] = 0.053$	344 parameters
$wR(F^2) = 0.110$	H-atom parameters constrained
$S = 1.05$	$\Delta\rho_{\max} = 0.51\text{ e \AA}^{-3}$
6152 reflections	$\Delta\rho_{\min} = -0.54\text{ e \AA}^{-3}$

**Table 1**  
Hydrogen-bond geometry ( $\text{\AA}$ ,  $^\circ$ ).

$D-\text{H}\cdots A$	$D-\text{H}$	$\text{H}\cdots A$	$D\cdots A$	$D-\text{H}\cdots A$
$\text{Cl}2-\text{H}12\cdots\text{N}3$	0.94	2.53	3.436 (8)	162

Data collection: *CrystalClear* (Rigaku, 2001); cell refinement: *CrystalClear*; data reduction: *CrystalStructure* (Rigaku/MSC, 2004); program(s) used to solve structure: *SHELXTL* (Sheldrick, 2008); program(s) used to refine structure: *SHELXTL*; molecular graphics: *SHELXTL*; software used to prepare material for publication: *SHELXTL* and *PLATON* (Spek, 2009).

This work was supported by the research start-up fund for new staff of HuaiBei Normal University (600581).

Supplementary data and figures for this paper are available from the IUCr electronic archives (Reference: BQ2311).

## References

- Gao, Y., Yang, X., Zhang, J., Li, G. & Zhang, H. (2010). *J. Coord. Chem.* **63**, 3413–3422.
- Jacobson, R. (1998). *REQAB*. Private communication to the Rigaku Corporation, Tokyo, Japan.
- Kwak, H., Lee, S. H., Kim, S. H., Lee, Y. M., Park, B. K., Lee, Y. J., Jun, J. Y., Kim, C., Kim, S. J. & Kim, Y. (2009). *Polyhedron*, **28**, 553–561.
- Ng, M. T., Deivaraj, T. C., Klooster, W. T., McIntyre, G. J. & Vittal, J. J. (2004). *Chem. Eur. J.* **10**, 5853–5859.
- Rigaku (2001). *CrystalClear*. Rigaku Corporation, Tokyo, Japan.
- Rigaku/MSC (2004). *CrystalStructure*. Rigaku/MSC, The Woodlands, Texas, USA.
- Sheldrick, G. M. (2008). *Acta Cryst. A* **64**, 112–122.
- Spek, A. L. (2009). *Acta Cryst. D* **65**, 148–155.

## **supplementary materials**

*Acta Cryst.* (2011). E67, m1553 [doi:10.1107/S1600536811042188]

**[*catena-Poly[[[bis(4-bromobenzoato- $\kappa O$ )zinc]- $\mu$ -1,2-bis(4-pyridyl)ethene- $\kappa^2 N;N'$ ] monosolvate*]** acetonitrile

**N.-Y. Li and D.-M. Sun**

### Comment

In recent years, a large number of coordination polymers assembled from carboxylates and pyridyl-like ligands have been extensively investigated. Among these coordination polymers, most of them are constructed by polycarboxylates and dipyridyl ligands, the complexes assembled from mono-carboxylate and dipyridyl ligands are still rare.

In this work, 4-bromobenzoate (BBA) and 1,2-bis(4-pyridyl)ethene (bpe) were employed to react with Zn<sup>II</sup> and thus afford the title complex,  $\{[\text{Zn}(\text{C}_7\text{H}_4\text{O}_2\text{Br})_2(\text{C}_{12}\text{H}_{10}\text{N}_2)].(\text{C}_2\text{H}_3\text{N})\}_n$  (I). In (I), the Zn<sup>II</sup> atom lies on a twofold rotation axis that relates one BBA ligand to the other as well as one bpe ligand to the other; the coordination geometry is a distorted tetrahedron (Fig. 1). The Zn—O (1.958 (3)–1.985 (4) Å) and Zn—N (2.049 (3)–2.060 (3) Å) bond lengths are comparable to those reported for similar complexes (Gao *et al.*, 2010; Kwak *et al.*, 2009; Ng *et al.*, 2004). The Zn<sup>II</sup> centres are linked by the bpe ligands to form a one-dimensional zigzag chain (Fig. 2). A non-coordinated solvent molecule of acetonitrile occupies the interstitial voids within the unit cell. It is noted that there is a weak C12—H12···N3 (Table 1) intermolecular hydrogen bond in the structure (Fig. 2). This weak interaction connects the main molecule with the solvent molecule.

### Experimental

To a 25 ml Teflon-lined stainless steel autoclave was loaded Zn(NO<sub>3</sub>)<sub>2</sub>.6H<sub>2</sub>O (149 mg, 0.5 mmol), 4-bromobenzoic acid (200 mg, 1 mmol), 1,2-bis(4-pyridyl)ethene (91 mg, 0.5 mmol), H<sub>2</sub>O (8 ml) and acetonitrile (8 ml). The autoclave was sealed and heated in an oven to 423 K for three days, and then cooled to ambient temperature at the rate of 5 K/h to form yellow crystals of (I). Yield: 238 mg (69% yield based on Zn). Anal. calcd. for C<sub>28</sub>H<sub>21</sub>Br<sub>2</sub>N<sub>3</sub>O<sub>4</sub>Zn: C, 48.83; H, 3.07; N, 6.10. Found: C, 50.05; H, 3.22; N, 6.37.

### Refinement

The C-bound H atoms were positioned geometrically, with C—H = 0.97 Å (methyl) or 0.94 Å (phenyl, pyridyl and vinyl), and refined as riding, with  $U_{\text{iso}}(\text{H}) = 1.5U_{\text{eq}}(\text{C})$  for methyl groups or  $1.2U_{\text{eq}}(\text{C})$  otherwise.

# supplementary materials

---

## Figures




Fig. 1. Coordination environment of  $\text{Zn}^{\text{II}}$  atom in the compound with nonhydrogen atoms represented by thermal ellipsoids draw at 30% probability level. [Symmetry codes, i:  $-x - 1, -y + 1, -z + 1$ ; ii:  $-x, -y + 2, -z + 2$ .]




Fig. 2. View of the unit-cell contents of the title compound showing weak intermolecular H-bond with dashed lines.

## **catena-Poly[[[bis(4-bromobenzoato- $\kappa O$ )zinc]- $\mu$ -1,2-bis(4-pyridyl)ethene- $\kappa^2 N:N'$ ] acetonitrile monosolvate]**

### Crystal data

$[\text{Zn}(\text{C}_7\text{H}_4\text{BrO}_2)_2(\text{C}_{12}\text{H}_{10}\text{N}_2)] \cdot \text{C}_2\text{H}_3\text{N}$	$Z = 2$
$M_r = 688.67$	$F(000) = 684$
Triclinic, $P\bar{1}$	$D_x = 1.675 \text{ Mg m}^{-3}$
Hall symbol: -P 1	Mo $K\alpha$ radiation, $\lambda = 0.71073 \text{ \AA}$
$a = 6.2738 (13) \text{ \AA}$	Cell parameters from 5974 reflections
$b = 11.852 (2) \text{ \AA}$	$\theta = 3.2\text{--}27.5^\circ$
$c = 19.496 (4) \text{ \AA}$	$\mu = 3.86 \text{ mm}^{-1}$
$\alpha = 105.25 (3)^\circ$	$T = 223 \text{ K}$
$\beta = 94.63 (3)^\circ$	Block, yellow
$\gamma = 99.87 (3)^\circ$	$0.20 \times 0.10 \times 0.10 \text{ mm}$
$V = 1365.8 (5) \text{ \AA}^3$	

### Data collection

Rigaku MercuryCCD area-detector diffractometer	6152 independent reflections
Radiation source: fine-focus sealed tube graphite	3934 reflections with $I > 2\sigma(I)$
$\omega$ scans	$R_{\text{int}} = 0.045$
Absorption correction: multi-scan (REQAB; Jacobson, 1998)	$\theta_{\text{max}} = 27.5^\circ, \theta_{\text{min}} = 3.2^\circ$
$T_{\text{min}} = 0.512, T_{\text{max}} = 0.699$	$h = -7 \rightarrow 8$
13203 measured reflections	$k = -15 \rightarrow 15$
	$l = -25 \rightarrow 21$

## *Refinement*

Refinement on $F^2$	Primary atom site location: structure-invariant direct methods
Least-squares matrix: full	Secondary atom site location: difference Fourier map
$R[F^2 > 2\sigma(F^2)] = 0.053$	Hydrogen site location: inferred from neighbouring sites
$wR(F^2) = 0.110$	H-atom parameters constrained
$S = 1.05$	$w = 1/[\sigma^2(F_o^2) + (0.0378P)^2]$ where $P = (F_o^2 + 2F_c^2)/3$
6152 reflections	$(\Delta/\sigma)_{\max} = 0.001$
344 parameters	$\Delta\rho_{\max} = 0.51 \text{ e \AA}^{-3}$
0 restraints	$\Delta\rho_{\min} = -0.54 \text{ e \AA}^{-3}$

## *Special details*

**Geometry.** All e.s.d.'s (except the e.s.d. in the dihedral angle between two l.s. planes) are estimated using the full covariance matrix. The cell e.s.d.'s are taken into account individually in the estimation of e.s.d.'s in distances, angles and torsion angles; correlations between e.s.d.'s in cell parameters are only used when they are defined by crystal symmetry. An approximate (isotropic) treatment of cell e.s.d.'s is used for estimating e.s.d.'s involving l.s. planes.

**Refinement.** Refinement of  $F^2$  against ALL reflections. The weighted  $R$ -factor  $wR$  and goodness of fit  $S$  are based on  $F^2$ , conventional  $R$ -factors  $R$  are based on  $F$ , with  $F$  set to zero for negative  $F^2$ . The threshold expression of  $F^2 > \sigma(F^2)$  is used only for calculating  $R$ -factors(gt) etc. and is not relevant to the choice of reflections for refinement.  $R$ -factors based on  $F^2$  are statistically about twice as large as those based on  $F$ , and  $R$ -factors based on ALL data will be even larger.

## *Fractional atomic coordinates and isotropic or equivalent isotropic displacement parameters ( $\text{\AA}^2$ )*

	<i>x</i>	<i>y</i>	<i>z</i>	$U_{\text{iso}}^*/U_{\text{eq}}$
Zn1	0.44987 (7)	0.71792 (4)	0.71734 (3)	0.03921 (15)
Br1	1.37712 (10)	1.24685 (5)	0.59537 (3)	0.0756 (2)
Br2	0.94349 (9)	0.17043 (4)	0.87084 (3)	0.06577 (18)
N1	0.1772 (5)	0.6343 (3)	0.64483 (16)	0.0374 (8)
N2	0.3188 (5)	0.8162 (3)	0.80109 (18)	0.0401 (8)
O1	0.6984 (7)	0.8429 (3)	0.7125 (3)	0.0948 (15)
O2	0.4838 (8)	0.8466 (5)	0.6190 (4)	0.148 (3)
O3	0.6068 (4)	0.5980 (2)	0.73677 (15)	0.0438 (7)
O4	0.3088 (5)	0.5406 (3)	0.78312 (16)	0.0513 (8)
C1	0.1024 (7)	0.5156 (4)	0.6262 (2)	0.0431 (10)
H1	0.1853	0.4677	0.6438	0.052*
C2	-0.0907 (7)	0.4620 (4)	0.5825 (2)	0.0412 (10)
H2	-0.1379	0.3788	0.5711	0.049*
C3	-0.2171 (6)	0.5292 (3)	0.5550 (2)	0.0387 (10)
C4	-0.1384 (7)	0.6510 (4)	0.5738 (2)	0.0449 (11)
H4	-0.2182	0.7006	0.5567	0.054*
C5	0.0578 (7)	0.6997 (4)	0.6177 (2)	0.0455 (11)
H5	0.1099	0.7825	0.6291	0.055*

## supplementary materials

---

C6	-0.4231 (6)	0.4716 (4)	0.5084 (2)	0.0398 (10)
H6	-0.4489	0.3883	0.4887	0.048*
C7	0.4393 (7)	0.9177 (3)	0.8455 (2)	0.0448 (11)
H7	0.5797	0.9455	0.8355	0.054*
C8	0.3661 (7)	0.9821 (4)	0.9046 (2)	0.0445 (11)
H8	0.4549	1.0537	0.9334	0.053*
C9	0.1640 (7)	0.9436 (4)	0.9224 (2)	0.0419 (10)
C10	0.0358 (7)	0.8385 (4)	0.8758 (2)	0.0500 (11)
H10	-0.1054	0.8095	0.8846	0.060*
C11	0.1194 (7)	0.7783 (4)	0.8169 (2)	0.0495 (11)
H11	0.0329	0.7075	0.7863	0.059*
C12	0.0908 (7)	1.0147 (4)	0.9876 (2)	0.0475 (11)
H12	0.1836	1.0872	1.0132	0.057*
C13	0.6547 (13)	0.8825 (6)	0.6615 (5)	0.094 (2)
C14	0.8324 (9)	0.9765 (4)	0.6480 (3)	0.0626 (15)
C15	0.7939 (9)	1.0245 (5)	0.5934 (3)	0.0739 (16)
H15	0.6559	1.0015	0.5656	0.089*
C16	0.9529 (8)	1.1066 (4)	0.5776 (3)	0.0633 (14)
H16	0.9235	1.1401	0.5401	0.076*
C17	1.1540 (8)	1.1382 (4)	0.6178 (3)	0.0521 (12)
C18	1.1993 (8)	1.0931 (4)	0.6739 (3)	0.0566 (12)
H18	1.3377	1.1165	0.7015	0.068*
C19	1.0344 (9)	1.0115 (4)	0.6889 (3)	0.0620 (14)
H19	1.0615	0.9800	0.7275	0.074*
C20	0.4967 (7)	0.5330 (3)	0.7693 (2)	0.0388 (10)
C21	0.6034 (6)	0.4408 (3)	0.7914 (2)	0.0381 (9)
C22	0.5102 (7)	0.3804 (4)	0.8367 (2)	0.0505 (11)
H22	0.3778	0.3959	0.8525	0.061*
C23	0.6067 (7)	0.2979 (4)	0.8591 (2)	0.0529 (12)
H23	0.5431	0.2579	0.8904	0.063*
C24	0.7997 (7)	0.2758 (3)	0.8343 (2)	0.0439 (10)
C25	0.8937 (7)	0.3311 (4)	0.7881 (2)	0.0484 (11)
H25	1.0233	0.3130	0.7710	0.058*
C26	0.7951 (7)	0.4148 (4)	0.7667 (2)	0.0451 (10)
H26	0.8590	0.4541	0.7351	0.054*
C27	0.8391 (11)	1.4073 (6)	1.0550 (3)	0.099 (2)
H27A	0.8972	1.4411	1.1053	0.148*
H27B	0.8351	1.4712	1.0327	0.148*
H27C	0.9315	1.3557	1.0313	0.148*
C28	0.6195 (13)	1.3385 (6)	1.0485 (3)	0.092 (2)
N3	0.4488 (11)	1.2851 (5)	1.0428 (3)	0.121 (2)

Atomic displacement parameters ( $\text{\AA}^2$ )

	$U^{11}$	$U^{22}$	$U^{33}$	$U^{12}$	$U^{13}$	$U^{23}$
Zn1	0.0397 (3)	0.0422 (3)	0.0404 (3)	0.0136 (2)	0.0121 (2)	0.0142 (2)
Br1	0.0754 (4)	0.0562 (3)	0.1030 (5)	0.0027 (3)	0.0328 (3)	0.0365 (3)
Br2	0.0704 (4)	0.0566 (3)	0.0812 (4)	0.0200 (3)	0.0006 (3)	0.0365 (3)

N1	0.046 (2)	0.0385 (18)	0.0321 (18)	0.0153 (17)	0.0117 (16)	0.0110 (15)
N2	0.0378 (19)	0.0420 (19)	0.043 (2)	0.0104 (17)	0.0123 (16)	0.0129 (16)
O1	0.129 (4)	0.051 (2)	0.130 (4)	0.029 (2)	0.092 (3)	0.041 (2)
O2	0.072 (3)	0.126 (4)	0.265 (8)	-0.003 (3)	0.060 (4)	0.093 (5)
O3	0.0402 (16)	0.0477 (17)	0.0550 (18)	0.0112 (14)	0.0136 (14)	0.0302 (15)
O4	0.0408 (17)	0.064 (2)	0.063 (2)	0.0195 (15)	0.0179 (15)	0.0325 (16)
C1	0.045 (2)	0.046 (3)	0.044 (2)	0.018 (2)	0.015 (2)	0.016 (2)
C2	0.046 (2)	0.036 (2)	0.044 (2)	0.008 (2)	0.015 (2)	0.0129 (19)
C3	0.041 (2)	0.042 (2)	0.036 (2)	0.009 (2)	0.0153 (19)	0.0131 (19)
C4	0.045 (2)	0.041 (2)	0.051 (3)	0.015 (2)	0.006 (2)	0.012 (2)
C5	0.050 (3)	0.035 (2)	0.051 (3)	0.015 (2)	0.012 (2)	0.006 (2)
C6	0.041 (2)	0.042 (2)	0.036 (2)	0.0057 (19)	0.0120 (19)	0.0109 (18)
C7	0.044 (2)	0.038 (2)	0.051 (3)	0.004 (2)	0.014 (2)	0.010 (2)
C8	0.054 (3)	0.038 (2)	0.043 (2)	0.014 (2)	0.012 (2)	0.0110 (19)
C9	0.046 (3)	0.047 (2)	0.041 (2)	0.022 (2)	0.011 (2)	0.016 (2)
C10	0.036 (2)	0.056 (3)	0.059 (3)	0.012 (2)	0.019 (2)	0.012 (2)
C11	0.044 (3)	0.051 (3)	0.052 (3)	0.012 (2)	0.008 (2)	0.009 (2)
C12	0.052 (3)	0.042 (2)	0.052 (3)	0.012 (2)	0.012 (2)	0.017 (2)
C13	0.091 (5)	0.056 (4)	0.156 (7)	0.031 (4)	0.083 (5)	0.037 (4)
C14	0.064 (3)	0.043 (3)	0.097 (4)	0.021 (3)	0.050 (3)	0.030 (3)
C15	0.053 (3)	0.066 (3)	0.108 (5)	0.013 (3)	0.021 (3)	0.030 (3)
C16	0.063 (3)	0.060 (3)	0.080 (4)	0.017 (3)	0.017 (3)	0.037 (3)
C17	0.056 (3)	0.041 (2)	0.070 (3)	0.013 (2)	0.027 (3)	0.027 (2)
C18	0.065 (3)	0.047 (3)	0.062 (3)	0.012 (2)	0.017 (3)	0.021 (2)
C19	0.088 (4)	0.046 (3)	0.068 (3)	0.023 (3)	0.038 (3)	0.030 (3)
C20	0.041 (2)	0.037 (2)	0.036 (2)	0.0059 (19)	0.0045 (19)	0.0066 (19)
C21	0.033 (2)	0.041 (2)	0.041 (2)	0.0015 (18)	0.0040 (18)	0.0168 (19)
C22	0.039 (2)	0.064 (3)	0.058 (3)	0.012 (2)	0.018 (2)	0.030 (2)
C23	0.053 (3)	0.061 (3)	0.057 (3)	0.012 (2)	0.014 (2)	0.037 (2)
C24	0.046 (3)	0.038 (2)	0.050 (3)	0.007 (2)	-0.003 (2)	0.018 (2)
C25	0.041 (2)	0.051 (3)	0.059 (3)	0.009 (2)	0.012 (2)	0.025 (2)
C26	0.043 (2)	0.045 (2)	0.055 (3)	0.008 (2)	0.014 (2)	0.026 (2)
C27	0.115 (5)	0.091 (4)	0.079 (4)	-0.020 (4)	0.035 (4)	0.023 (3)
C28	0.127 (6)	0.076 (4)	0.057 (4)	-0.017 (4)	0.019 (4)	0.014 (3)
N3	0.120 (5)	0.114 (5)	0.089 (4)	-0.053 (4)	0.012 (4)	0.011 (3)

*Geometric parameters (Å, °)*

Zn1—O3	1.958 (3)	C10—H10	0.9400
Zn1—O1	1.985 (4)	C11—H11	0.9400
Zn1—N1	2.049 (3)	C12—C12 <sup>ii</sup>	1.302 (8)
Zn1—N2	2.060 (3)	C12—H12	0.9400
Br1—C17	1.894 (5)	C13—C14	1.526 (8)
Br2—C24	1.906 (4)	C14—C15	1.357 (7)
N1—C5	1.338 (5)	C14—C19	1.377 (7)
N1—C1	1.345 (5)	C15—C16	1.382 (7)
N2—C7	1.342 (5)	C15—H15	0.9400
N2—C11	1.343 (5)	C16—C17	1.367 (7)
O1—C13	1.236 (9)	C16—H16	0.9400

## supplementary materials

---

O2—C13	1.238 (9)	C17—C18	1.367 (7)
O3—C20	1.272 (5)	C18—C19	1.396 (7)
O4—C20	1.243 (5)	C18—H18	0.9400
C1—C2	1.372 (6)	C19—H19	0.9400
C1—H1	0.9400	C20—C21	1.510 (5)
C2—C3	1.386 (5)	C21—C22	1.380 (6)
C2—H2	0.9400	C21—C26	1.384 (5)
C3—C4	1.382 (5)	C22—C23	1.378 (6)
C3—C6	1.464 (5)	C22—H22	0.9400
C4—C5	1.382 (6)	C23—C24	1.381 (6)
C4—H4	0.9400	C23—H23	0.9400
C5—H5	0.9400	C24—C25	1.360 (6)
C6—C6 <sup>i</sup>	1.333 (7)	C25—C26	1.389 (6)
C6—H6	0.9400	C25—H25	0.9400
C7—C8	1.365 (5)	C26—H26	0.9400
C7—H7	0.9400	C27—C28	1.453 (9)
C8—C9	1.375 (6)	C27—H27A	0.9700
C8—H8	0.9400	C27—H27B	0.9700
C9—C10	1.403 (6)	C27—H27C	0.9700
C9—C12	1.484 (6)	C28—N3	1.128 (8)
C10—C11	1.377 (6)		
O3—Zn1—O1	100.35 (16)	O1—C13—O2	124.7 (7)
O3—Zn1—N1	109.30 (12)	O1—C13—C14	116.8 (8)
O1—Zn1—N1	129.94 (19)	O2—C13—C14	118.4 (8)
O3—Zn1—N2	117.36 (13)	C15—C14—C19	118.9 (5)
O1—Zn1—N2	98.90 (14)	C15—C14—C13	119.7 (7)
N1—Zn1—N2	101.82 (13)	C19—C14—C13	121.3 (6)
C5—N1—C1	117.4 (3)	C14—C15—C16	121.6 (5)
C5—N1—Zn1	119.6 (3)	C14—C15—H15	119.2
C1—N1—Zn1	122.9 (3)	C16—C15—H15	119.2
C7—N2—C11	117.4 (4)	C17—C16—C15	118.6 (5)
C7—N2—Zn1	120.1 (3)	C17—C16—H16	120.7
C11—N2—Zn1	122.3 (3)	C15—C16—H16	120.7
C13—O1—Zn1	109.7 (5)	C18—C17—C16	121.8 (5)
C20—O3—Zn1	111.2 (2)	C18—C17—Br1	118.8 (4)
N1—C1—C2	122.2 (4)	C16—C17—Br1	119.4 (4)
N1—C1—H1	118.9	C17—C18—C19	118.1 (5)
C2—C1—H1	118.9	C17—C18—H18	120.9
C1—C2—C3	120.8 (4)	C19—C18—H18	120.9
C1—C2—H2	119.6	C14—C19—C18	120.9 (5)
C3—C2—H2	119.6	C14—C19—H19	119.6
C4—C3—C2	116.7 (4)	C18—C19—H19	119.6
C4—C3—C6	122.7 (4)	O4—C20—O3	124.0 (4)
C2—C3—C6	120.6 (4)	O4—C20—C21	119.0 (4)
C5—C4—C3	119.8 (4)	O3—C20—C21	117.0 (3)
C5—C4—H4	120.1	C22—C21—C26	118.7 (4)
C3—C4—H4	120.1	C22—C21—C20	120.3 (4)
N1—C5—C4	123.0 (4)	C26—C21—C20	121.0 (4)

N1—C5—H5	118.5	C23—C22—C21	121.6 (4)
C4—C5—H5	118.5	C23—C22—H22	119.2
C6 <sup>i</sup> —C6—C3	124.9 (5)	C21—C22—H22	119.2
C6 <sup>i</sup> —C6—H6	117.5	C22—C23—C24	118.0 (4)
C3—C6—H6	117.5	C22—C23—H23	121.0
N2—C7—C8	122.6 (4)	C24—C23—H23	121.0
N2—C7—H7	118.7	C25—C24—C23	122.2 (4)
C8—C7—H7	118.7	C25—C24—Br2	119.9 (3)
C7—C8—C9	120.8 (4)	C23—C24—Br2	117.9 (4)
C7—C8—H8	119.6	C24—C25—C26	118.9 (4)
C9—C8—H8	119.6	C24—C25—H25	120.6
C8—C9—C10	117.0 (4)	C26—C25—H25	120.6
C8—C9—C12	119.5 (4)	C21—C26—C25	120.5 (4)
C10—C9—C12	123.5 (4)	C21—C26—H26	119.7
C11—C10—C9	119.1 (4)	C25—C26—H26	119.7
C11—C10—H10	120.5	C28—C27—H27A	109.5
C9—C10—H10	120.5	C28—C27—H27B	109.5
N2—C11—C10	123.1 (4)	H27A—C27—H27B	109.5
N2—C11—H11	118.5	C28—C27—H27C	109.5
C10—C11—H11	118.5	H27A—C27—H27C	109.5
C12 <sup>ii</sup> —C12—C9	125.7 (6)	H27B—C27—H27C	109.5
C12 <sup>ii</sup> —C12—H12	117.2	N3—C28—C27	179.3 (7)
C9—C12—H12	117.2		

Symmetry codes: (i)  $-x-1, -y+1, -z+1$ ; (ii)  $-x, -y+2, -z+2$ .

#### *Hydrogen-bond geometry ( $\text{\AA}$ , $^\circ$ )*

$D\text{—H}\cdots A$	$D\text{—H}$	$H\cdots A$	$D\cdots A$	$D\text{—H}\cdots A$
C12—H12 $\cdots$ N3	0.94	2.53	3.436 (8)	162.

## supplementary materials

---

Fig. 1




Fig. 2

