

N-(1-Naphthyl)benzenesulfonamide

Sifang Zhang, Yuwen Zhang, Chuntao Wang and Ruitao Zhu*

Department of Chemistry, Taiyuan Normal University, Taiyuan 030031, People's Republic of China

Correspondence e-mail: ruitaozhu@126.com


Received 14 September 2011; accepted 24 September 2011

Key indicators: single-crystal X-ray study; $T = 298\text{ K}$; mean $\sigma(\text{C}-\text{C}) = 0.005\text{ \AA}$; R factor = 0.035; wR factor = 0.084; data-to-parameter ratio = 13.4.

In the title compound, $\text{C}_{16}\text{H}_{13}\text{NO}_2\text{S}$, the $\text{C}-\text{SO}_2-\text{NH}-\text{C}$ torsion angle is $-70.1(2)^\circ$. The dihedral angle between the planes of the naphthyl ring system and the phenyl ring is $34.67(4)^\circ$. In the crystal, molecules are linked by intermolecular $\text{N}-\text{H}\cdots\text{O}$ hydrogen bonds into chains along [100]. There are also $\pi-\pi$ interactions between adjacent naphthyl groups [interplanar spacing = $3.541(3)\text{ \AA}$] for molecules stacked along [100].

Related literature

For hydrogen-bonding modes of sulfonamides, see: Adsmond & Grant (2001). For related structures, see: Shakuntala *et al.* (2011). For standard bond-length data, see: Allen *et al.* (1987).

**Experimental***Crystal data*

$\text{C}_{16}\text{H}_{13}\text{NO}_2\text{S}$
 $M_r = 283.33$

Orthorhombic, $P2_12_12_1$
 $a = 4.9232(5)\text{ \AA}$

$b = 15.4162(15)\text{ \AA}$
 $c = 18.2102(17)\text{ \AA}$
 $V = 1382.1(2)\text{ \AA}^3$
 $Z = 4$

Mo $K\alpha$ radiation
 $\mu = 0.23\text{ mm}^{-1}$
 $T = 298\text{ K}$
 $0.43 \times 0.33 \times 0.32\text{ mm}$

Data collection

Bruker SMART CCD area-detector diffractometer
Absorption correction: multi-scan (*SADABS*; Bruker, 2007)
 $T_{\min} = 0.906$, $T_{\max} = 0.929$

6917 measured reflections
2438 independent reflections
2178 reflections with $I > \sigma(I)$
 $R_{\text{int}} = 0.032$

Refinement

$R[F^2 > 2\sigma(F^2)] = 0.035$
 $wR(F^2) = 0.084$
 $S = 1.09$
2438 reflections
182 parameters
H-atom parameters constrained

$\Delta\rho_{\max} = 0.18\text{ e \AA}^{-3}$
 $\Delta\rho_{\min} = -0.20\text{ e \AA}^{-3}$
Absolute structure: Flack (1983),
983 Friedel pairs
Flack parameter: 0.06 (9)

Table 1
Hydrogen-bond geometry (\AA , $^\circ$).

$D-\text{H}\cdots A$	$D-\text{H}$	$\text{H}\cdots A$	$D\cdots A$	$D-\text{H}\cdots A$
$\text{N1}-\text{H1}\cdots\text{O1}^i$	0.90	2.05	2.911 (3)	159

Symmetry code: (i) $x + 1, y, z$.

Data collection: *SMART* (Bruker, 2007); cell refinement: *SAINT* (Bruker, 2007); data reduction: *SAINT*; program(s) used to solve structure: *SHELXS97* (Sheldrick, 2008); program(s) used to refine structure: *SHELXL97* (Sheldrick, 2008); molecular graphics: *SHELXTL* (Sheldrick, 2008); software used to prepare material for publication: *SHELXTL*.

We would like to thank the Natural Science Foundation of Shanxi Province for financial support (2011011007-2).

Supplementary data and figures for this paper are available from the IUCr electronic archives (Reference: PK2348).

References

- Adsmond, D. A. & Grant, D. W. (2001). *J. Pharm. Sci.* **90**, 2058–2077.
- Allen, F. H., Kennard, O., Watson, D. G., Brammer, L., Orpen, A. G. & Taylor, R. (1987). *J. Chem. Soc. Perkin Trans. 2*, pp. S1–19.
- Bruker (2007). *SMART*, *SAINT* and *SADABS*. Bruker AXS Inc., Madison, Wisconsin, USA.
- Flack, H. D. (1983). *Acta Cryst. A* **39**, 876–881.
- Shakuntala, K., Foro, S. & Gowda, B. T. (2011). *Acta Cryst. E* **67**, o1540.
- Sheldrick, G. M. (2008). *Acta Cryst. A* **64**, 112–122.

supplementary materials

Acta Cryst. (2011). E67, o2831 [doi:10.1107/S1600536811039201]

N-(1-Naphthyl)benzenesulfonamide

S. Zhang, Y. Zhang, C. Wang and R. Zhu

Comment

Sulfonamide moieties are constituents of many biologically important compounds. The hydrogen bonding preferences of sulfonamides has been investigated (Adsmond & Grant, 2001). In this paper, we present the crystal structure of the title compound.

The molecular structure of is shown in Fig. 1. The bond lengths (Allen *et al.*, 1987) and angles are normal. The molecule is twisted at the S atom with C—SO₂—NH—C torsion angle of -70.14 (2) °. The dihedral between the planes of the naphthyl and benzene groups is 34.67 (4) °. In the crystal, molecules are linked by intermolecular N—H···O hydrogen bonds into chains along [100]. There are also π – π interactions between adjacent naphthyl groups (interplanar spacing 3.541 (3) Å) for molecules stacked along [100].

Experimental

To a 100 ml round flask fitted with a condenser was added 1-naphthylamine (1.43 g, 10 mmol), dichloromethane (15 ml) and triethylamine(0.5 ml) with magnetic stirring. Benzenesulfonyl chloride (1.76 g, 10 mmol) was added gradually. The reaction mixture was stirred at room temperature for 1 h and then refluxed for 2 h. The product precipitated as a white powder, which was washed three times with water and dichloromethane. Recrystallization from ethyl alcohol produced the crystals of the title compound.

Refinement

H atoms were placed in idealized positions and allowed to ride on their respective parent atoms, with C—H = 0.93 Å, N—H = 0.90 Å and $U_{\text{iso}}(\text{H}) = 1.2U_{\text{eq}}(\text{C}, \text{N})$.

Figures




Fig. 1. The molecular structure of the title compound with displacement ellipsoids are drawn at the 30% probability level.




Fig. 2. Part of the crystal structure with hydrogen bonds drawn as dashed lines. Only H atoms involved in hydrogen bonds are shown.

supplementary materials

N-(1-Naphthyl)benzenesulfonamide

Crystal data

C ₁₆ H ₁₃ NO ₂ S	<i>F</i> (000) = 592
<i>M_r</i> = 283.33	<i>D_x</i> = 1.362 Mg m ⁻³
Orthorhombic, <i>P</i> 2 ₁ 2 ₁ 2 ₁	Mo <i>Kα</i> radiation, λ = 0.71073 Å
Hall symbol: P 2ac 2ab	Cell parameters from 3370 reflections
<i>a</i> = 4.9232 (5) Å	θ = 2.6–26.0°
<i>b</i> = 15.4162 (15) Å	μ = 0.23 mm ⁻¹
<i>c</i> = 18.2102 (17) Å	<i>T</i> = 298 K
<i>V</i> = 1382.1 (2) Å ³	Block-like, colorless
<i>Z</i> = 4	0.43 × 0.33 × 0.32 mm

Data collection

Bruker SMART CCD area-detector diffractometer	2438 independent reflections
Radiation source: fine-focus sealed tube graphite	2178 reflections with $I > \sigma(I)$
φ and ω scans	R_{int} = 0.032
Absorption correction: multi-scan (<i>SADABS</i> ; Bruker, 2007)	$\theta_{\text{max}} = 25.0^\circ$, $\theta_{\text{min}} = 2.6^\circ$
$T_{\text{min}} = 0.906$, $T_{\text{max}} = 0.929$	$h = -5 \rightarrow 5$
6917 measured reflections	$k = -17 \rightarrow 18$
	$l = -14 \rightarrow 21$

Refinement

Refinement on F^2	Hydrogen site location: inferred from neighbouring sites
Least-squares matrix: full	H-atom parameters constrained
$R[F^2 > 2\sigma(F^2)] = 0.035$	$w = 1/[\sigma^2(F_o^2) + (0.037P)^2 + 0.2812P]$
$wR(F^2) = 0.084$	where $P = (F_o^2 + 2F_c^2)/3$
$S = 1.09$	$(\Delta/\sigma)_{\text{max}} < 0.001$
2438 reflections	$\Delta\rho_{\text{max}} = 0.18 \text{ e \AA}^{-3}$
182 parameters	$\Delta\rho_{\text{min}} = -0.20 \text{ e \AA}^{-3}$
0 restraints	Extinction correction: <i>SHELXL97</i> (Sheldrick, 2008), $F_c^* = kF_c[1 + 0.001x F_c^2 \lambda^3 / \sin(2\theta)]^{-1/4}$
Primary atom site location: structure-invariant direct methods	Extinction coefficient: 0.038 (2)
Secondary atom site location: difference Fourier map	Absolute structure: Flack (1983), 983 Friedel pairs
	Flack parameter: 0.06 (9)

Special details

Geometry. All e.s.d.'s (except the e.s.d. in the dihedral angle between two l.s. planes) are estimated using the full covariance matrix. The cell e.s.d.'s are taken into account individually in the estimation of e.s.d.'s in distances, angles and torsion angles; correlations between e.s.d.'s in cell parameters are only used when they are defined by crystal symmetry. An approximate (isotropic) treatment of cell e.s.d.'s is used for estimating e.s.d.'s involving l.s. planes.

Refinement. Refinement of F^2 against ALL reflections. The weighted R -factor wR and goodness of fit S are based on F^2 , conventional R -factors R are based on F , with F set to zero for negative F^2 . The threshold expression of $F^2 > \sigma(F^2)$ is used only for calculating R -factors(gt) etc. and is not relevant to the choice of reflections for refinement. R -factors based on F^2 are statistically about twice as large as those based on F , and R -factors based on ALL data will be even larger.

Fractional atomic coordinates and isotropic or equivalent isotropic displacement parameters (\AA^2)

	<i>x</i>	<i>y</i>	<i>z</i>	$U_{\text{iso}}^*/U_{\text{eq}}$
S1	0.63000 (11)	0.47539 (3)	0.28894 (3)	0.03525 (17)
N1	0.7982 (4)	0.50023 (12)	0.36336 (10)	0.0388 (5)
H1	0.9726	0.4850	0.3559	0.047*
O1	0.3486 (3)	0.48108 (11)	0.30720 (9)	0.0496 (4)
O2	0.7374 (3)	0.52862 (11)	0.23221 (9)	0.0486 (4)
C1	0.7144 (5)	0.46260 (17)	0.43237 (13)	0.0430 (6)
C2	0.8252 (6)	0.38695 (18)	0.45628 (15)	0.0574 (7)
H2	0.9547	0.3589	0.4277	0.069*
C3	0.7467 (9)	0.3508 (2)	0.52341 (18)	0.0800 (11)
H3	0.8277	0.2999	0.5400	0.096*
C4	0.5555 (9)	0.3894 (3)	0.56351 (18)	0.0824 (12)
H4	0.4994	0.3634	0.6070	0.099*
C5	0.4352 (6)	0.4686 (3)	0.54214 (14)	0.0673 (9)
C6	0.5182 (5)	0.50785 (18)	0.47415 (13)	0.0503 (7)
C7	0.4088 (6)	0.5887 (2)	0.45330 (16)	0.0617 (8)
H7	0.4671	0.6155	0.4103	0.074*
C8	0.2160 (7)	0.6281 (3)	0.4966 (2)	0.0894 (12)
H8	0.1439	0.6815	0.4831	0.107*
C9	0.1278 (8)	0.5871 (4)	0.5617 (2)	0.1007 (15)
H9	-0.0060	0.6132	0.5902	0.121*
C10	0.2347 (8)	0.5107 (4)	0.5831 (2)	0.0960 (14)
H10	0.1733	0.4853	0.6264	0.115*
C11	0.6966 (5)	0.36643 (14)	0.26609 (12)	0.0375 (6)
C12	0.8977 (6)	0.34830 (16)	0.21627 (15)	0.0541 (6)
H12	0.9960	0.3927	0.1942	0.065*
C13	0.9509 (7)	0.2619 (2)	0.19960 (19)	0.0767 (10)
H13	1.0869	0.2477	0.1663	0.092*
C14	0.7995 (8)	0.1971 (2)	0.2331 (2)	0.0815 (12)
H14	0.8348	0.1393	0.2220	0.098*
C15	0.6022 (8)	0.21681 (18)	0.28156 (19)	0.0747 (9)
H15	0.5027	0.1725	0.3034	0.090*
C16	0.5467 (6)	0.30098 (16)	0.29891 (15)	0.0553 (7)
H16	0.4100	0.3143	0.3323	0.066*

supplementary materials

Atomic displacement parameters (\AA^2)

	U^{11}	U^{22}	U^{33}	U^{12}	U^{13}	U^{23}
S1	0.0321 (3)	0.0373 (3)	0.0364 (3)	0.0023 (3)	0.0012 (2)	0.0003 (3)
N1	0.0274 (10)	0.0482 (11)	0.0407 (10)	-0.0036 (7)	0.0033 (8)	-0.0040 (8)
O1	0.0304 (8)	0.0613 (10)	0.0570 (10)	0.0060 (9)	-0.0004 (8)	-0.0087 (9)
O2	0.0576 (10)	0.0435 (8)	0.0448 (9)	0.0013 (9)	0.0034 (8)	0.0097 (8)
C1	0.0377 (13)	0.0569 (15)	0.0344 (12)	-0.0124 (12)	-0.0021 (10)	-0.0027 (11)
C2	0.0598 (18)	0.0633 (17)	0.0491 (15)	-0.0032 (14)	-0.0107 (14)	0.0038 (13)
C3	0.105 (3)	0.079 (2)	0.055 (2)	-0.016 (2)	-0.009 (2)	0.0166 (17)
C4	0.101 (3)	0.104 (3)	0.0423 (18)	-0.044 (2)	-0.0106 (19)	0.0179 (18)
C5	0.0537 (18)	0.114 (3)	0.0340 (13)	-0.0348 (19)	0.0049 (13)	-0.0198 (17)
C6	0.0384 (13)	0.0705 (18)	0.0420 (13)	-0.0158 (13)	0.0000 (11)	-0.0140 (12)
C7	0.0512 (18)	0.083 (2)	0.0513 (16)	-0.0034 (15)	0.0029 (14)	-0.0252 (15)
C8	0.067 (2)	0.119 (3)	0.082 (2)	0.020 (2)	-0.0036 (19)	-0.046 (2)
C9	0.058 (2)	0.178 (4)	0.066 (2)	-0.009 (3)	0.020 (2)	-0.067 (3)
C10	0.070 (2)	0.161 (4)	0.057 (2)	-0.042 (3)	0.0155 (18)	-0.034 (3)
C11	0.0370 (13)	0.0366 (11)	0.0389 (12)	0.0012 (10)	-0.0069 (10)	-0.0002 (9)
C12	0.0507 (15)	0.0522 (13)	0.0594 (15)	0.0053 (13)	0.0022 (15)	-0.0121 (13)
C13	0.064 (2)	0.075 (2)	0.091 (2)	0.0250 (17)	-0.0058 (19)	-0.0371 (19)
C14	0.089 (3)	0.0419 (15)	0.113 (3)	0.0147 (17)	-0.046 (2)	-0.0218 (18)
C15	0.092 (2)	0.0398 (14)	0.092 (2)	-0.0081 (16)	-0.021 (2)	0.0030 (16)
C16	0.0582 (17)	0.0461 (14)	0.0615 (17)	-0.0063 (12)	-0.0032 (15)	0.0029 (13)

Geometric parameters (\AA , $^\circ$)

S1—O2	1.4214 (16)	C7—H7	0.9300
S1—O1	1.4273 (17)	C8—C9	1.411 (6)
S1—N1	1.6337 (19)	C8—H8	0.9300
S1—C11	1.761 (2)	C9—C10	1.348 (6)
N1—C1	1.444 (3)	C9—H9	0.9300
N1—H1	0.9000	C10—H10	0.9300
C1—C2	1.359 (4)	C11—C12	1.372 (3)
C1—C6	1.414 (4)	C11—C16	1.386 (3)
C2—C3	1.398 (4)	C12—C13	1.391 (4)
C2—H2	0.9300	C12—H12	0.9300
C3—C4	1.332 (5)	C13—C14	1.388 (5)
C3—H3	0.9300	C13—H13	0.9300
C4—C5	1.411 (5)	C14—C15	1.347 (5)
C4—H4	0.9300	C14—H14	0.9300
C5—C10	1.397 (5)	C15—C16	1.363 (4)
C5—C6	1.437 (4)	C15—H15	0.9300
C6—C7	1.410 (4)	C16—H16	0.9300
C7—C8	1.376 (4)		
O2—S1—O1	119.67 (11)	C6—C7—H7	120.0
O2—S1—N1	106.20 (10)	C7—C8—C9	119.8 (4)
O1—S1—N1	106.52 (10)	C7—C8—H8	120.1

O2—S1—C11	108.04 (11)	C9—C8—H8	120.1
O1—S1—C11	107.11 (11)	C10—C9—C8	121.0 (3)
N1—S1—C11	108.98 (10)	C10—C9—H9	119.5
C1—N1—S1	118.87 (14)	C8—C9—H9	119.5
C1—N1—H1	107.4	C9—C10—C5	121.8 (4)
S1—N1—H1	107.3	C9—C10—H10	119.1
C2—C1—C6	121.7 (2)	C5—C10—H10	119.1
C2—C1—N1	120.6 (2)	C12—C11—C16	121.4 (2)
C6—C1—N1	117.7 (2)	C12—C11—S1	119.00 (18)
C1—C2—C3	120.8 (3)	C16—C11—S1	119.57 (19)
C1—C2—H2	119.6	C11—C12—C13	118.4 (3)
C3—C2—H2	119.6	C11—C12—H12	120.8
C4—C3—C2	119.7 (3)	C13—C12—H12	120.8
C4—C3—H3	120.1	C14—C13—C12	119.5 (3)
C2—C3—H3	120.1	C14—C13—H13	120.3
C3—C4—C5	122.2 (3)	C12—C13—H13	120.3
C3—C4—H4	118.9	C15—C14—C13	120.9 (3)
C5—C4—H4	118.9	C15—C14—H14	119.6
C10—C5—C4	123.4 (4)	C13—C14—H14	119.6
C10—C5—C6	117.7 (4)	C14—C15—C16	120.7 (3)
C4—C5—C6	118.8 (3)	C14—C15—H15	119.6
C7—C6—C1	123.5 (2)	C16—C15—H15	119.6
C7—C6—C5	119.7 (3)	C15—C16—C11	119.1 (3)
C1—C6—C5	116.8 (3)	C15—C16—H16	120.4
C8—C7—C6	119.9 (3)	C11—C16—H16	120.4
C8—C7—H7	120.0		
O2—S1—N1—C1	173.69 (18)	C5—C6—C7—C8	-2.5 (4)
O1—S1—N1—C1	45.1 (2)	C6—C7—C8—C9	-0.2 (5)
C11—S1—N1—C1	-70.1 (2)	C7—C8—C9—C10	1.6 (6)
S1—N1—C1—C2	91.7 (2)	C8—C9—C10—C5	-0.3 (6)
S1—N1—C1—C6	-89.3 (2)	C4—C5—C10—C9	178.8 (3)
C6—C1—C2—C3	0.4 (4)	C6—C5—C10—C9	-2.4 (5)
N1—C1—C2—C3	179.3 (3)	O2—S1—C11—C12	19.9 (2)
C1—C2—C3—C4	2.0 (5)	O1—S1—C11—C12	150.09 (19)
C2—C3—C4—C5	-2.8 (5)	N1—S1—C11—C12	-95.1 (2)
C3—C4—C5—C10	-179.9 (3)	O2—S1—C11—C16	-160.45 (19)
C3—C4—C5—C6	1.2 (4)	O1—S1—C11—C16	-30.3 (2)
C2—C1—C6—C7	176.5 (2)	N1—S1—C11—C16	84.6 (2)
N1—C1—C6—C7	-2.4 (3)	C16—C11—C12—C13	-0.6 (4)
C2—C1—C6—C5	-1.9 (3)	S1—C11—C12—C13	179.0 (2)
N1—C1—C6—C5	179.2 (2)	C11—C12—C13—C14	0.4 (4)
C10—C5—C6—C7	3.7 (4)	C12—C13—C14—C15	-0.1 (5)
C4—C5—C6—C7	-177.4 (3)	C13—C14—C15—C16	-0.1 (5)
C10—C5—C6—C1	-177.9 (2)	C14—C15—C16—C11	-0.1 (5)
C4—C5—C6—C1	1.1 (4)	C12—C11—C16—C15	0.5 (4)
C1—C6—C7—C8	179.2 (3)	S1—C11—C16—C15	-179.2 (2)


supplementary materials

Hydrogen-bond geometry (Å, °)

$D\text{---H}\cdots A$	$D\text{---H}$	$H\cdots A$	$D\cdots A$	$D\text{---H}\cdots A$
N1—H1 ⁱ —O1 ⁱ	0.90	2.05	2.911 (3)	159.

Symmetry codes: (i) $x+1, y, z$.

Fig. 1



supplementary materials

Fig. 2

