

2-(2-Hydroxyphenyl)-1,3-benzothiazole-6-carbaldehyde

Kew-Yu Chen,* Tzu-Chien Fang, Ming-Jen Chang,
Hsing-Yang Tsai and Ming-Hui Luo

Department of Chemical Engineering, Feng Chia University, 40724 Taichung,
Taiwan

Correspondence e-mail: kyuchen@fcu.edu.tw


Received 30 September 2011; accepted 3 October 2011

Key indicators: single-crystal X-ray study; $T = 150\text{ K}$; mean $\sigma(\text{C}-\text{C}) = 0.002\text{ \AA}$;
 R factor = 0.029; wR factor = 0.058; data-to-parameter ratio = 11.6.

The molecule of the title compound, $\text{C}_{14}\text{H}_9\text{NO}_2\text{S}$, is nearly planar, the maximum atomic deviation being $0.081(2)\text{ \AA}$. An intramolecular $\text{O}-\text{H}\cdots\text{N}$ bond generates an $S(6)$ ring motif. In the crystal, inversion-related molecules linked by a pair of weak $\text{C}-\text{H}\cdots\text{O}$ hydrogen bonds form a supramolecular dimer. $\pi-\pi$ stacking is observed between the thiazole and benzene rings of adjacent molecules, the centroid–centroid distance being $3.7679(9)\text{ \AA}$.

Related literature

For the spectroscopy and preparation of the title compound, see: Hsieh *et al.* (2008). For the spectroscopy and applications of benzoxazole and benzothiazole derivatives, see: Chen & Pang (2009, 2010); Hrobáriková *et al.* (2010); Kim *et al.* (2010a,b); Lijima *et al.* (2010); Lim *et al.* (2011); López-Ruiz *et al.* (2011); Tanaka *et al.* (2001). For related structures, see: Tong (2005); Hahn *et al.* (1998). For graph-set theory, see: Bernstein *et al.* (1995).



Experimental

Crystal data

$\text{C}_{14}\text{H}_9\text{NO}_2\text{S}$

$M_r = 255.28$

Monoclinic, $P2_1/n$

$a = 8.2645(3)\text{ \AA}$

$b = 5.6449(2)\text{ \AA}$

$c = 23.8341(9)\text{ \AA}$

$\beta = 98.147(2)^\circ$

$V = 1100.69(7)\text{ \AA}^3$

$Z = 4$

Mo $K\alpha$ radiation

$\mu = 0.29\text{ mm}^{-1}$

$T = 150\text{ K}$

$0.38 \times 0.14 \times 0.04\text{ mm}$

Data collection

Bruker SMART CCD area-detector

diffractometer

Absorption correction: multi-scan

(*SADABS*; Bruker, 2001)

$T_{\min} = 0.882$, $T_{\max} = 0.992$

8427 measured reflections

1943 independent reflections

1333 reflections with $I > 2\sigma(I)$

$R_{\text{int}} = 0.042$

Refinement

$R[F^2 > 2\sigma(F^2)] = 0.029$

$wR(F^2) = 0.058$

$S = 0.90$

1943 reflections

168 parameters

1 restraint

H atoms treated by a mixture of
independent and constrained
refinement

$\Delta\rho_{\max} = 0.22\text{ e \AA}^{-3}$

$\Delta\rho_{\min} = -0.27\text{ e \AA}^{-3}$

Table 1
Hydrogen-bond geometry (\AA , $^\circ$).

$D-\text{H}\cdots A$	$D-\text{H}$	$\text{H}\cdots A$	$D\cdots A$	$D-\text{H}\cdots A$
O2—H2 \cdots N1	0.89 (2)	1.81 (2)	2.6228 (18)	150 (2)
C5—H5 \cdots O2 ⁱ	0.93	2.61	3.293 (2)	130

Symmetry code: (i) $-x + 2, -y + 3, -z + 1$.

Data collection: *SMART* (Bruker, 2001); cell refinement: *SAINT* (Bruker, 2001); data reduction: *SAINT*; program(s) used to solve structure: *SHELXS97* (Sheldrick, 2008); program(s) used to refine structure: *SHELXL97* (Sheldrick, 2008); molecular graphics: *ORTEP-3 for Windows* (Farrugia, 1997); software used to prepare material for publication: *WinGX* (Farrugia, 1999).

This work was supported by the National Science Council and Feng Chia University in Taiwan.

Supplementary data and figures for this paper are available from the IUCr electronic archives (Reference: XU5345).

References

- Bernstein, J., Davis, R. E., Shimoni, L. & Chang, N.-L. (1995). *Angew. Chem. Int. Ed. Engl.* **34**, 1555–1573.
Bruker (2001). *SMART*, *SAINT* and *SADABS*. Bruker AXS Inc., Madison, Wisconsin, USA.
Chen, W.-H. & Pang, Y. (2009). *Tetrahedron Lett.* **50**, 6680–6683.
Chen, W.-H. & Pang, Y. (2010). *Tetrahedron Lett.* **51**, 1914–1918.
Farrugia, L. J. (1997). *J. Appl. Cryst.* **30**, 565.
Farrugia, L. J. (1999). *J. Appl. Cryst.* **32**, 837–838.
Hahn, F. E., Imhof, L. & Lügger, T. (1998). *Acta Cryst. C* **54**, 668–669.
Hrobáriková, V., Hrobárik, P., Gajdos, P., Fililis, I., Fakis, M., Persephonis, P. & Zahradník, P. (2010). *J. Org. Chem.* **75**, 3053–3068.
Hsieh, C.-C., Cheng, Y.-M., Hsu, C.-J., Chen, K.-Y. & Chou, P.-T. (2008). *J. Phys. Chem. A*, **112**, 8323–8332.
Kim, T. H., Kwon, N. Y. & Lee, T. S. (2010a). *Tetrahedron Lett.* **51**, 5596–5600.
Kim, C. H., Park, J., Seo, J., Park, S. Y. & Joo, T. (2010b). *J. Phys. Chem. A*, **114**, 5618–5629.
Lijima, T., Momotake, A., Shinohara, Y., Sato, T., Nishimura, Y. & Arai, T. (2010). *J. Phys. Chem. A*, **114**, 1603–1609.
Lim, C. K., Seo, J., Kim, S., Kwon, I. C., Ahn, C. H. & Park, S. Y. (2011). *Dyes Pigments*, **90**, 284–289.
López-Ruiz, H., Briseno-Ortega, H., Rojas-Lima, S., Santillan, R. & Farfán, N. (2011). *Tetrahedron Lett.* **52**, 4308–4312.
Sheldrick, G. M. (2008). *Acta Cryst. A* **64**, 112–122.
Tanaka, K., Kumagai, T., Aoki, H., Deguchi, M. & Iwata, S. (2001). *J. Org. Chem.* **66**, 7328–7333.
Tong, Y.-P. (2005). *Acta Cryst. E* **61**, o3076–o3078.

supplementary materials

Acta Cryst. (2011). E67, o2862 [doi:10.1107/S1600536811040712]

2-(2-Hydroxyphenyl)-1,3-benzothiazole-6-carbaldehyde

K.-Y. Chen, T.-C. Fang, M.-J. Chang, H.-Y. Tsai and M.-H. Luo

Comment

The excited-state intramolecular proton transfer (ESIPT) reaction of 2-(2-hydroxyphenyl)benzoxazole and 2-(2-hydroxyphenyl)benzothiazole derivatives has been investigated for past years (Hsieh *et al.*, 2008; Kim *et al.*, 2010*a,b*; Lijima *et al.*, 2010; López-Ruiz *et al.*, 2011), which incorporates transfer of a hydroxy proton to the imine nitrogen through a intramolecular six-membered-ring hydrogen-bonding system (Chen *et al.*, 2009, 2010). The unusual photophysical property of the resulting proton-transfer tautomer has found many important applications (Hrobáriková *et al.*, 2010; Lim *et al.*, 2011; Tanaka *et al.*, 2001).

The molecular structure of the title compound (HBT) is shown in Figure 1. The molecule is nearly planar, which is consistent with previous studies (Tong, 2005; Hahn *et al.*, 1998). HBT possesses an intramolecular O—H···N hydrogen bond (Table 1), which generates an S(6) ring motif (Bernstein *et al.*, 1995). In the crystal (Figure 2), inversion-related molecules are linked by a pair of weak C—H···O hydrogen bonds, forming a cyclic dimers with $R_2^2(18)$ graph-set motif. $\pi\cdots\pi$ stacking is observed between thiazole and C1¹-benzene rings of adjacent molecules [symmetry code: (i): 2-x,2-y,1-z], the centroid-to-centroid distance being 3.7679 (9) Å.

Experimental

The title compound was synthesized according to the literature (Hsieh *et al.*, 2008). Yellow needle-shaped crystals suitable for the crystallographic studies reported here were isolated over a period of five weeks by slow evaporation from the chloroform solution.

Refinement

H atoms bonded to O and C atoms were located in a difference electron density map. The hydroxy H atom and the C_{sp3} H atoms were freely refined, and the C_{sp2} H atoms repositioned geometrically and refined using a riding model, [C—H = 0.93 Å and $U_{\text{iso}}(\text{H}) = 1.2U_{\text{eq}}(\text{C})$].

Figures




Fig. 1. The molecular structure of the title compound, showing 50% probability displacement ellipsoids.




Fig. 2. A section of the crystal packing of the title compound, viewed down the *b* axis. Green dashed lines denote the intermolecular C—H···O hydrogen bonds.

supplementary materials

2-(2-Hydroxyphenyl)-1,3-benzothiazole-6-carbaldehyde

Crystal data

C ₁₄ H ₉ NO ₂ S	<i>F</i> (000) = 528
<i>M_r</i> = 255.28	<i>D_x</i> = 1.541 Mg m ⁻³
Monoclinic, <i>P2₁/n</i>	Mo <i>Kα</i> radiation, λ = 0.71073 Å
Hall symbol: -p 2yn	Cell parameters from 2297 reflections
<i>a</i> = 8.2645 (3) Å	θ = 2.5–25.7°
<i>b</i> = 5.6449 (2) Å	μ = 0.29 mm ⁻¹
<i>c</i> = 23.8341 (9) Å	<i>T</i> = 150 K
β = 98.147 (2)°	Plate, yellow
<i>V</i> = 1100.69 (7) Å ³	0.38 × 0.14 × 0.04 mm
<i>Z</i> = 4	

Data collection

Bruker SMART CCD area-detector diffractometer	1943 independent reflections
Radiation source: fine-focus sealed tube graphite	1333 reflections with $I > 2\sigma(I)$
φ and ω scans	$R_{\text{int}} = 0.042$
Absorption correction: multi-scan (<i>SADABS</i> ; Bruker, 2001)	$\theta_{\text{max}} = 25.0^\circ$, $\theta_{\text{min}} = 1.7^\circ$
$T_{\text{min}} = 0.882$, $T_{\text{max}} = 0.992$	$h = -9 \rightarrow 9$
8427 measured reflections	$k = -6 \rightarrow 3$
	$l = -28 \rightarrow 28$

Refinement

Refinement on F^2	Secondary atom site location: difference Fourier map
Least-squares matrix: full	Hydrogen site location: inferred from neighbouring sites
$R[F^2 > 2\sigma(F^2)] = 0.029$	H atoms treated by a mixture of independent and constrained refinement
$wR(F^2) = 0.058$	$w = 1/[\sigma^2(F_o^2) + (0.0251P)^2]$ where $P = (F_o^2 + 2F_c^2)/3$
$S = 0.90$	$(\Delta/\sigma)_{\text{max}} < 0.001$
1943 reflections	$\Delta\rho_{\text{max}} = 0.22 \text{ e \AA}^{-3}$
168 parameters	$\Delta\rho_{\text{min}} = -0.27 \text{ e \AA}^{-3}$
1 restraint	Extinction correction: <i>SHELXL97</i> (Sheldrick, 2008), $F_c^* = kF_c[1 + 0.001xF_c^2\lambda^3/\sin(2\theta)]^{1/4}$
Primary atom site location: structure-invariant direct methods	Extinction coefficient: 0.0014 (2)

Special details

Geometry. All e.s.d.'s (except the e.s.d. in the dihedral angle between two l.s. planes) are estimated using the full covariance matrix. The cell e.s.d.'s are taken into account individually in the estimation of e.s.d.'s in distances, angles and torsion angles; correlations between e.s.d.'s in cell parameters are only used when they are defined by crystal symmetry. An approximate (isotropic) treatment of cell e.s.d.'s is used for estimating e.s.d.'s involving l.s. planes.

Refinement. Refinement of F^2 against ALL reflections. The weighted R -factor wR and goodness of fit S are based on F^2 , conventional R -factors R are based on F , with F set to zero for negative F^2 . The threshold expression of $F^2 > \sigma(F^2)$ is used only for calculating R -factors(gt) etc. and is not relevant to the choice of reflections for refinement. R -factors based on F^2 are statistically about twice as large as those based on F , and R -factors based on ALL data will be even larger.

Fractional atomic coordinates and isotropic or equivalent isotropic displacement parameters (\AA^2)

	x	y	z	$U_{\text{iso}}^*/U_{\text{eq}}$
S	0.74344 (6)	0.79801 (8)	0.392532 (18)	0.02213 (16)
O1	0.51794 (15)	0.6205 (2)	0.59718 (5)	0.0317 (4)
O2	1.05490 (15)	1.4470 (2)	0.35839 (5)	0.0272 (3)
N1	0.88683 (16)	1.1973 (2)	0.42282 (6)	0.0183 (3)
C1	0.81872 (19)	1.1165 (3)	0.46928 (7)	0.0167 (4)
C2	0.7362 (2)	0.8986 (3)	0.46067 (7)	0.0169 (4)
C3	0.66326 (19)	0.7940 (3)	0.50315 (7)	0.0189 (4)
H3	0.6076	0.6510	0.4970	0.023*
C4	0.67509 (19)	0.9071 (3)	0.55528 (7)	0.0183 (4)
C5	0.7578 (2)	1.1247 (3)	0.56406 (7)	0.0211 (4)
H5	0.7644	1.1987	0.5992	0.025*
C6	0.82896 (19)	1.2299 (3)	0.52162 (7)	0.0204 (4)
H6	0.8830	1.3741	0.5277	0.024*
C7	0.5997 (2)	0.7988 (3)	0.60128 (7)	0.0252 (4)
H7	0.6162	0.8736	0.6364	0.030*
C11	0.91907 (19)	1.0821 (3)	0.32631 (7)	0.0168 (4)
C12	1.0157 (2)	1.2802 (3)	0.31764 (7)	0.0190 (4)
C13	1.07588 (19)	1.3083 (3)	0.26659 (7)	0.0219 (4)
H13	1.1400	1.4394	0.2611	0.026*
C14	1.04146 (19)	1.1439 (3)	0.22413 (7)	0.0229 (5)
H14	1.0826	1.1644	0.1901	0.027*
C15	0.9459 (2)	0.9477 (3)	0.23149 (7)	0.0239 (5)
H15	0.9225	0.8371	0.2026	0.029*
C16	0.8862 (2)	0.9183 (3)	0.28209 (7)	0.0222 (4)
H16	0.8223	0.7863	0.2870	0.027*
C17	0.85778 (19)	1.0478 (3)	0.38021 (7)	0.0180 (4)
H2	1.005 (2)	1.407 (4)	0.3877 (7)	0.071 (8)*

Atomic displacement parameters (\AA^2)

	U^{11}	U^{22}	U^{33}	U^{12}	U^{13}	U^{23}
S	0.0278 (3)	0.0195 (3)	0.0199 (3)	-0.0044 (2)	0.0062 (2)	-0.0013 (2)

supplementary materials

O1	0.0349 (8)	0.0318 (8)	0.0295 (8)	-0.0074 (7)	0.0085 (7)	0.0053 (6)
O2	0.0346 (8)	0.0245 (8)	0.0233 (8)	-0.0092 (6)	0.0070 (7)	-0.0026 (6)
N1	0.0198 (9)	0.0164 (8)	0.0186 (8)	0.0005 (7)	0.0025 (7)	0.0009 (7)
C1	0.0154 (10)	0.0160 (10)	0.0181 (10)	0.0040 (8)	0.0003 (8)	0.0037 (8)
C2	0.0172 (10)	0.0169 (10)	0.0165 (10)	0.0029 (8)	0.0016 (8)	0.0006 (8)
C3	0.0162 (10)	0.0168 (10)	0.0233 (11)	0.0011 (8)	0.0016 (8)	0.0021 (9)
C4	0.0170 (10)	0.0194 (10)	0.0187 (10)	0.0043 (8)	0.0034 (8)	0.0037 (8)
C5	0.0226 (11)	0.0225 (11)	0.0179 (11)	0.0051 (9)	0.0019 (9)	-0.0020 (8)
C6	0.0209 (11)	0.0178 (10)	0.0220 (11)	-0.0004 (8)	0.0011 (8)	-0.0001 (8)
C7	0.0241 (12)	0.0299 (11)	0.0215 (11)	0.0069 (10)	0.0030 (9)	0.0008 (9)
C11	0.0162 (10)	0.0173 (10)	0.0171 (10)	0.0008 (8)	0.0028 (8)	0.0023 (8)
C12	0.0183 (10)	0.0184 (10)	0.0194 (10)	0.0019 (9)	-0.0003 (8)	-0.0006 (9)
C13	0.0207 (11)	0.0220 (10)	0.0233 (11)	-0.0037 (8)	0.0048 (9)	0.0048 (9)
C14	0.0202 (11)	0.0292 (12)	0.0201 (11)	0.0032 (9)	0.0060 (9)	0.0051 (9)
C15	0.0279 (12)	0.0240 (11)	0.0198 (11)	0.0016 (9)	0.0040 (9)	-0.0030 (8)
C16	0.0238 (11)	0.0184 (10)	0.0244 (11)	-0.0023 (9)	0.0038 (9)	0.0011 (9)
C17	0.0146 (10)	0.0163 (10)	0.0228 (11)	0.0030 (8)	0.0016 (8)	0.0022 (8)

Geometric parameters (\AA , $^\circ$)

S—C2	1.7296 (16)	C5—H5	0.9300
S—C17	1.7454 (17)	C6—H6	0.9300
O1—C7	1.2087 (19)	C7—H7	0.9300
O2—C12	1.3588 (19)	C11—C16	1.400 (2)
O2—H2	0.890 (14)	C11—C12	1.406 (2)
N1—C17	1.3160 (18)	C11—C17	1.459 (2)
N1—C1	1.3882 (19)	C12—C13	1.387 (2)
C1—C6	1.394 (2)	C13—C14	1.373 (2)
C1—C2	1.407 (2)	C13—H13	0.9300
C2—C3	1.382 (2)	C14—C15	1.386 (2)
C3—C4	1.388 (2)	C14—H14	0.9300
C3—H3	0.9300	C15—C16	1.376 (2)
C4—C5	1.407 (2)	C15—H15	0.9300
C4—C7	1.469 (2)	C16—H16	0.9300
C5—C6	1.375 (2)		
C2—S—C17	89.10 (8)	C4—C7—H7	117.4
C12—O2—H2	107.2 (14)	C16—C11—C12	117.93 (15)
C17—N1—C1	110.79 (14)	C16—C11—C17	121.39 (15)
N1—C1—C6	125.67 (16)	C12—C11—C17	120.67 (15)
N1—C1—C2	114.43 (15)	O2—C12—C13	117.94 (15)
C6—C1—C2	119.88 (16)	O2—C12—C11	121.94 (15)
C3—C2—C1	121.30 (16)	C13—C12—C11	120.11 (16)
C3—C2—S	128.62 (14)	C14—C13—C12	120.42 (16)
C1—C2—S	110.07 (12)	C14—C13—H13	119.8
C2—C3—C4	118.44 (16)	C12—C13—H13	119.8
C2—C3—H3	120.8	C13—C14—C15	120.61 (16)
C4—C3—H3	120.8	C13—C14—H14	119.7
C3—C4—C5	120.40 (16)	C15—C14—H14	119.7
C3—C4—C7	119.55 (16)	C16—C15—C14	119.28 (16)

supplementary materials

C5—C4—C7	120.05 (16)	C16—C15—H15	120.4
C6—C5—C4	121.13 (16)	C14—C15—H15	120.4
C6—C5—H5	119.4	C15—C16—C11	121.64 (16)
C4—C5—H5	119.4	C15—C16—H16	119.2
C5—C6—C1	118.84 (16)	C11—C16—H16	119.2
C5—C6—H6	120.6	N1—C17—C11	123.12 (15)
C1—C6—H6	120.6	N1—C17—S	115.59 (12)
O1—C7—C4	125.29 (17)	C11—C17—S	121.27 (12)
O1—C7—H7	117.4		

Hydrogen-bond geometry (Å, °)

<i>D</i> —H··· <i>A</i>	<i>D</i> —H	H··· <i>A</i>	<i>D</i> ··· <i>A</i>	<i>D</i> —H··· <i>A</i>
O2—H2···N1	0.89 (2)	1.81 (2)	2.6228 (18)	150 (2)
C5—H5···O2 ⁱ	0.93	2.61	3.293 (2)	130

Symmetry codes: (i) $-x+2, -y+3, -z+1$.

supplementary materials

Fig. 1




Fig. 2

