

4-Chloro-*N*-(3-methylphenyl)benzamideVinola Z. Rodrigues,^a Marek Fronc,^b B. Thimme Gowda^{a*} and Jozef Kožíšek^b^aDepartment of Chemistry, Mangalore University, Mangalagangothri 574 199, Mangalore, India, and ^bInstitute of Physical Chemistry and Chemical Physics, Slovak University of Technology, Radlinského 9, SK-812 37 Bratislava, Slovak Republic
Correspondence e-mail: gowdabt@yahoo.com


Received 2 October 2011; accepted 4 October 2011

Key indicators: single-crystal X-ray study; $T = 293$ K; mean $\sigma(\text{C}-\text{C}) = 0.003$ Å; R factor = 0.040; wR factor = 0.117; data-to-parameter ratio = 14.0.

In the title compound, $\text{C}_{14}\text{H}_{12}\text{ClNO}$, the *meta*-methyl substituent in the aniline ring is positioned *anti* to the $\text{N}-\text{H}$ bond. The dihedral angle between the rings is 12.4 (1)°. The crystal structure is stabilized by intermolecular $\text{N}-\text{H}\cdots\text{O}$ hydrogen bonds, which link the molecules into $C(4)$ chains running along the c -axis direction.

Related literature

For the preparation of the title compound, see: Gowda *et al.* (2003). For studies on the effects of substituents on the structures and other aspects of *N*-(aryl)-amides, see: Bowes *et al.* (2003); Gowda *et al.* (2000); Saeed *et al.* (2010), on *N*-(aryl)-methanesulfonamides, see: Gowda *et al.* (2007), on *N*-(aryl)-arylsulfonamides, see: Shetty & Gowda (2005) and on *N*-chloro-arylsulfonamides, see: Gowda & Shetty (2004).



Experimental

Crystal data

 $\text{C}_{14}\text{H}_{12}\text{ClNO}$
 $M_r = 245.70$
Monoclinic, $P2_1/c$
 $a = 13.4379$ (9) Å
 $b = 10.2493$ (11) Å
 $c = 9.2600$ (7) Å
 $\beta = 92.893$ (6)° $V = 1273.74$ (19) Å³
 $Z = 4$
Mo $K\alpha$ radiation
 $\mu = 0.28$ mm⁻¹
 $T = 293$ K
 $0.81 \times 0.17 \times 0.04$ mm

Data collection

Oxford Diffraction Xcalibur diffractometer with a Ruby (Gemini Cu) detector
Absorption correction: analytical [CrysAlis RED (Oxford Diffraction, 2009)], based on expressions derived from Clark &Reid (1995)]
 $T_{\min} = 0.877$, $T_{\max} = 0.988$
17187 measured reflections
2156 independent reflections
1299 reflections with $I > 2\sigma(I)$
 $R_{\text{int}} = 0.040$

Refinement

 $R[F^2 > 2\sigma(F^2)] = 0.040$
 $wR(F^2) = 0.117$
 $S = 0.94$
2156 reflections154 parameters
H-atom parameters constrained
 $\Delta\rho_{\max} = 0.22$ e Å⁻³
 $\Delta\rho_{\min} = -0.13$ e Å⁻³

Table 1

Hydrogen-bond geometry (Å, °).

$D-\text{H}\cdots A$	$D-\text{H}$	$\text{H}\cdots A$	$D\cdots A$	$D-\text{H}\cdots A$
$\text{N1}-\text{H1A}\cdots\text{O1}^i$	0.86	2.06	2.888 (2)	161

Symmetry code: (i) $x, -y + \frac{1}{2}, z + \frac{1}{2}$.

Data collection: CrysAlis CCD (Oxford Diffraction, 2009); cell refinement: CrysAlis CCD; data reduction: CrysAlis RED (Oxford Diffraction, 2009); program(s) used to solve structure: SHELXS97 (Sheldrick, 2008); program(s) used to refine structure: SHELXL97 (Sheldrick, 2008); molecular graphics: Mercury (Macrae *et al.*, 2008); software used to prepare material for publication: enCIFer (Allen *et al.*, 2004).

MF and JK thank the Grant Agencies for their financial support: VEGA Grant Agency of Slovak Ministry of Education 1/0679/11; Research and Development Agency (Slovakia) APVV-0202-10 and the Structural Funds, Interreg IIIA, for financial support in purchasing the diffractometer. VZR thanks the University Grants Commission, Government of India, New Delhi, for the award of an RFSMS research fellowship.

Supplementary data and figures for this paper are available from the IUCr electronic archives (Reference: BT5661).

References

- Allen, F. H., Johnson, O., Shields, G. P., Smith, B. R. & Towler, M. (2004). *J. Appl. Cryst.* **37**, 335–338.
- Bowes, K. F., Glidewell, C., Low, J. N., Skakle, J. M. S. & Wardell, J. L. (2003). *Acta Cryst.* **C59**, o1–o3.
- Clark, R. C. & Reid, J. S. (1995). *Acta Cryst.* **A51**, 887–897.
- Gowda, B. T., Foro, S. & Fuess, H. (2007). *Acta Cryst.* **E63**, o2339.
- Gowda, B. T., Jyothi, K., Paulus, H. & Fuess, H. (2003). *Z. Naturforsch. Teil A*, **58**, 225–230.
- Gowda, B. T. & Shetty, M. (2004). *J. Phys. Org. Chem.* **17**, 848–864.
- Gowda, B. T., Svoboda, I. & Fuess, H. (2000). *Z. Naturforsch. Teil A*, **55**, 779–790.
- Macrae, C. F., Bruno, I. J., Chisholm, J. A., Edgington, P. R., McCabe, P., Pidcock, E., Rodriguez-Monge, L., Taylor, R., van de Streek, J. & Wood, P. A. (2008). *J. Appl. Cryst.* **41**, 466–470.
- Oxford Diffraction (2009). *CrysAlis CCD* and *CrysAlis RED*. Oxford Diffraction Ltd, Yarnton, England.
- Saeed, A., Arshad, M. & Simpson, J. (2010). *Acta Cryst.* **E66**, o2808–o2809.
- Sheldrick, G. M. (2008). *Acta Cryst.* **A64**, 112–122.
- Shetty, M. & Gowda, B. T. (2005). *Z. Naturforsch. Teil A*, **60**, 113–120.

supplementary materials

Acta Cryst. (2011). E67, o2903 [doi:10.1107/S1600536811040864]

4-Chloro-*N*-(3-methylphenyl)benzamide

V. Z. Rodrigues, M. Fronc, B. T. Gowda and J. Kozisek

Comment

The amide and sulfonamide moieties are the constituents of many biologically important compounds. As part of our studies on the substituent effects on the structures and other aspects of *N*-(aryl)-amides (Bowes *et al.*, 2003; Gowda *et al.*, 2000; Saeed *et al.*, 2010), *N*-(aryl)-methanesulfonamides (Gowda *et al.*, 2007), *N*-(aryl)-arylsulfonamides (Shetty & Gowda, 2005) and *N*-chloro-arylsulfonamides (Gowda & Shetty, 2004), in the present work, the crystal structure of 4-chloro-*N*-(3-methylphenyl)benzamide (I) has been determined (Fig. 1). In (I), the *meta*-methyl substituent in the anilino ring is positioned *anti* to the N–H bond.

The central amide group –NHCO– is tilted to the anilino ring with the C9–C8–N1–C7 and C13–C8–N1–C7 torsion angles of 30.4 (3)° and -151.2 (2)°. The C2–C1–C7–N1 and C6–C1–C7–N1 torsion angles are -15.7 (3)° and 166.8 (2)°, respectively, while the C2–C1–C7–O1 and C6–C1–C7–O1 torsion angles are 162.8 (2)° and -14.7 (3)°, respectively. But the C1–C7–N1–C8 and C8–N1–C7–O1 torsion angles are 173.6 (2)° and -4.9 (3)°, respectively.

The packing of molecules linked by N–H···O hydrogen bonds is shown in Fig. 2.

Experimental

The title compound was prepared according to the method described by Gowda *et al.* (2003). The purity of the compound was checked by determining its melting point. It was characterized by recording its infrared and NMR spectra. Plate like colourless single crystals of the title compound were obtained by slow evaporation of an ethanol solution of the compound (0.5 g in about 30 ml of ethanol) at room temperature.

Refinement

All H atoms were visible in difference maps and then treated as riding atoms with C–H distances of 0.93 Å (C-aromatic), 0.96 Å (C-methyl) and N–H = 0.86 Å. The $U_{\text{iso}}(\text{H})$ values were set at 1.2 $U_{\text{eq}}(\text{C-aromatic, N})$ and 1.5 $U_{\text{eq}}(\text{C-methyl})$.

Figures




Fig. 1. Molecular structure of the title compound showing the atom labelling scheme. Displacement ellipsoids are drawn at the 50% probability level. H atoms are represented as small spheres of arbitrary radii.




Fig. 2. Part of the crystal structure of the title compound generated by N—H...O hydrogen bonds which are shown by dashed lines. H atoms not involved in intermolecular bonding have been omitted.

4-Chloro-*N*-(3-methylphenyl)benzamide

Crystal data

$C_{14}H_{12}ClNO$

$M_r = 245.70$

Monoclinic, $P2_1/c$

Hall symbol: -p 2ybc

$a = 13.4379$ (9) Å

$b = 10.2493$ (11) Å

$c = 9.2600$ (7) Å

$\beta = 92.893$ (6)°

$V = 1273.74$ (19) Å³

$Z = 4$

$F(000) = 512$

$D_x = 1.281$ Mg m⁻³

Mo $K\alpha$ radiation, $\lambda = 0.71073$ Å

Cell parameters from 4939 reflections

$\theta = 3.4$ – 29.4 °

$\mu = 0.28$ mm⁻¹

$T = 293$ K

Plate, colorless

$0.81 \times 0.17 \times 0.04$ mm

Data collection

Oxford Diffraction Xcalibur
diffractometer with a Ruby (Gemini Cu) detector

Radiation source: fine-focus sealed tube
graphite

ω scans

Absorption correction: analytical

[*CrysAlis RED* (Oxford Diffraction, 2009), based on
expressions derived from Clark & Reid (1995)]

$T_{\min} = 0.877$, $T_{\max} = 0.988$

17187 measured reflections

2156 independent reflections

1299 reflections with $I > 2\sigma(I)$

$R_{\text{int}} = 0.040$

$\theta_{\max} = 24.7$ °, $\theta_{\min} = 4.2$ °

$h = -15 \rightarrow 15$

$k = -12 \rightarrow 12$

$l = -10 \rightarrow 10$

Refinement

Refinement on F^2

Least-squares matrix: full

$R[F^2 > 2\sigma(F^2)] = 0.040$

$wR(F^2) = 0.117$

$S = 0.94$

Primary atom site location: structure-invariant direct
methods

Secondary atom site location: difference Fourier map

Hydrogen site location: inferred from neighbouring
sites

H-atom parameters constrained

$w = 1/[\sigma^2(F_o^2) + (0.0699P)^2]$

2156 reflections

154 parameters

0 restraints

$$\text{where } P = (F_o^2 + 2F_c^2)/3$$

$$(\Delta/\sigma)_{\max} < 0.001$$

$$\Delta\rho_{\max} = 0.22 \text{ e } \text{\AA}^{-3}$$

$$\Delta\rho_{\min} = -0.13 \text{ e } \text{\AA}^{-3}$$

Special details

Experimental. CrysAlis RED (Oxford Diffraction, 2009) Analytical numeric absorption correction using a multifaceted crystal model based on expressions derived (Clark & Reid, 1995).

Geometry. All e.s.d.'s (except the e.s.d. in the dihedral angle between two l.s. planes) are estimated using the full covariance matrix. The cell e.s.d.'s are taken into account individually in the estimation of e.s.d.'s in distances, angles and torsion angles; correlations between e.s.d.'s in cell parameters are only used when they are defined by crystal symmetry. An approximate (isotropic) treatment of cell e.s.d.'s is used for estimating e.s.d.'s involving l.s. planes.

Refinement. Refinement of F^2 against ALL reflections. The weighted R -factor wR and goodness of fit S are based on F^2 , conventional R -factors R are based on F , with F set to zero for negative F^2 . The threshold expression of $F^2 > \sigma(F^2)$ is used only for calculating R -factors(gt) *etc.* and is not relevant to the choice of reflections for refinement. R -factors based on F^2 are statistically about twice as large as those based on F , and R -factors based on ALL data will be even larger.

Fractional atomic coordinates and isotropic or equivalent isotropic displacement parameters (\AA^2)

	<i>x</i>	<i>y</i>	<i>z</i>	$U_{\text{iso}}^*/U_{\text{eq}}$
C1	0.44789 (14)	0.3375 (2)	-0.05692 (18)	0.0570 (5)
C2	0.40134 (16)	0.2716 (2)	0.0505 (2)	0.0749 (6)
H2A	0.4350	0.2045	0.0999	0.090*
C3	0.30600 (17)	0.3031 (3)	0.0860 (3)	0.0850 (7)
H3A	0.2759	0.2590	0.1600	0.102*
C4	0.25583 (15)	0.4008 (3)	0.0105 (3)	0.0746 (6)
C5	0.30042 (16)	0.4691 (2)	-0.0950 (2)	0.0715 (6)
H5A	0.2665	0.5361	-0.1441	0.086*
C6	0.39659 (15)	0.4374 (2)	-0.1280 (2)	0.0651 (6)
H6A	0.4274	0.4841	-0.1994	0.078*
C7	0.54954 (14)	0.3041 (2)	-0.10552 (19)	0.0587 (5)
C8	0.70086 (15)	0.1721 (2)	-0.0475 (2)	0.0636 (6)
C9	0.76789 (15)	0.2328 (2)	-0.1344 (2)	0.0695 (6)
H9A	0.7512	0.3126	-0.1768	0.083*
C10	0.85939 (16)	0.1769 (3)	-0.1594 (2)	0.0806 (7)
C11	0.88194 (19)	0.0584 (3)	-0.0961 (3)	0.0932 (8)
H11A	0.9423	0.0185	-0.1139	0.112*
C12	0.8172 (2)	-0.0028 (3)	-0.0066 (3)	0.0966 (8)
H12A	0.8347	-0.0820	0.0366	0.116*
C13	0.72518 (18)	0.0546 (3)	0.0188 (2)	0.0811 (7)
H13A	0.6812	0.0144	0.0793	0.097*
C14	0.93185 (19)	0.2429 (3)	-0.2533 (3)	0.1117 (10)
H14C	0.9906	0.1900	-0.2584	0.134*
H14B	0.9496	0.3265	-0.2129	0.134*
H14A	0.9018	0.2544	-0.3487	0.134*
N1	0.60639 (12)	0.22675 (17)	-0.01846 (16)	0.0636 (5)

supplementary materials

H1A	0.5833	0.2081	0.0640	0.076*
O1	0.57677 (10)	0.34502 (16)	-0.22211 (13)	0.0762 (5)
C11	0.13343 (4)	0.43547 (8)	0.04942 (9)	0.1151 (4)

Atomic displacement parameters (\AA^2)

	U^{11}	U^{22}	U^{33}	U^{12}	U^{13}	U^{23}
C1	0.0537 (11)	0.0700 (14)	0.0472 (10)	-0.0039 (11)	0.0032 (8)	-0.0056 (10)
C2	0.0575 (13)	0.0910 (17)	0.0767 (14)	0.0007 (11)	0.0087 (11)	0.0183 (12)
C3	0.0605 (14)	0.108 (2)	0.0887 (16)	-0.0038 (14)	0.0215 (12)	0.0212 (14)
C4	0.0506 (12)	0.0875 (17)	0.0864 (14)	-0.0042 (11)	0.0101 (11)	-0.0100 (13)
C5	0.0633 (13)	0.0758 (16)	0.0751 (13)	0.0058 (11)	0.0019 (11)	-0.0011 (11)
C6	0.0650 (13)	0.0747 (15)	0.0562 (11)	-0.0025 (11)	0.0096 (10)	-0.0003 (10)
C7	0.0562 (12)	0.0761 (14)	0.0438 (10)	-0.0039 (10)	0.0031 (9)	-0.0071 (9)
C8	0.0598 (12)	0.0813 (16)	0.0498 (10)	0.0084 (11)	0.0022 (9)	-0.0116 (11)
C9	0.0598 (13)	0.0931 (17)	0.0558 (11)	0.0087 (12)	0.0043 (10)	-0.0080 (10)
C10	0.0590 (14)	0.123 (2)	0.0590 (12)	0.0198 (14)	-0.0026 (10)	-0.0145 (13)
C11	0.0670 (15)	0.131 (3)	0.0807 (15)	0.0254 (16)	-0.0042 (13)	-0.0256 (17)
C12	0.091 (2)	0.095 (2)	0.1014 (19)	0.0246 (16)	-0.0190 (16)	-0.0062 (15)
C13	0.0794 (16)	0.0897 (19)	0.0739 (14)	0.0095 (14)	0.0015 (12)	-0.0021 (13)
C14	0.0646 (15)	0.179 (3)	0.0930 (17)	0.0155 (17)	0.0231 (13)	-0.0068 (17)
N1	0.0575 (10)	0.0846 (13)	0.0494 (9)	0.0063 (9)	0.0104 (7)	-0.0001 (8)
O1	0.0624 (8)	0.1183 (13)	0.0485 (8)	0.0053 (8)	0.0093 (6)	0.0064 (8)
Cl1	0.0586 (4)	0.1374 (7)	0.1517 (7)	0.0078 (4)	0.0288 (4)	0.0031 (5)

Geometric parameters (\AA , $^\circ$)

C1—C2	1.378 (3)	C8—C9	1.386 (3)
C1—C6	1.382 (3)	C8—N1	1.426 (3)
C1—C7	1.499 (3)	C9—C10	1.387 (3)
C2—C3	1.377 (3)	C9—H9A	0.9300
C2—H2A	0.9300	C10—C11	1.375 (4)
C3—C4	1.378 (3)	C10—C14	1.499 (4)
C3—H3A	0.9300	C11—C12	1.382 (4)
C4—C5	1.365 (3)	C11—H11A	0.9300
C4—C11	1.738 (2)	C12—C13	1.400 (4)
C5—C6	1.381 (3)	C12—H12A	0.9300
C5—H5A	0.9300	C13—H13A	0.9300
C6—H6A	0.9300	C14—H14C	0.9600
C7—O1	1.231 (2)	C14—H14B	0.9600
C7—N1	1.342 (2)	C14—H14A	0.9600
C8—C13	1.383 (3)	N1—H1A	0.8600
C2—C1—C6	118.24 (19)	C8—C9—C10	121.3 (2)
C2—C1—C7	123.92 (19)	C8—C9—H9A	119.4
C6—C1—C7	117.79 (17)	C10—C9—H9A	119.4
C3—C2—C1	121.3 (2)	C11—C10—C9	118.2 (2)
C3—C2—H2A	119.4	C11—C10—C14	120.7 (2)
C1—C2—H2A	119.4	C9—C10—C14	121.2 (3)

C2—C3—C4	119.1 (2)	C10—C11—C12	121.6 (2)
C2—C3—H3A	120.4	C10—C11—H11A	119.2
C4—C3—H3A	120.4	C12—C11—H11A	119.2
C5—C4—C3	121.0 (2)	C11—C12—C13	119.9 (3)
C5—C4—C11	119.86 (19)	C11—C12—H12A	120.0
C3—C4—C11	119.17 (18)	C13—C12—H12A	120.0
C4—C5—C6	119.1 (2)	C8—C13—C12	118.8 (2)
C4—C5—H5A	120.4	C8—C13—H13A	120.6
C6—C5—H5A	120.4	C12—C13—H13A	120.6
C5—C6—C1	121.23 (19)	C10—C14—H14C	109.5
C5—C6—H6A	119.4	C10—C14—H14B	109.5
C1—C6—H6A	119.4	H14C—C14—H14B	109.5
O1—C7—N1	122.88 (18)	C10—C14—H14A	109.5
O1—C7—C1	120.08 (17)	H14C—C14—H14A	109.5
N1—C7—C1	117.03 (16)	H14B—C14—H14A	109.5
C13—C8—C9	120.2 (2)	C7—N1—C8	127.20 (16)
C13—C8—N1	116.8 (2)	C7—N1—H1A	116.4
C9—C8—N1	123.0 (2)	C8—N1—H1A	116.4
C6—C1—C2—C3	0.4 (3)	C13—C8—C9—C10	1.3 (3)
C7—C1—C2—C3	-177.07 (19)	N1—C8—C9—C10	179.60 (17)
C1—C2—C3—C4	1.3 (4)	C8—C9—C10—C11	0.4 (3)
C2—C3—C4—C5	-2.2 (4)	C8—C9—C10—C14	-179.9 (2)
C2—C3—C4—C11	176.97 (18)	C9—C10—C11—C12	-1.7 (3)
C3—C4—C5—C6	1.3 (3)	C14—C10—C11—C12	178.6 (2)
C11—C4—C5—C6	-177.83 (15)	C10—C11—C12—C13	1.4 (4)
C4—C5—C6—C1	0.5 (3)	C9—C8—C13—C12	-1.6 (3)
C2—C1—C6—C5	-1.3 (3)	N1—C8—C13—C12	179.96 (18)
C7—C1—C6—C5	176.33 (17)	C11—C12—C13—C8	0.3 (3)
C2—C1—C7—O1	162.80 (19)	O1—C7—N1—C8	-4.9 (3)
C6—C1—C7—O1	-14.7 (3)	C1—C7—N1—C8	173.59 (17)
C2—C1—C7—N1	-15.7 (3)	C13—C8—N1—C7	-151.2 (2)
C6—C1—C7—N1	166.77 (17)	C9—C8—N1—C7	30.4 (3)

Hydrogen-bond geometry (\AA , $^\circ$)

$D-H\cdots A$	$D-H$	$H\cdots A$	$D\cdots A$	$D-H\cdots A$
N1—H1A \cdots O1 ⁱ	0.86	2.06	2.888 (2)	161.

Symmetry codes: (i) $x, -y+1/2, z+1/2$.

Fig. 1




Fig. 2

