

2,3-Dichloro-1,4-hydroquinone 2,3-dichloro-1,4-benzoquinone monohydrate: a quinhydrone-type 1:1 donor-acceptor [D—A] charge-transfer complex

Xavier Guégano, Jürg Hauser, Shi-Xia Liu* and Silvio Decurtins

Departement für Chemie und Biochemie, Universität Bern, Freiestrasse 3, CH-3012 Bern, Switzerland

Correspondence e-mail: liu@iac.unibe.ch


Received 27 September 2011; accepted 7 October 2011

Key indicators: single-crystal X-ray study; $T = 173$ K; mean $\sigma(\text{C}-\text{C}) = 0.003$ Å; R factor = 0.031; wR factor = 0.077; data-to-parameter ratio = 15.8.

In the crystal structure of the title compound (systematic name: 2,3-dichlorobenzene-1,4-diol 2,3-dichlorocyclohexa-2,5-diene-1,4-dione monohydrate), $\text{C}_6\text{H}_4\text{Cl}_2\text{O}_2 \cdot \text{C}_6\text{H}_2\text{Cl}_2\text{O}_2 \cdot \text{H}_2\text{O}$, the 2,3-dichloro-1,4-hydroquinone donor (*D*) and the 2,3-dichloro-1,4-benzoquinone acceptor (*A*) molecules form alternating stacks along [100]. Their molecular planes [maximum deviations for non-H atoms: 0.0133 (14) (*D*) and 0.0763 (14) Å (*A*)] are inclined to one another by 1.45 (3)° and are thus almost parallel. There are π - π interactions involving the *D* and *A* molecules, with centroid-centroid distances of 3.5043 (9) and 3.9548 (9) Å. Intermolecular O—H...O hydrogen bonds involving the water molecule and the hydroxy and ketone groups lead to the formation of two-dimensional networks lying parallel to (001). These networks are linked by C—H...O interactions, forming a three-dimensional structure.

Related literature

For prototypical examples of similar organic redox systems, see: Yi *et al.* (2009*a,b*). For details concerning quinhydrone, a 1:1 hydroquinone-quinone adduct, and a well known molecular charge-transfer (CT) complex, see: Foster (1969). For structural studies of different polymorphs of quinhydrone, see: Matsuda *et al.* (1958); Sakurai (1965,1968). For details concerning quinhydrone analogues, see: Bouvet *et al.* (2006,2007); Patil *et al.* (1984); Yamamura *et al.* (2007). For a detailed computational study on the stacking energies and the electron density topology in quinhydrone, see: Gonzalez Moa *et al.* (2007).



Experimental

Crystal data

$\text{C}_6\text{H}_4\text{Cl}_2\text{O}_2 \cdot \text{C}_6\text{H}_2\text{Cl}_2\text{O}_2 \cdot \text{H}_2\text{O}$
 $M_r = 373.98$
 Monoclinic, $P2_1/c$
 $a = 7.15329$ (14) Å
 $b = 7.19541$ (15) Å
 $c = 27.2811$ (5) Å
 $\beta = 92.9738$ (18)°

$V = 1402.29$ (5) Å³
 $Z = 4$
 Mo $K\alpha$ radiation
 $\mu = 0.86$ mm⁻¹
 $T = 173$ K
 $0.3 \times 0.2 \times 0.07$ mm

Data collection

Siemens SMART 1K CCD area-detector diffractometer
 Absorption correction: multi-scan (SADABS; Sheldrick, 1996)
 $T_{\min} = 0.859$, $T_{\max} = 0.942$

19255 measured reflections
 3138 independent reflections
 2653 reflections with $I > 2\sigma(I)$
 $R_{\text{int}} = 0.026$

Refinement

$R[F^2 > 2\sigma(F^2)] = 0.031$
 $wR(F^2) = 0.077$
 $S = 1.04$
 3138 reflections
 198 parameters

H atoms treated by a mixture of independent and constrained refinement
 $\Delta\rho_{\max} = 0.49$ e Å⁻³
 $\Delta\rho_{\min} = -0.19$ e Å⁻³

Table 1

Hydrogen-bond geometry (Å, °).

<i>D</i> —H... <i>A</i>	<i>D</i> —H	H... <i>A</i>	<i>D</i> ... <i>A</i>	<i>D</i> —H... <i>A</i>
O11—H11...O20	0.84	1.76	2.5947 (18)	173
O14—H14...O4 ⁱ	0.84	2.03	2.8381 (18)	161
O20—H20A...O1 ⁱⁱ	0.79 (3)	2.12 (3)	2.914 (2)	177 (3)
O20—H20B...O11 ⁱⁱⁱ	0.74 (3)	2.06 (3)	2.7899 (19)	169 (3)
C6—H6...O14 ^{iv}	0.95	2.48	3.209 (2)	134

Symmetry codes: (i) $x - 1, y + 1, z$; (ii) $-x + 1, y - \frac{1}{2}, -z + \frac{1}{2}$; (iii) $-x + 1, y + \frac{1}{2}, -z + \frac{1}{2}$; (iv) $-x + 1, -y + 2, -z + 1$.

Data collection: SMART (Bruker, 2003); cell refinement: SAINT (Bruker, 2003); data reduction: SAINT; program(s) used to solve structure: SHELXS97 (Sheldrick, 2008); program(s) used to refine structure: SHELXL97 (Sheldrick, 2008); molecular graphics: Mercury (Macrae *et al.*, 2008); software used to prepare material for publication: publCIF (Westrip, 2010).

The X-ray diffraction analysis was possible thanks to the Swiss National Science Foundation (R'Equip project 206021–128724).

Supplementary data and figures for this paper are available from the IUCr electronic archives (Reference: SU2322).

References

- Bouvet, M., Malézieux, B. & Herson, P. (2006). *Chem. Commun.* pp. 1751–1753.
- Bouvet, M., Malézieux, B., Herson, P. & Villain, F. (2007). *CrystEngComm*, **9**, 270–272.
- Bruker (2003). *SMART* and *SAINT*. Bruker AXS Inc., Madison, Wisconsin, USA.
- Foster, F. (1969). In *Organic Charge-Transfer Complexes*. Academic Press: New York.
- Gonzalez Moa, M. J., Mandado, M. & Mosquera, R. A. (2007). *J. Phys. Chem. A*, **111**, 1998–2001.
- Macrae, C. F., Bruno, I. J., Chisholm, J. A., Edgington, P. R., McCabe, P., Pidcock, E., Rodriguez-Monge, L., Taylor, R., van de Streek, J. & Wood, P. A. (2008). *J. Appl. Cryst.* **41**, 466–470.
- Matsuda, H., Osaki, K. & Nitta, I. (1958). *Bull. Chem. Soc. Jpn*, **31**, 611–620.
- Patil, A. O., Curtin, D. Y. & Paul, I. C. (1984). *J. Am. Chem. Soc.* **106**, 4010–4015.
- Sakurai, T. (1965). *Acta Cryst.* **19**, 320–330.
- Sakurai, T. (1968). *Acta Cryst.* **B24**, 403–412.
- Sheldrick, G. M. (1996). *SADABS*. University of Göttingen, Germany.
- Sheldrick, G. M. (2008). *Acta Cryst.* **A64**, 112–122.
- Westrip, S. P. (2010). *J. Appl. Cryst.* **43**, 920–925.
- Yamamura, K., Yamane, J., Eda, K., Tajima, F., Yamada, Y. & Hashimoto, M. (2007). *J. Mol. Struct.* **842**, 12–16.
- Yi, C., Blum, C., Liu, S.-X., Keene, T. D., Frei, G., Neels, A. & Decurtins, S. (2009a). *Org. Lett.* **11**, 2261–2264.
- Yi, C., Liu, S.-X., Neels, A., Renaud, P. & Decurtins, S. (2009b). *Org. Lett.* **11**, 5530–5533.

supplementary materials

Acta Cryst. (2011). E67, o2967-o2968 [doi:10.1107/S1600536811041377]

2,3-Dichloro-1,4-hydroquinone 2,3-dichloro-1,4-benzoquinone monohydrate: a quinhydrone-type 1:1 donor-acceptor [D-A] charge-transfer complex

X. Guégano, J. Hauser, S.-X. Liu and S. Decurtins

Comment

Quinones as electron acceptors (A) and their reduced forms, semiquinones and dihydroquinones as electron donors (D) represent prototypical examples of organic redox systems (Yi *et al.*, 2009*a,b*). Furthermore, it is certainly true that interactions between aromatic π -systems have long been observed in crystal structures and rationales for the occurrence of π - π interactions have included explanations which are based on electron donor-acceptor [D—A] models. Such charge-transfer (CT) complexes may form when good electron donors and acceptors lie in close proximity, and this situation is typically associated with intense electronic CT transitions in the UV-vis spectral region. Basically, the filled donor molecular orbital (HOMO) and the vacant acceptor molecular orbital (LUMO) maximize their overlap. Quinhydrone, a 1:1 hydroquinone-quinone adduct, is a well known molecular CT complex (Foster, 1969) and its complex stabilization is based on the CT between the electron donor (hydroquinone) and the electron acceptor (quinone). Structural studies of different polymorphs of quinhydrone have been undertaken (Matsuda *et al.*, 1958; Sakurai, 1965, 1968). Quinhydrone analogues have also been studied (Bouvet *et al.*, 2006, 2007, Patil *et al.*, 1984 and Yamamura *et al.*, 2007). A detailed computational study on the stacking energies and the electron density topology in quinhydrone has recently been given (Gonzalez Moa *et al.*, 2007). Additionally, hydrogen bonds contribute further stability, both in the solid state as well as in solution. We report herein on the crystal structure of the title quinhydrone analogue.

The molecular structure of the title compound is illustrated in Fig. 1. The 2,3-Dichloro-1,4-hydroquinone donor (D) and the 2,3-Dichloro-1,4-benzoquinone acceptor (A) molecules form alternate stacks along the [100] direction with their molecular planes [max. deviations for non-H atoms: (D) = 0.0133 (14) Å and (A) = 0.0763 (14) Å] are tilted by ca. 22.8° and 23.1° about the [100] direction. Their molecular planes are inclined to one another by 1.45 (3)° and are thus almost parallel (Fig. 2). The π - π interactions involving the six-membered rings of the (D) [C11-C16] and (A) [C1-C6] molecules have centroid-centroid distances of 3.5043 (9) Å [D \cdots Aⁱ; symmetry code (i) x-1, y, z] and 3.9548 (9) Å [D \cdots A].

Intermolecular O—H \cdots O hydrogen bonds between water and the hydroxyl and ketone groups of adjacent stacks form a complex two-dimensional network lying parallel to (001) [Fig. 3]. Chlorine atoms, Cl2 and Cl3ⁱ, of (A) in adjacent stacks are in short contact with a distance of 3.3902 (7) Å [symmetry code: (i) = 2-x, 0.5+y, 0.5-z]. The two-dimensional networks are linked via C-H \cdots O contacts to form a three-dimensional structure.

Experimental

Deep-red coloured, elongated (up to 5 mm long) plate-shaped crystals of the 1:1 donor-acceptor adduct [D—A].H₂O (D = 2,3-Dichloro-1,4-hydroquinone; A = 2,3-Dichloro-1,4-benzoquinone) were formed by slow sublimation of the single component (A) within a closed flask at room-temperature under aerobic conditions.

Refinement

H atoms of the water molecule were located in a difference Fourier map and refined with $U_{\text{iso}}(\text{H}) = 1.5$ times $U_{\text{eq}}(\text{O})$. Other H atoms were positioned geometrically and refined using a riding model (including free rotation about the hydroxy C—O bond): O—H = 0.84 Å, C—H = 0.95 Å, with $U_{\text{iso}}(\text{H}) = k \times U_{\text{eq}}(\text{O,C})$, where $k = 1.5$ for OH H atoms, and $k = 1.2$ for all other H atoms.

Figures




Fig. 1. The molecular structure of the title compound, with atom labels and displacement ellipsoids drawn at the 50% probability level [left donor (D); right acceptor (A)].




Fig. 2. A view perpendicular to the molecular planes, of the stacking of the acceptor (A) and donor (D) molecules in the title compound.




Fig. 3. The crystal packing of the title compound, viewed along the *a* axis. Stacks of donor (D) and acceptor (A) molecules are connected by O—H...O hydrogen bonds (dashed blue lines).

2,3-dichlorobenzene-1,4-diol 2,3-dichlorocyclohexa-2,5-diene-1,4-dione monohydrate

Crystal data

$\text{C}_6\text{H}_4\text{Cl}_2\text{O}_2 \cdot \text{C}_6\text{H}_2\text{Cl}_2\text{O}_2 \cdot \text{H}_2\text{O}$

$M_r = 373.98$

Monoclinic, $P2_1/c$

$F(000) = 752$

$D_x = 1.771 \text{ Mg m}^{-3}$

Mo $K\alpha$ radiation, $\lambda = 0.71073 \text{ \AA}$

Hall symbol: -P 2ybc
 $a = 7.15329$ (14) Å
 $b = 7.19541$ (15) Å
 $c = 27.2811$ (5) Å
 $\beta = 92.9738$ (18)°
 $V = 1402.29$ (5) Å³
 $Z = 4$

Cell parameters from 7958 reflections
 $\theta = 2.8$ – 27.5 °
 $\mu = 0.86$ mm⁻¹
 $T = 173$ K
 Plate, red
 $0.3 \times 0.2 \times 0.07$ mm

Data collection

Siemens SMART 1K CCD area-detector diffractometer
 Radiation source: fine-focus sealed tube graphite
 rotation method scans
 Absorption correction: multi-scan (SADABS; Sheldrick, 1996)
 $T_{\min} = 0.859$, $T_{\max} = 0.942$
 19255 measured reflections

3138 independent reflections
 2653 reflections with $I > 2\sigma(I)$
 $R_{\text{int}} = 0.026$
 $\theta_{\max} = 27.9$ °, $\theta_{\min} = 2.9$ °
 $h = -8$ → 9
 $k = -9$ → 9
 $l = -35$ → 35

Refinement

Refinement on F^2
 Least-squares matrix: full
 $R[F^2 > 2\sigma(F^2)] = 0.031$
 $wR(F^2) = 0.077$
 $S = 1.04$
 3138 reflections
 198 parameters
 0 restraints

Primary atom site location: structure-invariant direct methods
 Secondary atom site location: difference Fourier map
 Hydrogen site location: inferred from neighbouring sites
 H atoms treated by a mixture of independent and constrained refinement
 $w = 1/[\sigma^2(F_o^2) + (0.0429P)^2 + 0.583P]$
 where $P = (F_o^2 + 2F_c^2)/3$
 $(\Delta/\sigma)_{\max} = 0.001$
 $\Delta\rho_{\max} = 0.49$ e Å⁻³
 $\Delta\rho_{\min} = -0.19$ e Å⁻³

Special details

Geometry. All s.u.'s (except the s.u. in the dihedral angle between two l.s. planes) are estimated using the full covariance matrix. The cell s.u.'s are taken into account individually in the estimation of s.u.'s in distances, angles and torsion angles; correlations between s.u.'s in cell parameters are only used when they are defined by crystal symmetry. An approximate (isotropic) treatment of cell s.u.'s is used for estimating s.u.'s involving l.s. planes.

Refinement. Refinement of F^2 against ALL reflections. The weighted R -factor wR and goodness of fit S are based on F^2 , conventional R -factors R are based on F , with F set to zero for negative F^2 . The threshold expression of $F^2 > \sigma(F^2)$ is used only for calculating R -factors(gt) *etc.* and is not relevant to the choice of reflections for refinement. R -factors based on F^2 are statistically about twice as large as those based on F , and R -factors based on ALL data will be even larger.

supplementary materials

Fractional atomic coordinates and isotropic or equivalent isotropic displacement parameters (\AA^2)

	x	y	z	$U_{\text{iso}}^*/U_{\text{eq}}$
C1	0.7866 (2)	0.9165 (2)	0.38598 (7)	0.0220 (4)
O1	0.72734 (19)	1.07053 (18)	0.37504 (5)	0.0299 (3)
C2	0.8432 (2)	0.7822 (3)	0.34779 (6)	0.0216 (4)
Cl2	0.81893 (7)	0.85660 (7)	0.288391 (16)	0.03021 (13)
C3	0.9107 (2)	0.6140 (3)	0.36016 (6)	0.0220 (4)
Cl3	0.97542 (7)	0.45526 (7)	0.317735 (18)	0.03252 (13)
C4	0.9362 (2)	0.5565 (2)	0.41269 (7)	0.0229 (4)
O4	1.00840 (19)	0.40868 (18)	0.42426 (5)	0.0305 (3)
C5	0.8731 (2)	0.6879 (3)	0.44963 (7)	0.0252 (4)
H5	0.8808	0.6525	0.4832	0.030*
C6	0.8052 (3)	0.8555 (3)	0.43743 (7)	0.0251 (4)
H6	0.7682	0.9370	0.4625	0.030*
C11	0.3767 (2)	0.7229 (2)	0.33596 (6)	0.0194 (3)
O11	0.43186 (18)	0.60154 (17)	0.30131 (4)	0.0245 (3)
H11	0.4304	0.6550	0.2739	0.037*
C12	0.3915 (2)	0.6700 (2)	0.38514 (6)	0.0185 (3)
Cl12	0.47890 (6)	0.45165 (6)	0.399712 (15)	0.02360 (11)
C13	0.3366 (2)	0.7902 (2)	0.42204 (6)	0.0196 (3)
Cl13	0.35592 (7)	0.72408 (6)	0.482946 (16)	0.02805 (12)
C14	0.2652 (2)	0.9655 (2)	0.40960 (6)	0.0206 (4)
O14	0.2109 (2)	1.07750 (18)	0.44656 (5)	0.0289 (3)
H14	0.1681	1.1775	0.4347	0.043*
C15	0.2513 (2)	1.0172 (2)	0.36060 (6)	0.0220 (4)
H15	0.2035	1.1366	0.3519	0.026*
C16	0.3058 (2)	0.8979 (2)	0.32418 (6)	0.0213 (4)
H16	0.2947	0.9361	0.2908	0.026*
O20	0.4387 (2)	0.7413 (2)	0.21375 (5)	0.0325 (3)
H20A	0.393 (3)	0.691 (4)	0.1903 (10)	0.049*
H20B	0.472 (4)	0.834 (4)	0.2061 (10)	0.049*

Atomic displacement parameters (\AA^2)

	U^{11}	U^{22}	U^{33}	U^{12}	U^{13}	U^{23}
C1	0.0203 (8)	0.0230 (9)	0.0224 (9)	-0.0010 (7)	-0.0011 (7)	0.0017 (7)
O1	0.0371 (7)	0.0239 (7)	0.0283 (7)	0.0054 (6)	-0.0032 (6)	0.0030 (5)
C2	0.0203 (8)	0.0282 (9)	0.0160 (8)	-0.0027 (7)	-0.0002 (7)	0.0021 (7)
Cl2	0.0329 (2)	0.0399 (3)	0.0178 (2)	0.0027 (2)	0.00070 (17)	0.00536 (19)
C3	0.0190 (8)	0.0246 (9)	0.0226 (9)	-0.0017 (7)	0.0023 (7)	-0.0043 (7)
Cl3	0.0360 (3)	0.0321 (3)	0.0298 (3)	0.0031 (2)	0.00496 (19)	-0.00926 (19)
C4	0.0195 (8)	0.0235 (9)	0.0254 (9)	-0.0020 (7)	-0.0005 (7)	0.0027 (7)
O4	0.0315 (7)	0.0243 (7)	0.0356 (8)	0.0055 (6)	-0.0008 (6)	0.0056 (6)
C5	0.0266 (9)	0.0297 (9)	0.0193 (9)	0.0005 (8)	0.0012 (7)	0.0028 (7)
C6	0.0267 (9)	0.0281 (9)	0.0206 (9)	0.0021 (8)	0.0022 (7)	-0.0022 (7)
C11	0.0206 (8)	0.0185 (8)	0.0190 (8)	-0.0021 (7)	0.0013 (6)	-0.0019 (6)

O11	0.0373 (7)	0.0196 (6)	0.0169 (6)	0.0037 (5)	0.0047 (5)	0.0002 (5)
C12	0.0188 (8)	0.0144 (7)	0.0223 (9)	0.0003 (6)	-0.0001 (6)	0.0023 (6)
Cl12	0.0305 (2)	0.0166 (2)	0.0236 (2)	0.00314 (17)	0.00040 (17)	0.00279 (16)
C13	0.0214 (8)	0.0210 (8)	0.0163 (8)	-0.0018 (7)	0.0002 (6)	0.0022 (7)
Cl13	0.0374 (3)	0.0288 (2)	0.0179 (2)	0.0034 (2)	0.00093 (17)	0.00261 (17)
C14	0.0219 (8)	0.0178 (8)	0.0221 (9)	-0.0007 (7)	0.0021 (7)	-0.0025 (7)
O14	0.0412 (8)	0.0213 (6)	0.0245 (7)	0.0081 (6)	0.0043 (6)	-0.0023 (5)
C15	0.0241 (9)	0.0161 (8)	0.0257 (9)	0.0005 (7)	-0.0009 (7)	0.0034 (7)
C16	0.0232 (9)	0.0220 (8)	0.0186 (9)	-0.0001 (7)	0.0004 (7)	0.0041 (7)
O20	0.0539 (9)	0.0238 (7)	0.0192 (7)	-0.0103 (7)	-0.0027 (6)	0.0009 (5)

Geometric parameters (Å, °)

C1—O1	1.218 (2)	C11—C12	1.393 (2)
C1—C6	1.470 (3)	O11—H11	0.8400
C1—C2	1.493 (2)	C12—C13	1.400 (2)
C2—C3	1.340 (3)	C12—Cl12	1.7292 (16)
C2—Cl2	1.7070 (17)	C13—C14	1.396 (2)
C3—C4	1.494 (2)	C13—Cl13	1.7270 (17)
C3—Cl3	1.7069 (18)	C14—O14	1.363 (2)
C4—O4	1.217 (2)	C14—C15	1.386 (2)
C4—C5	1.470 (3)	O14—H14	0.8400
C5—C6	1.335 (3)	C15—C16	1.384 (2)
C5—H5	0.9500	C15—H15	0.9500
C6—H6	0.9500	C16—H16	0.9500
C11—O11	1.360 (2)	O20—H20A	0.79 (3)
C11—C16	1.389 (2)	O20—H20B	0.74 (3)
O1—C1—C6	121.26 (17)	C16—C11—C12	118.64 (16)
O1—C1—C2	121.44 (16)	C11—O11—H11	109.5
C6—C1—C2	117.30 (15)	C11—C12—C13	120.89 (15)
C3—C2—C1	121.09 (15)	C11—C12—Cl12	118.58 (13)
C3—C2—Cl2	122.70 (14)	C13—C12—Cl12	120.53 (13)
C1—C2—Cl2	116.21 (13)	C14—C13—C12	119.78 (15)
C2—C3—C4	121.08 (16)	C14—C13—Cl13	119.55 (13)
C2—C3—Cl3	122.71 (14)	C12—C13—Cl13	120.68 (13)
C4—C3—Cl3	116.20 (13)	O14—C14—C15	123.05 (15)
O4—C4—C5	121.72 (17)	O14—C14—C13	117.98 (15)
O4—C4—C3	121.32 (17)	C15—C14—C13	118.96 (16)
C5—C4—C3	116.95 (15)	C14—O14—H14	109.5
C6—C5—C4	122.04 (16)	C16—C15—C14	121.10 (16)
C6—C5—H5	119.0	C16—C15—H15	119.4
C4—C5—H5	119.0	C14—C15—H15	119.4
C5—C6—C1	121.43 (17)	C15—C16—C11	120.63 (16)
C5—C6—H6	119.3	C15—C16—H16	119.7
C1—C6—H6	119.3	C11—C16—H16	119.7
O11—C11—C16	122.47 (15)	H20A—O20—H20B	108 (3)
O11—C11—C12	118.89 (15)		
O1—C1—C2—C3	179.07 (17)	O11—C11—C12—C13	179.97 (15)
C6—C1—C2—C3	-0.9 (2)	C16—C11—C12—C13	-0.1 (3)

supplementary materials

O1—C1—C2—C12	-0.3 (2)	O11—C11—C12—C112	-0.2 (2)
C6—C1—C2—C12	179.69 (13)	C16—C11—C12—C112	179.74 (13)
C1—C2—C3—C4	-1.5 (3)	C11—C12—C13—C14	0.2 (3)
C12—C2—C3—C4	177.83 (13)	C112—C12—C13—C14	-179.57 (13)
C1—C2—C3—C13	179.46 (13)	C11—C12—C13—C113	-179.82 (13)
C12—C2—C3—C13	-1.2 (2)	C112—C12—C13—C113	0.4 (2)
C2—C3—C4—O4	-175.28 (17)	C12—C13—C14—O14	179.13 (15)
C13—C3—C4—O4	3.8 (2)	C113—C13—C14—O14	-0.8 (2)
C2—C3—C4—C5	3.7 (2)	C12—C13—C14—C15	-0.4 (3)
C13—C3—C4—C5	-177.23 (13)	C113—C13—C14—C15	179.68 (13)
O4—C4—C5—C6	175.42 (18)	O14—C14—C15—C16	-179.12 (16)
C3—C4—C5—C6	-3.5 (3)	C13—C14—C15—C16	0.4 (3)
C4—C5—C6—C1	1.2 (3)	C14—C15—C16—C11	-0.2 (3)
O1—C1—C6—C5	-178.88 (17)	O11—C11—C16—C15	180.00 (16)
C2—C1—C6—C5	1.1 (3)	C12—C11—C16—C15	0.0 (3)

Hydrogen-bond geometry (\AA , $^\circ$)

<i>D</i> —H \cdots <i>A</i>	<i>D</i> —H	H \cdots <i>A</i>	<i>D</i> \cdots <i>A</i>	<i>D</i> —H \cdots <i>A</i>
O11—H11 \cdots O20	0.84	1.76	2.5947 (18)	173
O14—H14 \cdots O4 ⁱ	0.84	2.03	2.8381 (18)	161
O20—H20A \cdots O1 ⁱⁱ	0.79 (3)	2.12 (3)	2.914 (2)	177 (3)
O20—H20B \cdots O11 ⁱⁱⁱ	0.74 (3)	2.06 (3)	2.7899 (19)	169 (3)
C6—H6 \cdots O14 ^{iv}	0.95	2.48	3.209 (2)	134

Symmetry codes: (i) $x-1, y+1, z$; (ii) $-x+1, y-1/2, -z+1/2$; (iii) $-x+1, y+1/2, -z+1/2$; (iv) $-x+1, -y+2, -z+1$.

Fig. 1




Fig. 2




Fig. 3

