

Opipramol dihydrochloride

Richard Betz,^{a*} Thomas Gerber,^a Eric Hosten,^a
Budanoor P. Siddaraju^b and Hemmige S. Yathirajan^b

^aNelson Mandela Metropolitan University, Summerstrand Campus, Department of Chemistry, University Way, Summerstrand, PO Box 77000, Port Elizabeth 6031, South Africa, and ^bUniversity of Mysore, Department of Studies in Chemistry, Manasagangotri, Mysore 570 006, India

Correspondence e-mail: richard.betz@webmail.co.za


Received 27 September 2011; accepted 13 October 2011

Key indicators: single-crystal X-ray study; $T = 200$ K; mean $\sigma(\text{C}-\text{C}) = 0.002$ Å; R factor = 0.027; wR factor = 0.071; data-to-parameter ratio = 19.2.

The title compound (systematic name: 4-[3-[2-azatricyclo-[9.4.0.0^{3,8}]pentadeca-1(15),3,5,7,11,13-hexaen-2-yl]propyl]-1-(2-hydroxyethyl)piperazine-1,4-dium dichloride), $\text{C}_{23}\text{H}_{31}\text{N}_3\text{O}^+\cdot 2\text{Cl}^-$, is the dihydrochloride of a piperazine derivative bearing a bulky 3-(5*H*-dibenz[*b,f*]azepin-5-yl)propyl substituent. Protonation took place on both N atoms of the piperazine unit. The diazacyclohexane ring adopts a chair conformation. $\text{N}-\text{H}\cdots\text{Cl}$, $\text{O}-\text{H}\cdots\text{Cl}$ and $\text{C}-\text{H}\cdots\text{Cl}$ hydrogen bonding as well as $\text{C}-\text{H}\cdots\text{O}$ contacts connect the components into a three-dimensional network in the crystal. Two $\text{C}-\text{H}\cdots\pi$ contacts are also observed.

Related literature

For applications of opipramol, see: Moller *et al.* (2001). For related structures, see: Jasinski *et al.* (2010); Fun *et al.* (2011); Siddegowda, Butcher *et al.* (2011); Siddegowda, Jasinski *et al.* (2011); Swamy *et al.* (2007). For graph-set analysis of hydrogen bonds, see: Etter *et al.* (1990); Bernstein *et al.* (1995). For puckering analysis, see: Cremer & Pople (1975); Boessenkool & Boeyens (1980).



Experimental

Crystal data

$\text{C}_{23}\text{H}_{31}\text{N}_3\text{O}^{2+}\cdot 2\text{Cl}^-$

$M_r = 436.41$

Orthorhombic, $Pna2_1$

$a = 33.6581$ (6) Å

$b = 9.4265$ (2) Å

$c = 6.8978$ (1) Å

$V = 2188.52$ (7) Å³

$Z = 4$

Mo $K\alpha$ radiation

$\mu = 0.32$ mm⁻¹

$T = 200$ K

$0.51 \times 0.27 \times 0.14$ mm

Data collection

Bruker APEXII CCD

diffractometer

20068 measured reflections

5253 independent reflections

4854 reflections with $I > 2\sigma(I)$

$R_{\text{int}} = 0.030$

Refinement

$R[F^2 > 2\sigma(F^2)] = 0.027$

$wR(F^2) = 0.071$

$S = 1.03$

5253 reflections

274 parameters

1 restraint

H atoms treated by a mixture of independent and constrained refinement

$\Delta\rho_{\text{max}} = 0.24$ e Å⁻³

$\Delta\rho_{\text{min}} = -0.21$ e Å⁻³

Absolute structure: Flack (1983),

2293 Friedel pairs

Flack parameter: -0.004 (33)

Table 1

Hydrogen-bond geometry (Å, °).

Cg2 is the centroid of the C11–C16 ring.

$D-\text{H}\cdots A$	$D-\text{H}$	$\text{H}\cdots A$	$D\cdots A$	$D-\text{H}\cdots A$
O1–H81 \cdots Cl1 ⁱ	0.79 (2)	2.39 (2)	3.1701 (14)	169 (2)
N2–H72 \cdots Cl1	0.82 (2)	2.212 (19)	3.0057 (13)	163.0 (15)
N3–H73 \cdots Cl2 ⁱⁱ	0.98 (3)	2.03 (3)	2.9972 (12)	171 (2)
C5–H5A \cdots Cl1 ⁱⁱⁱ	0.99	2.82	3.7065 (15)	149
C5–H5A \cdots O1 ⁱⁱ	0.99	2.59	3.3373 (18)	133
C14–H14 \cdots Cl1 ^{iv}	0.95	2.83	3.7233 (14)	157
C31–H31A \cdots O1 ⁱⁱ	0.99	2.54	3.2446 (18)	128
C31–H31B \cdots Cl2 ^v	0.99	2.75	3.5518 (14)	139
C32–H32B \cdots Cl1 ⁱⁱ	0.99	2.85	3.8246 (13)	169
C32–H32A \cdots Cl2 ^{vi}	0.99	2.85	3.7213 (16)	148
C6–H6B \cdots Cl2 ^{vi}	0.99	2.76	3.6634 (18)	152
C34–H34A \cdots Cl2	0.99	2.82	3.5471 (13)	131
C16–H16 \cdots Cg2 ^{vii}	0.95	2.98	3.6402 (16)	128
C23–H23 \cdots Cg2 ^{vii}	0.95	2.67	3.4805 (18)	143

Symmetry codes: (i) $-x + \frac{1}{2}, y - \frac{1}{2}, z + \frac{1}{2}$; (ii) $-x + \frac{1}{2}, y + \frac{1}{2}, z + \frac{1}{2}$; (iii) $x, y, z + 1$; (iv) $-x, -y + 1, z + \frac{1}{2}$; (v) $x, y + 1, z$; (vi) $-x + \frac{1}{2}, y + \frac{1}{2}, z - \frac{1}{2}$; (vii) $-x, -y + 2, z - \frac{1}{2}$.

Data collection: APEX2 (Bruker, 2010); cell refinement: SAINT (Bruker, 2010); data reduction: SAINT; program(s) used to solve structure: SHELXS97 (Sheldrick, 2008); program(s) used to refine structure: SHELXL97 (Sheldrick, 2008); molecular graphics: ORTEP-3 (Farrugia, 1997) and Mercury (Macrae *et al.*, 2008); software used to prepare material for publication: SHELXL97 and PLATON (Spek, 2009).

BPS thanks the University of Mysore for research facilities. HSY thanks R. L. Fine Chem., Bengaluru, for the gift sample of the title compound.

Supplementary data and figures for this paper are available from the IUCr electronic archives (Reference: MW2030).

References

Bernstein, J., Davis, R. E., Shimon, L. & Chang, N.-L. (1995). *Angew. Chem. Int. Ed. Engl.* **34**, 1555–1573.

- Boessenkool, I. K. & Boeyens, J. C. A. (1980). *J. Cryst. Mol. Struct.* **10**, 11–18.
- Bruker (2010). *APEX2* and *SAINT* Bruker AXS Inc., Madison, Wisconsin, USA.
- Cremer, D. & Pople, J. A. (1975). *J. Am. Chem. Soc.* **97**, 1354–1358.
- Etter, M. C., MacDonald, J. C. & Bernstein, J. (1990). *Acta Cryst.* **B46**, 256–262.
- Farrugia, L. J. (1997). *J. Appl. Cryst.* **30**, 565.
- Flack, H. D. (1983). *Acta Cryst.* **A39**, 876–881.
- Fun, H.-K., Loh, W.-S., Siddegowda, M. S., Yathirajan, H. S. & Narayana, B. (2011). *Acta Cryst.* **E67**, o1598.
- Jasinski, J. P., Pek, A. E., Siddaraju, B. P., Yathirajan, H. S. & Narayana, B. (2010). *Acta Cryst.* **E66**, o1979–o1980.
- Macrae, C. F., Bruno, I. J., Chisholm, J. A., Edgington, P. R., McCabe, P., Pidcock, E., Rodriguez-Monge, L., Taylor, R., van de Streek, J. & Wood, P. A. (2008). *J. Appl. Cryst.* **41**, 466–470.
- Moller, H. J., Volz, H. P., Reimann, I. W. & Stoll, K. D. (2001). *J. Clin. Psychopharmacol.* **21**, 59–65.
- Sheldrick, G. M. (2008). *Acta Cryst.* **A64**, 112–122.
- Siddegowda, M. S., Butcher, R. J., Akkurt, M., Yathirajan, H. S. & Narayana, B. (2011). *Acta Cryst.* **E67**, o2079–o2080.
- Siddegowda, M. S., Jasinski, J. P., Golen, J. A., Yathirajan, H. S. & Swamy, M. T. (2011). *Acta Cryst.* **E67**, o2296.
- Spek, A. L. (2009). *Acta Cryst.* **D65**, 148–155.
- Swamy, M. T., Ashok, M. A., Yathirajan, H. S., Narayana, B. & Bolte, M. (2007). *Acta Cryst.* **E63**, o4919.

supplementary materials

Acta Cryst. (2011). E67, o2994–o2995 [doi:10.1107/S1600536811042280]

Opipramol dihydrochloride

R. Betz, T. Gerber, E. Hosten, B. P. Siddaraju and H. S. Yathirajan

Comment

Opipramol (systematic IUPAC name: 4-(3-{2-azatricyclo[9.4.0.0^{3,8}]pentadeca- 1(15),3,5,7,11,13-hexaen-2-yl}propyl)-1-(2-hydroxyethyl)piperazin-1,4-dium dichloride is an antidepressant and anxiolytic typically used in the treatment of generalized anxiety disorder (Moller *et al.*, 2001). Opipramol is a tricyclic compound with no reuptake-inhibiting properties. However, it has pronounced D2-, 5-HT2-, and H1-blocking potential and high affinity to sigma receptors (sigma-1 and sigma-2). The crystal structures of opipramol dipicrate (Jasinski *et al.*, 2010), opipramol (Fun *et al.*, 2011), opipramolium fumarate (Siddegowda, Butcher *et al.*, 2011) and flupentixol dihydrochloride (Siddegowda, Jasinski *et al.*, 2011) have been reported recently. In view of the importance of the title compound we herein report its molecular and crystal structure.

Protonation took place exclusively on both nitrogen atoms of the piperazine unit. The diazacyclohexane ring adopts a ⁴C₁ (^N₃C_N₂) conformation (Cremer & Pople, 1975). The more rigid skeleton of the seven-membered ring adopts a conformation in between a TC2 and a C6 conformation (Q_2 : 0.6128 (14) Å, Q_3 : 0.2043 (14) Å, π_2 : 177.14 (14)°, π_3 : 178.9 (4)°) (Boessenkool & Boeyens, 1980). The nitrogen atom sticks out of the least-squares plane defined by its atoms by 0.451 (1) Å (Fig. 1).

Apart from classical hydrogen bonds of the N–H⋯Cl and the O–H⋯Cl type, the crystal structure features a multitude of C–H⋯Cl contacts and two C–H⋯O contacts whose ranges invariably fall at least 0.1 Å below the sum of van-der-Waals radii of the respective atoms participating. The C–H⋯Cl contacts involve hydrogen atoms of methylene groups – both intracyclic as well as those on the hydroxyethyl side-chain – as well as one hydrogen atom on a phenyl ring. The C–H⋯O contacts all involve hydrogen atoms of the methylene groups located in the piperazine moiety and in the hydrocarbon chain connecting this to the remainder of the molecule. Both chloride anions attain pentacoordination *via* their combined contacts with hydrogen atoms, however, one of the anions features four instead of three C–H-supported contacts. In terms of graph-set analysis (Etter *et al.*, 1990; Bernstein *et al.*, 1995), the descriptor for the classical hydrogen bonds is *DDD* on the unitary level while the C–H⋯O contacts require a *C*¹₁(7)*C*¹₁(9) descriptor on the same level. A *DDDDDDD* descriptor on the unitary level is necessary to capture the C–H⋯Cl contacts. Furthermore, two C–H⋯C_g contacts can be observed involving hydrogen atoms on the phenyl rings as donors and the phenyl moiety comprised of the carbon atoms C11–C16. All of these interactions serve to connect the cations and anions into a three-dimensional network in the crystal. The shortest intercentroid distance between two aromatic systems was measured at 4.7319 (9) Å and is apparent between two different phenyl moieties.

The packing of the title compound in the crystal is shown in Figure 2.

Experimental

The title compound was obtained as a gift sample from *R. L. Fine Chem. Ltd.*, Bangalore, India. The compound was recrystallized from a 1:1 mixture of butan-1-one and benzene.

Refinement

Carbon-bound H atoms were placed in calculated positions (C—H 0.95 Å for aromatic and vinylic hydrogen atoms, C—H 0.99 Å for methylene groups) and were included in the refinement in the riding model approximation, with $U(\text{H})$ set to $1.2U_{\text{eq}}(\text{C})$. The oxygen-bound H atom as well as both nitrogen-bound H atoms were located on a difference Fourier map and refined freely.

Figures




Fig. 1. The molecular structure of the title compound, with atom labels and anisotropic displacement ellipsoids (drawn at 50% probability level).




Fig. 2. Molecular packing of the title compound, viewed along [0 0 - 1] (anisotropic displacement ellipsoids drawn at 50% probability level).

4-{3-[2-azatricyclo[9.4.0.0^{3,8}]pentadeca-1(15),3,5,7,11,13-\ hexaen-2-yl]propyl}-1-(2-hydroxyethyl)piperazine-1,4-dium dichloride

Crystal data

$\text{C}_{23}\text{H}_{31}\text{N}_3\text{O}^{2+}\cdot 2\text{Cl}^-$

$M_r = 436.41$

Orthorhombic, $Pna2_1$

Hall symbol: $P\ 2c\ -2n$

$a = 33.6581\ (6)\ \text{\AA}$

$b = 9.4265\ (2)\ \text{\AA}$

$c = 6.8978\ (1)\ \text{\AA}$

$D_x = 1.324\ \text{Mg m}^{-3}$

Melting point = 482–484 K

Mo $K\alpha$ radiation, $\lambda = 0.71073\ \text{\AA}$

Cell parameters from 9954 reflections

$\theta = 2.4\text{--}28.3^\circ$

$\mu = 0.32\ \text{mm}^{-1}$

$T = 200\ \text{K}$

$V = 2188.52 (7) \text{ \AA}^3$

$Z = 4$

$F(000) = 928$

Platelet, green

$0.51 \times 0.27 \times 0.14 \text{ mm}$

Data collection

Bruker APEXII CCD
diffractometer

4854 reflections with $I > 2\sigma(I)$

Radiation source: fine-focus sealed tube

$R_{\text{int}} = 0.030$

graphite

$\theta_{\text{max}} = 28.3^\circ$, $\theta_{\text{min}} = 2.2^\circ$

φ and ω scans

$h = -44 \rightarrow 44$

20068 measured reflections

$k = -12 \rightarrow 9$

5253 independent reflections

$l = -9 \rightarrow 9$

Refinement

Refinement on F^2

Secondary atom site location: difference Fourier map

Least-squares matrix: full

Hydrogen site location: inferred from neighbouring sites

$R[F^2 > 2\sigma(F^2)] = 0.027$

H atoms treated by a mixture of independent and constrained refinement

$wR(F^2) = 0.071$

$w = 1/[\sigma^2(F_o^2) + (0.0469P)^2 + 0.0541P]$

where $P = (F_o^2 + 2F_c^2)/3$

$S = 1.03$

$(\Delta/\sigma)_{\text{max}} = 0.001$

5253 reflections

$\Delta\rho_{\text{max}} = 0.24 \text{ e \AA}^{-3}$

274 parameters

$\Delta\rho_{\text{min}} = -0.21 \text{ e \AA}^{-3}$

1 restraint

Absolute structure: Flack (1983), **2293 Friedel pairs**

Primary atom site location: structure-invariant direct methods

Flack parameter: $-0.004 (33)$

Fractional atomic coordinates and isotropic or equivalent isotropic displacement parameters (\AA^2)

	<i>x</i>	<i>y</i>	<i>z</i>	$U_{\text{iso}}^*/U_{\text{eq}}$
O1	0.33522 (3)	0.24837 (14)	0.58052 (17)	0.0359 (3)
H81	0.3356 (6)	0.173 (2)	0.631 (4)	0.044 (6)*
N1	0.05595 (3)	0.73088 (11)	0.60732 (17)	0.0210 (2)
N2	0.19011 (3)	0.51107 (11)	0.64833 (18)	0.0161 (2)
H72	0.1886 (5)	0.5036 (16)	0.530 (3)	0.020 (4)*
N3	0.27628 (3)	0.49616 (11)	0.69361 (16)	0.0174 (2)
H73	0.2750 (7)	0.500 (2)	0.835 (5)	0.074 (8)*
C1	0.00179 (4)	0.79746 (17)	0.2855 (2)	0.0294 (3)
H1	-0.0128	0.7742	0.1721	0.035*
C2	0.03133 (4)	0.89067 (17)	0.2622 (2)	0.0310 (3)
H2	0.0356	0.9237	0.1337	0.037*
C3	0.08286 (3)	0.64016 (15)	0.7199 (2)	0.0213 (3)
H3A	0.0719	0.5428	0.7253	0.026*
H3B	0.0846	0.6765	0.8543	0.026*
C4	0.12436 (4)	0.63572 (14)	0.6312 (2)	0.0239 (3)

supplementary materials

H4A	0.1224	0.6198	0.4896	0.029*
H4B	0.1379	0.7275	0.6529	0.029*
C5	0.14816 (3)	0.51695 (13)	0.7235 (2)	0.0186 (2)
H5A	0.1487	0.5309	0.8657	0.022*
H5B	0.1349	0.4253	0.6970	0.022*
C6	0.31889 (4)	0.49360 (15)	0.6266 (3)	0.0237 (3)
H6A	0.3322	0.5816	0.6702	0.028*
H6B	0.3195	0.4921	0.4832	0.028*
C7	0.34174 (4)	0.36661 (17)	0.7029 (2)	0.0285 (3)
H7A	0.3705	0.3890	0.7074	0.034*
H7B	0.3328	0.3441	0.8362	0.034*
C11	0.01455 (3)	0.69862 (13)	0.6216 (2)	0.0207 (2)
C12	-0.01079 (4)	0.72710 (15)	0.4633 (2)	0.0232 (3)
C13	-0.05001 (4)	0.67850 (16)	0.4742 (2)	0.0292 (3)
H13	-0.0675	0.6980	0.3694	0.035*
C14	-0.06431 (4)	0.60348 (16)	0.6307 (3)	0.0332 (3)
H14	-0.0910	0.5707	0.6328	0.040*
C15	-0.03922 (4)	0.57682 (16)	0.7842 (3)	0.0332 (4)
H15	-0.0485	0.5242	0.8924	0.040*
C16	-0.00025 (4)	0.62655 (16)	0.7816 (2)	0.0272 (3)
H16	0.0164	0.6110	0.8908	0.033*
C21	0.06763 (4)	0.87575 (14)	0.5829 (2)	0.0219 (3)
C22	0.05776 (4)	0.94775 (16)	0.4107 (2)	0.0250 (3)
C23	0.07445 (5)	1.08179 (16)	0.3787 (3)	0.0338 (4)
H23	0.0679	1.1315	0.2633	0.041*
C24	0.10005 (5)	1.14375 (17)	0.5089 (3)	0.0400 (4)
H24	0.1118	1.2331	0.4808	0.048*
C25	0.10856 (5)	1.07525 (18)	0.6806 (3)	0.0383 (4)
H25	0.1256	1.1186	0.7728	0.046*
C26	0.09204 (4)	0.94189 (17)	0.7189 (3)	0.0294 (3)
H26	0.0975	0.8961	0.8386	0.035*
C31	0.21421 (3)	0.63744 (14)	0.7062 (2)	0.0199 (3)
H31A	0.2154	0.6432	0.8494	0.024*
H31B	0.2012	0.7248	0.6578	0.024*
C32	0.25605 (3)	0.62810 (13)	0.6249 (2)	0.0199 (2)
H32A	0.2549	0.6281	0.4815	0.024*
H32B	0.2715	0.7121	0.6666	0.024*
C33	0.25223 (3)	0.37093 (13)	0.6327 (2)	0.0190 (2)
H33A	0.2653	0.2829	0.6782	0.023*
H33B	0.2508	0.3674	0.4895	0.023*
C34	0.21058 (4)	0.37928 (14)	0.7157 (2)	0.0191 (2)
H34A	0.1952	0.2952	0.6742	0.023*
H34B	0.2119	0.3790	0.8591	0.023*
Cl1	0.172378 (10)	0.43277 (4)	0.23488 (5)	0.03102 (9)
Cl2	0.217571 (10)	0.00951 (3)	0.62670 (6)	0.02579 (8)

Atomic displacement parameters (\AA^2)

	U^{11}	U^{22}	U^{33}	U^{12}	U^{13}	U^{23}
O1	0.0419 (6)	0.0321 (6)	0.0337 (7)	0.0108 (5)	-0.0089 (5)	-0.0078 (5)
N1	0.0168 (4)	0.0210 (5)	0.0252 (6)	0.0029 (4)	-0.0012 (4)	0.0058 (5)
N2	0.0176 (5)	0.0147 (5)	0.0161 (6)	0.0006 (4)	-0.0018 (4)	0.0003 (4)
N3	0.0177 (5)	0.0200 (6)	0.0146 (5)	0.0015 (4)	-0.0007 (4)	-0.0010 (4)
C1	0.0297 (7)	0.0346 (8)	0.0240 (7)	0.0056 (6)	-0.0057 (5)	-0.0003 (6)
C2	0.0354 (7)	0.0339 (8)	0.0237 (8)	0.0069 (6)	0.0000 (6)	0.0072 (6)
C3	0.0179 (5)	0.0240 (6)	0.0219 (6)	0.0043 (5)	0.0001 (5)	0.0045 (6)
C4	0.0197 (6)	0.0267 (7)	0.0253 (7)	0.0060 (5)	0.0020 (5)	0.0062 (6)
C5	0.0150 (5)	0.0198 (6)	0.0209 (6)	0.0013 (4)	-0.0003 (5)	0.0007 (6)
C6	0.0164 (5)	0.0307 (7)	0.0241 (6)	-0.0005 (5)	0.0006 (6)	-0.0026 (6)
C7	0.0212 (6)	0.0377 (8)	0.0268 (7)	0.0080 (6)	-0.0065 (5)	-0.0059 (6)
C11	0.0181 (5)	0.0179 (6)	0.0262 (6)	0.0036 (4)	0.0004 (5)	-0.0018 (6)
C12	0.0224 (6)	0.0200 (7)	0.0272 (7)	0.0040 (5)	-0.0030 (5)	-0.0045 (6)
C13	0.0223 (7)	0.0274 (7)	0.0378 (8)	0.0037 (6)	-0.0064 (6)	-0.0095 (7)
C14	0.0221 (6)	0.0259 (7)	0.0517 (10)	-0.0012 (5)	0.0044 (7)	-0.0092 (8)
C15	0.0259 (7)	0.0245 (7)	0.0492 (11)	0.0017 (6)	0.0117 (6)	0.0049 (7)
C16	0.0229 (6)	0.0276 (7)	0.0312 (8)	0.0053 (6)	0.0038 (5)	0.0044 (6)
C21	0.0166 (6)	0.0210 (6)	0.0283 (8)	0.0044 (5)	0.0031 (5)	0.0019 (6)
C22	0.0228 (6)	0.0242 (7)	0.0280 (7)	0.0052 (6)	0.0045 (6)	0.0045 (6)
C23	0.0374 (8)	0.0238 (8)	0.0403 (9)	0.0067 (6)	0.0072 (7)	0.0092 (7)
C24	0.0353 (8)	0.0211 (8)	0.0637 (12)	-0.0021 (7)	0.0059 (8)	0.0027 (8)
C25	0.0284 (7)	0.0265 (8)	0.0598 (13)	0.0009 (6)	-0.0061 (7)	-0.0085 (7)
C26	0.0258 (6)	0.0265 (7)	0.0359 (8)	0.0040 (6)	-0.0024 (6)	0.0002 (7)
C31	0.0190 (5)	0.0152 (6)	0.0256 (7)	0.0010 (5)	-0.0004 (5)	-0.0032 (5)
C32	0.0218 (6)	0.0159 (5)	0.0220 (6)	-0.0002 (4)	-0.0004 (5)	0.0013 (6)
C33	0.0190 (5)	0.0161 (5)	0.0218 (6)	0.0023 (4)	-0.0001 (5)	-0.0019 (6)
C34	0.0204 (5)	0.0146 (6)	0.0224 (6)	0.0019 (5)	-0.0002 (5)	0.0026 (6)
Cl1	0.02975 (16)	0.0431 (2)	0.02020 (15)	0.00569 (15)	-0.00392 (14)	-0.00588 (17)
Cl2	0.03646 (17)	0.02445 (16)	0.01645 (14)	-0.00133 (13)	0.00162 (14)	-0.00090 (14)

Geometric parameters (\AA , $^\circ$)

O1—C7	1.4155 (19)	C11—C16	1.388 (2)
O1—H81	0.79 (2)	C11—C12	1.4112 (19)
N1—C11	1.4296 (15)	C12—C13	1.3995 (19)
N1—C21	1.4311 (17)	C13—C14	1.377 (2)
N1—C3	1.4681 (16)	C13—H13	0.9500
N2—C34	1.4947 (16)	C14—C15	1.377 (2)
N2—C31	1.4955 (16)	C14—H14	0.9500
N2—C5	1.5049 (16)	C15—C16	1.393 (2)
N2—H72	0.82 (2)	C15—H15	0.9500
N3—C33	1.4918 (16)	C16—H16	0.9500
N3—C32	1.4950 (16)	C21—C26	1.394 (2)
N3—C6	1.5069 (16)	C21—C22	1.408 (2)
N3—H73	0.98 (3)	C22—C23	1.400 (2)

supplementary materials

C1—C2	1.336 (2)	C23—C24	1.375 (3)
C1—C12	1.457 (2)	C23—H23	0.9500
C1—H1	0.9500	C24—C25	1.379 (3)
C2—C22	1.460 (2)	C24—H24	0.9500
C2—H2	0.9500	C25—C26	1.400 (2)
C3—C4	1.5256 (17)	C25—H25	0.9500
C3—H3A	0.9900	C26—H26	0.9500
C3—H3B	0.9900	C31—C32	1.5183 (17)
C4—C5	1.5168 (18)	C31—H31A	0.9900
C4—H4A	0.9900	C31—H31B	0.9900
C4—H4B	0.9900	C32—H32A	0.9900
C5—H5A	0.9900	C32—H32B	0.9900
C5—H5B	0.9900	C33—C34	1.5160 (17)
C6—C7	1.517 (2)	C33—H33A	0.9900
C6—H6A	0.9900	C33—H33B	0.9900
C6—H6B	0.9900	C34—H34A	0.9900
C7—H7A	0.9900	C34—H34B	0.9900
C7—H7B	0.9900		
C7—O1—H81	116.3 (18)	C13—C12—C1	117.91 (13)
C11—N1—C21	118.62 (10)	C11—C12—C1	124.22 (13)
C11—N1—C3	116.17 (10)	C14—C13—C12	122.66 (14)
C21—N1—C3	116.65 (10)	C14—C13—H13	118.7
C34—N2—C31	109.20 (10)	C12—C13—H13	118.7
C34—N2—C5	110.86 (10)	C15—C14—C13	118.80 (13)
C31—N2—C5	112.80 (10)	C15—C14—H14	120.6
C34—N2—H72	105.4 (11)	C13—C14—H14	120.6
C31—N2—H72	111.5 (11)	C14—C15—C16	120.43 (15)
C5—N2—H72	106.8 (11)	C14—C15—H15	119.8
C33—N3—C32	108.78 (10)	C16—C15—H15	119.8
C33—N3—C6	114.68 (10)	C11—C16—C15	120.82 (14)
C32—N3—C6	110.47 (10)	C11—C16—H16	119.6
C33—N3—H73	106.9 (13)	C15—C16—H16	119.6
C32—N3—H73	105.3 (13)	C26—C21—C22	119.42 (13)
C6—N3—H73	110.3 (13)	C26—C21—N1	120.63 (12)
C2—C1—C12	128.07 (14)	C22—C21—N1	119.64 (13)
C2—C1—H1	116.0	C23—C22—C21	118.25 (14)
C12—C1—H1	116.0	C23—C22—C2	117.80 (14)
C1—C2—C22	127.67 (14)	C21—C22—C2	123.94 (14)
C1—C2—H2	116.2	C24—C23—C22	122.13 (15)
C22—C2—H2	116.2	C24—C23—H23	118.9
N1—C3—C4	111.63 (11)	C22—C23—H23	118.9
N1—C3—H3A	109.3	C23—C24—C25	119.48 (15)
C4—C3—H3A	109.3	C23—C24—H24	120.3
N1—C3—H3B	109.3	C25—C24—H24	120.3
C4—C3—H3B	109.3	C24—C25—C26	120.03 (16)
H3A—C3—H3B	108.0	C24—C25—H25	120.0
C5—C4—C3	109.60 (11)	C26—C25—H25	120.0
C5—C4—H4A	109.8	C21—C26—C25	120.56 (16)
C3—C4—H4A	109.8	C21—C26—H26	119.7

C5—C4—H4B	109.8	C25—C26—H26	119.7
C3—C4—H4B	109.8	N2—C31—C32	111.00 (11)
H4A—C4—H4B	108.2	N2—C31—H31A	109.4
N2—C5—C4	112.22 (11)	C32—C31—H31A	109.4
N2—C5—H5A	109.2	N2—C31—H31B	109.4
C4—C5—H5A	109.2	C32—C31—H31B	109.4
N2—C5—H5B	109.2	H31A—C31—H31B	108.0
C4—C5—H5B	109.2	N3—C32—C31	110.71 (11)
H5A—C5—H5B	107.9	N3—C32—H32A	109.5
N3—C6—C7	112.88 (12)	C31—C32—H32A	109.5
N3—C6—H6A	109.0	N3—C32—H32B	109.5
C7—C6—H6A	109.0	C31—C32—H32B	109.5
N3—C6—H6B	109.0	H32A—C32—H32B	108.1
C7—C6—H6B	109.0	N3—C33—C34	110.76 (10)
H6A—C6—H6B	107.8	N3—C33—H33A	109.5
O1—C7—C6	109.62 (12)	C34—C33—H33A	109.5
O1—C7—H7A	109.7	N3—C33—H33B	109.5
C6—C7—H7A	109.7	C34—C33—H33B	109.5
O1—C7—H7B	109.7	H33A—C33—H33B	108.1
C6—C7—H7B	109.7	N2—C34—C33	110.62 (10)
H7A—C7—H7B	108.2	N2—C34—H34A	109.5
C16—C11—C12	119.44 (12)	C33—C34—H34A	109.5
C16—C11—N1	120.57 (12)	N2—C34—H34B	109.5
C12—C11—N1	119.67 (12)	C33—C34—H34B	109.5
C13—C12—C11	117.80 (13)	H34A—C34—H34B	108.1
C12—C1—C2—C22	-1.6 (3)	C11—N1—C21—C26	120.11 (14)
C11—N1—C3—C4	154.28 (12)	C3—N1—C21—C26	-26.61 (18)
C21—N1—C3—C4	-58.17 (16)	C11—N1—C21—C22	-66.34 (17)
N1—C3—C4—C5	-167.31 (11)	C3—N1—C21—C22	146.94 (12)
C34—N2—C5—C4	-170.35 (11)	C26—C21—C22—C23	2.8 (2)
C31—N2—C5—C4	66.82 (15)	N1—C21—C22—C23	-170.80 (12)
C3—C4—C5—N2	-177.44 (11)	C26—C21—C22—C2	-177.67 (13)
C33—N3—C6—C7	-60.37 (17)	N1—C21—C22—C2	8.7 (2)
C32—N3—C6—C7	176.29 (12)	C1—C2—C22—C23	-153.31 (16)
N3—C6—C7—O1	84.54 (16)	C1—C2—C22—C21	27.2 (2)
C21—N1—C11—C16	-123.35 (14)	C21—C22—C23—C24	0.3 (2)
C3—N1—C11—C16	23.53 (18)	C2—C22—C23—C24	-179.21 (15)
C21—N1—C11—C12	63.22 (17)	C22—C23—C24—C25	-2.6 (3)
C3—N1—C11—C12	-149.90 (12)	C23—C24—C25—C26	1.7 (3)
C16—C11—C12—C13	-1.05 (19)	C22—C21—C26—C25	-3.7 (2)
N1—C11—C12—C13	172.46 (13)	N1—C21—C26—C25	169.85 (13)
C16—C11—C12—C1	-177.82 (14)	C24—C25—C26—C21	1.4 (2)
N1—C11—C12—C1	-4.3 (2)	C34—N2—C31—C32	57.10 (15)
C2—C1—C12—C13	155.85 (16)	C5—N2—C31—C32	-179.15 (11)
C2—C1—C12—C11	-27.4 (2)	C33—N3—C32—C31	58.15 (15)
C11—C12—C13—C14	-0.8 (2)	C6—N3—C32—C31	-175.15 (12)
C1—C12—C13—C14	176.16 (14)	N2—C31—C32—N3	-58.39 (15)
C12—C13—C14—C15	1.0 (2)	C32—N3—C33—C34	-58.79 (15)
C13—C14—C15—C16	0.8 (2)	C6—N3—C33—C34	176.97 (12)

supplementary materials

C12—C11—C16—C15	2.8 (2)	C31—N2—C34—C33	-57.48 (14)
N1—C11—C16—C15	-170.66 (13)	C5—N2—C34—C33	177.64 (11)
C14—C15—C16—C11	-2.7 (2)	N3—C33—C34—N2	59.46 (15)

Hydrogen-bond geometry (\AA , $^\circ$)

Cg2 is the centroid of the C11—C16 ring.

$D-H\cdots A$	$D-H$	$H\cdots A$	$D\cdots A$	$D-H\cdots A$
O1—H81 \cdots C11 ⁱ	0.79 (2)	2.39 (2)	3.1701 (14)	169 (2)
N2—H72 \cdots C11	0.82 (2)	2.212 (19)	3.0057 (13)	163.0 (15)
N3—H73 \cdots C12 ⁱⁱ	0.98 (3)	2.03 (3)	2.9972 (12)	171 (2)
C5—H5A \cdots C11 ⁱⁱⁱ	0.99	2.82	3.7065 (15)	149.
C5—H5A \cdots O1 ⁱⁱ	0.99	2.59	3.3373 (18)	133.
C14—H14 \cdots C11 ^{iv}	0.95	2.83	3.7233 (14)	157.
C31—H31A \cdots O1 ⁱⁱ	0.99	2.54	3.2446 (18)	128.
C31—H31B \cdots C12 ^v	0.99	2.75	3.5518 (14)	139.
C32—H32B \cdots C11 ⁱⁱ	0.99	2.85	3.8246 (13)	169.
C32—H32A \cdots C12 ^{vi}	0.99	2.85	3.7213 (16)	148.
C6—H6B \cdots C12 ^{vi}	0.99	2.76	3.6634 (18)	152.
C34—H34A \cdots C12	0.99	2.82	3.5471 (13)	131.
C16—H16 \cdots Cg2 ^{iv}	0.95	2.98	3.6402 (16)	128
C23—H23 \cdots Cg2 ^{vii}	0.95	2.67	3.4805 (18)	143

Symmetry codes: (i) $-x+1/2, y-1/2, z+1/2$; (ii) $-x+1/2, y+1/2, z+1/2$; (iii) $x, y, z+1$; (iv) $-x, -y+1, z+1/2$; (v) $x, y+1, z$; (vi) $-x+1/2, y+1/2, z-1/2$; (vii) $-x, -y+2, z-1/2$.

Fig. 1




Fig. 2

